

One year of ^{222}Rn concentration in the atmospheric surface layer

S. Galmarini

European Commission – DG Joint Research Centre, Institute for Environment and Sustainability, Ispra, Italy

Received: 22 July 2005 – Published in Atmos. Chem. Phys. Discuss.: 19 December 2005

Revised: 20 June 2006 – Accepted: 21 June 2006 – Published: 12 July 2006

Abstract. A one-year time series of ^{222}Rn measured in a rural area in the North of Italy in 1997 is analyzed. The scope of the investigation is to better understand the behavior of this common atmospheric tracer in relation to the meteorological conditions at the release site. Wavelet analysis is used as one of the investigation tools of the time series. The measurements and scalograms of ^{222}Rn are compared to those of wind-speed, pressure, relative humidity, temperature and NO_x . The use of wavelet analysis allows the identification of the various scales controlling the influence of the meteorological variables on ^{222}Rn dispersion in the surface layer that are not visible through classical Fourier analysis or direct time series inspection. The analysis of the time series has identified specific periods during which the usual diurnal variation of radon is superimposed to a linear growth thus indicating the build up of concentration at the measurement level. From these specific cases an estimate of the surface flux of ^{222}Rn is made. By means of a simple model these special cases are reproduced.

1 Introduction

Radon emission from the ground has long been studied over the last decades from several view points. As radioactive product of the uranium chain, it is abundant all over the Earth crust and constantly emitted from the ground. The noble gas chemical form makes radon invulnerable to wet deposition or chemical reaction thus allowing an undisturbed transport in the atmosphere as well as closed environments such as households or working places. Radioactive decay is the only removal process. We will not discuss here the potential health threat of radon as it has been debated over the years and it is still being debated intensively. Apart from this as-

pect, which is in any case very relevant, radon represents a very interesting natural tracer for a number of atmospheric research studies. It has long been studied in the atmospheric boundary layer (ABL) as constantly emitted from the surface (e.g. Israël et al., 1966; Ikebe and Shimo, 1972; Beck et al., 1979; Vinod Kumar et al., 1999; Sesana et al., 2003, 2005; Pearson and Moses, 1966; Marcazan et al., 1993, 1997; Kataoka et al., 2001, 2003), to characterize the turbulent diffusion properties of the lower atmospheric levels, and since some of its radio nuclides have decay timescales comparable to that of turbulent transport in the ABL (e.g. Beran and Assaf, 1970; Druilhet and Fontan, 1973; Kristensen et al., 1997). Similarly intensive research has been conducted at global scale where radon is used for global atmospheric chemistry models evaluation (e.g. Dentener et al., 1999), as well as for the estimate of the fluxes of the atmospheric constituents or pollutants. For a detailed review on the radon studies and sampling techniques in atmospheric science refer to Zahorowski et al. (2004).

In this study a one-year time series of radon-222 measured in the atmospheric surface layer (ASL) is analyzed. The data were collected in 1997 on the premises of the European Commission Joint Research Center within a collaboration project between the latter and the Faculty of Physics of the University of Milan (Facchini et al., 1999). Although the time series is obtained as 1-h average concentration at a fixed location and height, its analysis highlights a series of features that were not investigated before.

In the first part of this study we will focus on the relationship between the time evolution of radon and that of the meteorological parameters. Although the influence of the latter on radon emission and time evolution is rather well known, in this context we will focus on the relationship between the scales of occurrence of the various meteorological events and those of radon dispersion in the ASL. Towards this purpose, a time series analysis is conducted by means of wavelets (Farge, 2000; Farge and Schneider, 2002). Wavelet analysis

Correspondence to: S. Galmarini
(stefano.galmarini@jrc.it)

Table 1. Variable measured together with ^{222}Rn and correlation of all variables to radon concentration from the hourly data series.

	Variable	Correlation with radon
1	Wind speed	-0.457984
2	Wind direction	-
3	Temperature	-0.477703
4	Relative Humidity	0.634650
5	Precipitation	-0.0610427
6	Pressure	0.258573
7	NO_x	0.416235
8	Radon	1

is a technique for the determination of localized (in time or in space) power spectra which adapts very well to the analysis of signals with a strong time or space variability. Several studies in the past have clearly demonstrated the capacity of this technique to highlight fundamental features of a signal that would not be otherwise distinguished by classical time series analysis or Fourier transformation (e.g. Druhilet et al., 1994; Gao et al., 1993; Galmarini and Attié, 1999; Attié and Durand, 2003; Salmond, 2005). In the second part of the paper some specific features identified in the time series will be analyzed and modeled. In general the following research issues are addressed:

- Analysis of the one-year time series and investigation of the monthly variability of radon concentration;
- Dependence of hourly concentration at the surface on meteorological parameters;
- Comparison between radon concentration and the one of other atmospheric tracers;
- Analysis of specific peculiarities in the radon time series and provision of explanations, including the determination of the surface flux from mean concentration values.

2 Description of the data

The data analyzed consist of one year of continuous measurements with one hour resolution of the variables of Table 1. The sampling location is within the Joint Research Center (JRC) premises and corresponds to an official EMEP site (station IT04 (45°48' N; 8°38' E) – 209 m a.s.l.). The JRC is located in the Po Valley at the foothills of the western Alps on the eastern shore of Lago Maggiore. At the station, variables 1–7 are regularly measured together with a variety of other chemical compounds (Leyendecker et al., 1998). The radon measurements were conducted specifically over the period December 1996–January 1998, and are not

part of the ordinary station instrumentation. The radon sampling was performed by means of a monitor developed by the University of Milan and described in detail in Facchini et al. (1999) and Sesana et al. (2003). The instrument detects directly ^{222}Rn that is therefore not inferred from the analysis of its daughters as in other cases (Marcazzan et al., 1993, 1997). Air was sampled at 3 m from the surface. The time series from 1 January to 31 December 1997 is complete, except for 4 days in July (22–24 July 1997, 28–29 July 1997), 5 days in November (13–18 November 1997) during which no radon concentrations were collected. Occasionally measurements were not collected over time intervals of one hour. In this case the values were interpolated from the preceding and succeeding measurements.

Figure 1a shows the time evolution of the variables of Table 1 including a 24 h running mean thus providing a clear characterization of the sampling location for the year analyzed. The most important aspects are:

- The absence of strong winds exception for drainage flows connected to the vicinity of the Alps and local breezes due to the vicinity of the lake basin.
- The temperature evolution through out the year is typical of a mid latitude location with fairly cold winters and warm summers.
- Relative humidity is fairly high through out the year due to the presence of the lake basin and the stagnation of air masses within the Po Valley.
- Very small precipitation (a particularly dry year for the region) mainly characterized by short periods of relatively intense rain.
- Nitrogen oxides are fairly abundant due to the rural character of the area but also the relative vicinity of industrial areas and intense traffic roads.
- Radon's concentration is also fairly high ranging from a minimum value of few Bqm^{-3} in May to a maximum 50 Bqm^{-3} in December.

The correlation between radon concentration and the other variables is also given in Table 1. All correlations are consistent with the expectations. Namely, negative correlation with wind speed and precipitation responsible for reducing the concentration in the boundary layer and its emission at the surface, respectively; negative correlation with temperature which has an opposite diurnal cycle compared to radon. High positive correlation with relative humidity is found, which on the contrary has a similar diurnal cycle; and also a fairly large positive correlation with NO_x . The correlation with atmospheric pressure is positive in contrast with what found by other studies (e.g. Moses et al., 1973). For all correlation coefficient a significance test of 99% has been calculated thus making them all plausible values. However what




Fig. 1. (a): Time series of wind speed, temperature, relative humidity, pressure, NO_x and Radon-222 concentration measured during 1 year at the EMEP station. The red line represents a 24 h running mean. (b): Monthly evolution of correlation coefficients of radon concentration and all the variables of Table 1.




Fig. 2. (a) (yellow curves) Normalized power spectrum of wind speed, temperature, relative humidity, and NO_x , as a function of period expressed in hours; (black curves) 12 h running mean. (b) Same as (a) but for radon concentration. The red long-dashed line in panels (a) and (b) represents the red noise spectrum described in Eq. (1) in the text and the dot-dashed line the 95% confidence spectrum.

makes this global analysis a bit surprising is the very low figures of the correlation coefficient of precipitation and wind, in absolute value. A more detailed analysis has been conducted on a monthly basis as shown in Fig. 1b. In the case of wind a maximum value of anti-correlation is found at 0.6 in April and the largest values are found in general in the first part of the year. However, for precipitation values are very small also on a monthly basis. This is directly connected to the scarcity of precipitation events and the abundance of dry periods through out the year. One could obviously reduce further the time window of investigation and focus only on

the precipitation periods; however this would be a subjective procedure that would not lead much added value. What we can conclude from the correlation analysis is that the expected wind speed and precipitation signatures are found and that the low values are motivated in the case of precipitation by the scarcity of precipitation periods.

3 Wavelet analysis of radon monthly evolution

We start by analyzing the Fourier spectrum of the one-year time series of the variables of Table 1 in order to first identify globally the dominant scales. Figure 2a shows the power spectrum of the meteorological variables and NO_x and Fig. 2b the spectrum of radon. As pointed out by Torrence and Compo (1998) a power spectrum analysis of geophysical data requires the definition of a null-hypothesis to be used as term of comparison of the actual signal spectrum and for the identification of significant features within it. The figures also display the red noise spectrum obtained from the parameterization by Gilman et al. (1963) and the 95% confidence spectrum. The first is obtained from:

$$P_k = \frac{1 - \alpha^2}{1 + \alpha^2 - 2\alpha \cos\left(\frac{2\pi k}{N}\right)} \quad (1)$$

for $k=1, \dots, N/2$, where N is the number of samples, α is given by:

$$a = \frac{\alpha_1 + \sqrt{\alpha_2}}{2} \quad (2)$$

In Eq. (2), α_1 and α_2 are, respectively, the lag-1 and lag-2 autocorrelation coefficients of the variables analyzed. In other words with Eqs. (1) and (2) the spectrum of red-noise with the same autocorrelation signature of the original signal is created. By comparing the spectra of the signal with that of red-noise we can distinguish real features from “noisy” ones. Figures 2a and b also show the spectrum obtained by a running average of the original one with a pace of 12 h.

As far as the meteorological variables and NO_x are concerned we can notice the following:

- Wind speed: The spectrum is above the red noise one up to a period of 30–40 h, and is significant below the 12 h scale. Energy peaks can be noticed at 8 and 12 h. This clearly indicates the presence of periodic processes like lake breeze that have a diurnal or semi diurnal periodicity. The 24 h signal just peaks out of the confidence line.
- Temperature, relative humidity and NO_x : The spectra of these three variables are comparable as they are all atmospheric tracers transported and diffused in a similar way. They all show the presence of significant peaks at 24, 12, 8, 6 and 4 h. In the case of temperature peaks are also visible at 3 h time scale.

- Pressure (not shown): peaks are present at the diurnal and semi diurnal scale though the associated variance is much smaller compared to the other variables.
- Precipitation (not shown): No clear structure is present in the case of precipitation which shows a significant signal only at periods smaller than 10 h, due to the scarce and short precipitation events.

The similarities between the radon spectrum and the one of the other tracers can be seen by analyzing Fig. 2b. In general, up to a scale of 50 h the radon spectrum lays above the red noise power spectrum and it is above the 95% confidence spectrum up to a scale of 20 h. Within this range of scales one can clearly distinguish the presence of six energy peaks positioned at 24.063 h, 11.98 h, 7.99 h, 5.9 h, 4.79 h. All the peaks refer to the daily evolution of radon concentration which is governed by the daily evolution of the boundary layer depth and the constant emission from the surface. The shortest time scales are however indicating the occurrence of processes affecting the concentration and that produce a variance at smaller scales than the half day or 24 h cycle. The result presented in Fig. 2b is not completely surprising as the mechanism of radon emission and dispersion in the boundary layer are known and given the spectrum of the other variables described above. However, it represents just a global analysis which does not provide any insight on the actual concentration time series. Therefore, it is interesting to verify whether the peaks at periods smaller than 24 h are a constituting feature of the dataset or associated to specific processes and time periods.

In order to detail the behavior of radon's time evolution we make use of wavelet analysis. The latter consists of a localized Fourier transform by means of a wave confined, in this case, in time and with variable wavelength and amplitude. The wavelet adopted here is the classical Morlet function defined as:

$$\Psi(\eta) = \pi^{-\frac{1}{4}} e^{i\omega_0\eta} e^{-\frac{\eta^2}{2}} \quad (3)$$

where η is a non-dimensional time parameter and ω_0 is the non-dimensional frequency (assumed equal to 6 to guarantee admissibility of the function; Farge, 2000). Given the time series of radon concentration c_i with $i=0, \dots, N-1$ time spaced of δt , the wavelet transform reads:

$$W_n(s) = \sum_{n'=0}^{N-1} c_{n'} \Psi^* \left[\frac{(n' - n) \delta t}{s} \right], \quad (4)$$

where * indicates the complex conjugate of the wavelet function. In Eq. (4), n and s govern respectively the translation of the function along the time coordinate and the scales represented by the function. By moving the wavelet along the time dimension, it resonates with the time series c_i structures with scale s , and provides information about the signal variance, the scale and the location in time at which it occurs. By repeating the procedure for a range of scales one can obtain the

so-called scalogram i.e. a three-dimensional representation of the amplitude variability with scale and time. A number of aspects have to be considered when selecting the wavelet function, for its normalization and for the selection of the scale range s in relation to the time series analyzed. Here-with we refer the reader to the comprehensive description of the problem described by Torrence and Compo (1998).

Figures 3a–f show the wavelet analysis of the radon time series relating to the months of January (Fig. 3a), March (Fig. 3b), May (Fig. 3c), July (Fig. 3d), September (Fig. 3e), and October (Fig. 3f). For the sake of synthesis only these months were selected as they contain features that are also present in the others. In all plots the x-axis is time expressed in hours from the beginning of the month. The upper panel gives the time series of radon through out the month and the lower panel the corresponding scalogram. The y-axis of the scalogram refers to the time scale ranging from 2 to 200 h. The contours represent the square of the wavelet amplitude and therefore the variance or power of the signal. The contours in all scalograms of Figs. 3a–f are normalized by the global wavelet spectrum so that they can be compared with one another. Similarly to what has been done for the Fourier analysis of Fig. 2b, one can determine the 95% percent confidence that the wavelet will fall in a specific range of scales and times. The latter is obtained by means of a χ^2 test as described in Torrence and Compo (1998). The white contour superimposed to the colored ones is the 95% confidence contour. The hatched surface defines the so called cone of influence (COI), it is the region where the wavelet power drops by a factor e^{-2} . The power drop is a consequence of padding the time series with zero values at the edges in order to reach a number of components corresponding to a power of two and adopted to facilitate the wavelet analysis. The area outside the COI therefore defines the scale-time range in which the wavelet transform strictly refers to the actual time series values.

By inspection of the Figs. 3a–f one can immediately identify the differences with the power spectrum of Fig. 2b. In all scalograms the wavelet analysis identifies significant (included in the white contour) power levels in correspondence to the 24 h scale as determined by the Fourier analysis. This time the power peaks can also be related to the time coordinated. Therefore, for example in January (Fig. 3a), the diurnal variation is disrupted up until the end of the first week, while in the following week the 24 h scale is evident. During the months of March, July, September and October this feature is constantly identified. On top of the diurnal cycle which is also clearly evident by analyzing the time series, the wavelet analysis identifies relevant power contributions at lower scales. January is again an example in this respect. There is a secondary periodic contribution at 12 h which is not present in the other months and that was globally identified by the Fourier analysis. This contribution is visible in the second week of January as a constant contour at $s=10-12$ h and in a more patchy way in the third and fourth weeks.




Fig. 3. Time series and scalograms of radon concentration for January (a), March (b), May (c), July (d), September (e), and October (f) 1997. The black rectangles in panel (d) correspond to periods with missing data. White contour 95% confidence contour. Hatched surface Cone of Influence, see text for details.





Fig. 3. Continued.

**Fig. 3.** Continued.

Having identified the exact location in time of occurrence of the variance maximum, one can connect it to the actual portion of the signal that generates it and therefore identify its origin. The peak is caused by the fact that the concentration grows during the night, levels off by the end of it, before it reduces during the day. This produces the appearance of a second scale in the scalogram as the night and day evolution of concentration can be clearly separated by the analysis. The 12 h period is a typical feature of January and February (not shown) and does not appear systematically in any other month of the year. By comparing the January scalogram and time series with those of the other months, one can see that in the other cases the transition between night and day trends is very sharp and therefore no additional structure beyond the diurnal cycle is identified.

The eight hour timescale identified by the Fourier analysis can also be seen in the scalograms of January (5 cases), February (3 cases), March (4 cases), April and May (1 case), September (1 case), November (1 case) and December (2 cases). In October it is much more unfrequented and it disappears during the summer period. A more detail analysis of the time series of concentration in the specific time window where energy is found, indicates that it could be connected to the nighttime boundary layer formation and evolution. In correspondence to the time range in which this scale appears we can notice:

- i. A sudden increase of concentration due to the collapsing of the daytime ABL. This phase lasts between 6 to 8 h;
- ii. The growth of concentration between 10:00–12:00 p.m. and 08:00 a.m. of the following day with a different trend;
- iii. The sharp collapsing of concentration due to the rapid growth of the ABL between 09:00 a.m. and 02:00 p.m.;
- iv. After this phase the concentration remains at constant low levels until the cycle eventually repeats.

Clear examples of these cases are presented in Figs. 4a–d, 5a and b. The figures show how, with varying intensities in the peaks, the double growth trend feature clearly corresponds to the peak at 8 h. In the case of Fig. 4a for example the minimum at hour 515 corresponds to the concentration at 06:00 p.m. when the daytime boundary layer has totally collapsed, while the peak value corresponds to 06:00 a.m. In this case it is clear that 7 h after the collapse of the daytime boundary layer a new development occurs in the evolution of its nighttime counterpart that leads to a different growth rate of the concentration. The variation of growth rate could be connected to steady reduction of the boundary layer height occurring in the presence of a constant cooling rate at the surface (Nieuwstadt, 1981) as expected in winter months. In March the same feature can be noticed but it appears more sporadically. The investigation of wind speed

evolution with time shows the presence of calm of wind during the night thus endorsing the radiative cooling-based stability conditions which would lead to a constant decrease of the boundary layer depth. Unfortunately only micrometeorological data, that are not available, could allow a comprehensive explanation of the process.

Below the eight-hour scale, all scalograms allow also the identification of isolated features that are connected to sudden variations of the concentration level. The scalograms indicate that the lower time scale's peaks relate to sudden increases or decreases of concentration which are randomly distributed and do not constitute a constant feature of the power spectrum as Fig. 2b seems to indicate. This is typically the case for the first week of March where the sudden decrease of concentration produces large power variation at scales ranging from 2 to 10 h and extending over a time period of approximately 30 h. When the diurnal variation is not perturbed by external events, one can notice that the majority of small scale perturbations tend to occur rather systematically during the nighttime hours either on the first phase of the concentration growth or in the second as described in (I) and (II). A clear case of such event is the nighttime occurrence of double peaks as in the case of January (Fig. 3a, 5th concentration peak from the right), July (Fig. 3d, 5th, 6th and 10th concentration peaks from the right). The variability of concentration in these cases depends strongly on the variability of shallow nocturnal boundary layer which can be sensitive to sudden wind speed variations like the formation of low level jets. The double peak feature seems systematically connected to the 6 h scale identified by the Fourier spectrum. Figure 5c shows mixed conditions where the presence of energy contributions at scales ranging from 4 to 8 h are due to a variation of the concentration trend and sudden peaks of concentration, Figs. 5d and e show constant trend but sudden drop of the concentration at sunrise.

The role of external parameters in perturbing the radon concentration can be clearly seen in May. During this month the diurnal variation disappears across the first and the second week as a consequence of a clear reduction of the concentration levels. The reason for that will be analyzed in the next section.

The wavelet analysis has identified that while the 24 h scale is a constant feature of the daily evolution of the radon concentration, the 12 and 8 h scales appear mainly in correspondence to the nighttime period when the concentration changes growth rate after the collapse of daytime turbulence and in correspondence of concentration fluctuations therein. The smallest time scales (6 and 4 h) identified by the Fourier analysis cannot be attributed systematically to specific moments and therefore feature of the time series of radon but occur randomly in the time series.




Fig. 4. Single daily evolution of radon concentration and corresponding scalogram (blowup from monthly analysis). White contour 95% confidence.

4 Wavelet analysis of meteorological variables and relationship with the radon time series

The variability of radon concentration in the atmosphere is strictly connected to the meteorological conditions as the emission from the surface is in principle constant in time and space for a given soil type (Moses et al., 1963). In this section we will analyze the dependence of radon concentration on the meteorological situation at the measuring station during the year of measurements. The section will focus first of all on the identification of periods of time where a clear effect of precipitation and advection can contribute to the perturbation of radon concentration, further to that a hypothesis based on wavelet analysis will also be presented.

As pointed out by several works in the past, two meteorological variables affect radon concentration in the atmosphere or more specifically in the boundary layer, namely wind speed and precipitation. The impact of varying wind speed is due to two main reasons: advection of radon free or radon rich air, increase of shear driven turbulence with consequent effective dilution of the radio nuclide and concentration reduction. For the sake of completeness also the increase of surface heat flux and convective conditions should be accounted as a cause of concentration reduction. Precipitation affects radon concentration through rain out and wash out which according to few authors can remove the radio nuclide from the air (e.g. Gat et al., 1966) and mainly by reducing the exhalation flux from the surface. The presence of a water




Fig. 5. Same as Fig. 4.




Fig. 6. Time series of wind speed, precipitation, radon and corresponding scalograms for the month of May 1997. The red curve in the precipitation time series corresponds to time cumulated values. In the scalograms the white thick contour indicates 95% confidence level and the hatched surface the COI (see text). The two vertical dashed lines are explained in the text. Color bar shows wavelet power relative to global wavelet spectrum.

saturated layer at the surface, ice or snow cover can delay or prevent radon from being exhaled and therefore affect the emission rate. Some authors (e.g. Hatuda, 1953; Moses et al., 1963; Kranner et al., 1964; Schery et al., 1984) consider also atmospheric pressure variations as possibly responsible for variation of surface emissions. Namely increase (decrease) of pressure would damp (increase) the exhalation flux from the soil. As shown before, the present dataset does not show any clear feature in this respect and on the contrary a positive correlation between the pressure time evolution and that of radon concentration. For the sake of completeness

it should be mentioned that the detailed studies of Guedalia et al. (1970); Clements and Wilkening (1974); Guedalia et al. (1980); Ishimori et al. (1998) and Katoaka et al. (2003), indicate that only sudden drops or increases of pressure (of the order of 10 to 15 hPa) can cause a modification of the radon emission. Such occurrences are never recorded in the present data set.

The radon time series analyzed in this study is particularly interesting since the measurements were conducted in a location and time period where low wind conditions can be generally found apart from sporadic short term events. During

1997 few precipitation events were recorded and mainly concentrated in short-term intense rain periods.

An interesting case is presented in Fig. 6. The figure shows the time evolution of wind speed, precipitation and radon concentration measured at the EMEP station in May 1997. Each time series in the figure is presented together with the corresponding wavelet analysis. In the case of precipitation one can notice the superimposition of two time series: black corresponds to the actual time series, red is cumulated precipitation. This representation allows the identification of the sporadic events and also their contribution to the total precipitation trend. The scalogram of wind speed identifies the presence of the 24 h periodicity connected to local breeze event typical of the region and smaller scale variability. As one can see from the time series, wind speeds are relatively small. During the whole month only three major precipitation events occurred, for a total of 77 mm of rain. A peak of 20 mm can be noticed in the first week (56.8 mm in the first 300 h) and during the third and fourth week two short and less intense precipitation events took place. The constant diurnal variation of radon time series of the first four days of the first week is sensibly affected by the first precipitation event. The diurnal periodicity is disrupted; the concentration reduces to a half of the original average value for the following eight days.

The inspection of the precipitation scalogram seems to indicate a qualitative correspondence between the scales associated to the precipitation event and the perturbation of the radon time series. In the figure, the two vertical dashed lines indicate the time range of extension of significant scale of the precipitation. As one can see it corresponds to the period of time over which the radon periodicity is perturbed. Roughly beyond this time period the radon time series shows again the typical diurnal variation. From Fig. 6 we can notice that later in the month two small and short events caused short disruptions in the radon time series. The significant timescales indicated by the scalogram is of order 10–20 h. The former hypothesis should not be taken beyond the pure level of speculation as it would need a considerable amount of additional evidence before it is proven. There are in fact a number of additional factors that are expected to influence the exhalation flux subsequently to the precipitation event, like the soil porosity and consequently the run-off timescale of the precipitated water. Whether or not the timescales identified by the wavelet analysis are relevant to the identification of the exhalation suppression time scale will need a deeper verification.

In spite of the wavelet analysis however, what we can evince is that even a short but intense precipitation event can influence considerably the concentration of radon in the surface layer over a long period of time and this evidence is an important aspect that will be discussed in the next section.

The combined effect of wind speed and precipitation on radon concentration is shown in Fig. 7. Across the second and the third week of December (highlighted by the two ver-

tical dashed lines) one notices a build up of wind intensity with scales ranging from 2 h to 10–50 h which gives room later on to a precipitation event of reduced intensity and with comparable time scales. The effect on the radon concentration can immediately be evinced from the time series. The scarcity of the total precipitated rain does not lead to a long lasting inhibition effect on the exhalation flux, which is immediately restored as it can be evinced from the immediate recovery of the average concentration level after the precipitation (compare the maximum amplitude scale of this scalogram with that of Fig. 6).

An interesting case, though different from the one analyzed so far, relates to July and is presented in Figs. 8a and b. In Fig. 8a, during the first week wind speed increases consistently and suddenly over a time scale of 12–15 h while a precipitation event takes place with a scale ranging from 2 to 50 h. During the corresponding period of time radon concentration does not show any distinctive pattern apart from a sudden peak occurring during the wind intensity maximum period (highlighted by the black dotted line). This peak represents an anomaly also considering the scale and the intensity of the precipitation event. The analysis of other variables such as temperature and NO_x concentration (Fig. 8b) leads us to conclude that the peak is probably due to the advection of an air mass carrying a high concentration of radon that went past the monitor (do consider the sudden drop of temperature in the time window of the wind intensity peak). The radon peak in fact corresponds to a sudden and very large peak of NO_x (middle panel) which passes during the intense wind period from few tens of ppb to 150 ppb thus indicating the transport of the tracer from a NO_x rich region. The radon peak could very likely be connected to the same advection process. No particular feature can be deduced from the analysis of the scalograms.

5 A proposal to the global scale modeling community

A remarkable feature of radon atmospheric phenomenology is the response time of radon exhalation and therefore concentration to the start and end of the precipitation event. Beside the evidence provided in the previous section (regardless of the corresponding wavelet analysis) which only relates to the atmospheric concentration rather than the flux, Ishimori et al. (1998) determine a recovery time of the exhalation flux of 1.5 days after a precipitation event of 15 mm. Israel et al. (1966) found a comparable reduction due to precipitation events of less than 3 h and rate above 3 mm/h. Megumi and Mamuro (1973) found a reduction of the ground exhalation rate of a factor 2 after a precipitation event of 93 mm.

The sensitivity of radon exhalation and concentration to small amounts of precipitation, also evident in the present dataset, points clearly towards the need of a parameterization of the process within global models (e.g. Jacob et al., 1997; Dentener et al., 1999; Josse et al., 2004). The latter




Fig. 7. Time series of wind speed, precipitation, radon and corresponding scalograms for the month of December 1997. In the scalograms the black contour indicates 95% confidence level and the hatched surface the COI. The two vertical dashed lines are explained in the text. Color bar shows wavelet power relative to global wavelet spectrum.

in fact normally make use of radon as a scalar for evaluation purposes. A normal practice in this context consists in setting a uniform flux value in space and time with only a reduction of the value above higher latitude where snow cover is supposed abundant and persistent throughout the year.

The response of radon exhalation flux to the latitudinal, time and intensity variability of precipitation should be accounted for explicitly in atmospheric dispersion models for an appropriate description of radon's atmospheric concentration distribution in space and time. Global model could include simple parameterizations that would account for a modulation of the flux depending on the precipitation

amount. In a first approximation one could think of a binary approach that would switch off or reduce the flux of a fixed amount for a predefined amount of time whenever precipitation exceeds a certain threshold. Indications on the reduction in dependence of precipitation amount as well as on the recovery time are already given in the literature although not for all kind of soils. Although very crude, this approach would be much more realistic than the constant flux assumption. Most of the effect could be expected to originate from the time and space variability of the precipitation event rather than the actual parameterization of the process.




Fig. 8. (a) Time series of wind speed, precipitation, radon and corresponding scalograms for the month of July 1997. The dashed line in the precipitation plot is the precipitation time series plot at a different scale to emphasize the start and end time of all precipitation events. (b) Time series of relative humidity NO_x , radon and corresponding scalograms for the month of July 1997. In the scalograms the white thick contour indicates 95% confidence level and the hatched surface the COI (see text). The three vertical dashed lines are explained in the text. The black rectangles in the radon time series correspond to periods with missing data. Color bar shows wavelet power relative to global wavelet spectrum.




Fig. 9. (a) Radon and precipitation time series during the month of October 1997. (b) Blow up of the time series portion of panel (a) not hatched. Crosses correspond to daily average values.

The hypothesis that the model flux could be assumed constant as it represents a surface averaged value has become weaker with time given the sensible increase of model grid resolutions. Further to that datasets of analyzed or re-analyzed precipitation fields at global scale are available for the application of a simple parameterization as the one described above. What is certain is that radon flux is inhibited by the precipitation and therefore over a period of time as long as a year or several months the overall budget in the troposphere should be adjusted accordingly.

6 Estimate of the radon surface flux from concentration time series

The analysis of the time series of the various months outlines the presence of specific periods during which the radon concentration behaves in a special way. This is typically the case of August, September and October. Figures 9a and b refer to the case of October. The upper panel shows the time series of radon concentration during the whole month. The part of the plot that is not hatched is the period of interest in this analysis. The plot also shows the time evolution of precipitation as one of the governing factors of radon concentration as shown before. In the time window highlighted, no precipitation event occurs and the wind speed ranges below 1 ms^{-1} . Between the 250th and 300th hour, a sharp decrease in radon concentration occurs coinciding with a pe-




Fig. 10. (a) Radon and precipitation time series during the month of September 1997. (b) and (c) Blow ups of the time series portion of panel (a) not hatched. Crosses correspond to daily average values.

riod with high wind speeds ranging between 3 to 4 ms⁻¹ but stopping at hour 331. From that moment on and for the following 6 days, the radon concentration shows the usual daily evolution though superimposed to a linear growth. Similar

cases are found in September (2 sequences Figs. 10a–c) and August (Figs. 11a–b). All the time series indicate a build up of radon in the boundary layer that occurs for more than two consecutive days. The well-behaved character of such occur-




Fig. 11. (a) Radon and precipitation time series during the month of August 1997. (b) Blow up of the time series portion of panel (a) not hatched. Crosses correspond to daily average values.

rences, which are the only ones found through out the year, can easily be spotted by simple inspection of the time series. In all portions, the maximum and minimum concentration increase linearly from one day to the other.

The absence of precipitation events in the selected periods leads to the conclusion that the exhalation rate of radon from the ground is constant. The absence of significant wind speeds and the inspection of NO_x concentration, lead to excluding advection of radon from other areas. One can therefore deduce that the radon build up is connected to a rather regular boundary layer excursion and a negligible detraining

ment of radon at boundary layer top.

The daily average concentration has been calculated for the various periods and a linear fitting has been obtained as depicted in Figs. 9b, 10b–c and 11b. In spite of the different months, all the linear fitting show a common slope value, namely 0.11 in October; 0.091 and 0.092 in September; 0.11 in August. The linear growth of concentration superimposed to the daily evolution is also used by Vinod Kumar et al. (1999) to validate their one-dimensional model.

The results of this analysis are used herewith for a crude calculation of the surface flux of radon. It should be spec-

ified that this might be different from the actual exhalation flux which occurs at the direct surface, but it rather represents an estimate of the turbulent flux in the first portion of atmospheric surface layer.

The time evolution of radon concentration can be written as:

$$\frac{dC}{dt} = -\frac{(F_H - F_S)}{H} - \lambda C, \quad (5)$$

where, the vertical divergence of the turbulent flux has been approximated by the ratio of the net flux (given as difference between the flux at the surface layer top F_H , and that at the bottom F_S), and the boundary layer depth, H . In other words Eq. (5) is a bulk expression for the boundary layer which is represented as a layer of depth H with a radon in-flux at the surface and an out-flux at the top. In Eq. (5) λ is the decay frequency (equal to $\log(2)/3.8 \text{ day}^{-1}$).

In this context we assume that:

$$F_H \ll F_S, \quad (6)$$

as justified by the concentration build up in the absence of external factors that could perturb the surface emission. Expression (6) thus assumes a negligible entrainment of radon-free air and detrainment of radon at the boundary layer top.

The analytical solution of Eq. (5) reads:

$$C = C_0 e^{-\lambda(t-t_0)} + \frac{F_S}{\lambda H} \left(1 - e^{-\lambda(t-t_0)}\right) \quad (7)$$

By Taylor expansion of the exponential relationship, Eq. (7) can be linearized to:

$$C = C_0 \left[1 - \lambda t + o(t^2)\right] + \frac{F_S}{\lambda H} \left[\lambda t + o(t^2)\right] \quad (8)$$

where t_0 has been assumed equal to 0. By neglecting the higher order terms and re-arranging the expression, Eq. (8) reduces to:

$$C = \left(\frac{F_S}{H} - C_0 \lambda\right)t + C_0 \quad (9)$$

The linear interpolation of the time evolution of concentration given in Figs. 8–10, provides the following expression:

$$C = mt + C_0 \quad (10)$$

where $m=0.1 \text{ Bqm}^{-3} \text{ h}^{-1}$. In order to retrieve the magnitude of the flux at surface we can equate the slope of Eq. (9) and that of Eq. (10):

$$F_S = mH + \lambda C_0^H \quad (11)$$

Considering that a bulk value for the boundary layer depth (H) can be confidently assumed of order 10^2 m , for all time periods and the considered location and that C_0 is order 1, Eq. (11) reads:

$$F_S \approx \left(\frac{1}{360} + \frac{10^2 \log(2)}{3.8 \times 24 \times 3600}\right) \frac{Bq}{sm^2}, \quad (12)$$

Table 2. Values of boundary layer height in meters used to simulate the four episodes described in Sect. 6.

	August	September (Fig. 10b)	September (Fig. 10c)	October
Day	140	120	140	130
Night	110	90	110	90

which reduces to:

$$F_S \approx (0.0022) \frac{Bq}{sm^2} \quad (13)$$

The order of magnitude of F_S is well within the ranges directly measured in several other locations with characteristics similar to the EMEP station at JRC (e.g. Guedalia et al., 1970; Kataoka et al., 2001; Levin et al., 2002). It should be noticed that in Eq. (12) the dominant term is mH which is an order of magnitude larger than the second one.

The order of magnitude of the flux, determined by the previous analysis, is then used to reproduce the portion of time series depicted in Figs. 9–11. The only free parameter in Eq. (7) is H , however starting from the regular behavior of the concentration in time, we assume the surface layer depth varying as a sinus function and having an night-day excursion during for the four cases as from Table 2. The solution of Eq. (5) with the use of a time varying H and a flux value of $0.005 \text{ Bqm}^{-2} \text{ s}^{-1}$ gives the result shown in Figs. 12a–d. The figure shows that the flux estimate and expression (7) are able to reproduce the linear growth of the radon concentration. Small variability of maximum and minimum concentration values are obviously connected to the crude representation of the boundary layer depth evolution which according to the expression used assumes always the same maximum and minimum values while in reality one expects more variability within the range of values assumed. In time, if no external event would occur to perturb the radon emission and boundary layer evolution, the concentration growth rate would reduce to zero as the emission would compete at one stage with radioactive decay.

7 Conclusions

A one-year time series of radon-222 collected in 1997 in the North of Italy has been studied. The time series is particularly interesting as it was collected in region and time period where meteorological events controlling the radon concentration in air were moderate in intensity and sporadic.

Given the time variability of radon concentration in air, wavelet analysis was used to better identify feature that the classical Fourier analysis would only provide globally. The latter has in fact identified the presence of dominant time scales in the time series that range from 4 to 24 h but




Fig. 12. Time evolution of the radon concentration presented in Figs. 9–11 (black curve), and modeled time series (red curve).

that cannot be associated to specific occurrences in time. Through wavelet analysis however, the energy corresponding to scales smaller than 24 h were precisely connected to specific portions of the time series. In such a way a more clear identification of the processes leading to the occurrence of the specific timescale was made.

The role of wind speed and precipitation in controlling the radon variability in time was also investigated. Portions of the year-long time series where the effect of precipitation and wind (separately and combined) on radon concentration were identified. In particular it was found that even a short term intense precipitation event can have a prolonged effect to the concentration in the atmosphere as due to the saturation of the first layers of the soil by water.

The wavelet analysis of this portion of the time series has lead to the formulation of a speculation that will need further verification. Namely the apparent existence of a correlation between the timescale identified through the scalogram and the effect on the radon concentration even in the case of short term precipitation events or periods of intense wind.

Specific time periods within different months of the year that identified concentration build up in the surface layer were analyzed more in detail. The concentration growth rate was found to be the same regardless of the different months and seasons thus indicating a common behavior of radon dispersion in the region. The growth rate was then used to obtain an estimate of the radon surface flux which is matching the findings of more detailed studies.

Radon-222 is once more demonstrated to be an excellent tracer for boundary layer studies. The well-behaved character of surface emission and the effect of precipitation and boundary layer turbulence, make it a perfect candidate for atmospheric dispersion analysis and characterization. The availability of micro meteorological measurements in combination to radon concentration and flux data would allow an even deeper investigation of radon dispersion and boundary layer properties. Future investigation will be dedicated to determining the effect of precipitation intensity on the radon surface flux and surface layer concentration. Special attention will also be given to the open question of the role of atmospheric pressure variations on radon emission.

Acknowledgements. The author is grateful to B. Ottobriani that has originally collected the radon data. To M. De Cort (JRC/IES, I) and J. Vilà-Guerau de Arellano (WAU, NL) for the fruitful discussions during the preparation of this work. M. Grazia Marczan and U. Facchini (University of Milan, I) are thanked for providing additional data, literature information and fruitful discussions based on their long experience on the radon issue. The contribution of J. P. Putaud (JRC/IES, I) is also acknowledged for the provision of meteorological data collected at the EMEP station during the radon sampling campaign. The stimulating contribution from two anonymous reviewers is acknowledged.

M. G. Lawrence

References

- Attié, J. L. and Durand, P.: Conditional wavelet technique applied to aircraft data measured in the thermal internal boundary layer during sea-breeze events, *Bound. Layer Meteorol.*, 106, 359–382, 2003.
- Beck, H. L. and Gogolak, C. V.: Time dependent calculations of the vertical distribution of ^{222}Rn and its decay products in the atmosphere, *J. Geophys. Res.*, 86, 6, 3139–3148, 1979.
- Beran, M. and Assaf, G.: Use of isotropic decay rates in turbulent dispersion studies, *J. Geophys. Res.*, 73, 27, 5297–5295, 1970.
- Clements, W. E. and Wilkening, M. H.: Atmospheric pressure effects on ^{222}Rn transport across the Earth-Air Interface, *J. Geophys. Res.*, 79, 5025–5029, 1974.
- Dentener, F., Feichter, J., and Jeuken, A.: Simulation of the transport of Rn^{222} using on-line and off-line global models at different horizontal resolutions: a detailed comparison with measurements, *Tellus*, 51B, 573–602, 1999.
- Druilhet, A. and Fontan, J.: Utilisation du thoron pour la détermination du coefficient vertical de diffusion turbulente près de sol, *Tellus*, 25, 199–212, 1973.
- Druilhet, A., Attié, J. L., de Abreu Sá, S., Durand, P., and Bénech, B.: Experimental study of inhomogeneous turbulence in the lower troposphere by wavelet analysis, in: *Wavelets: theory, algorithms, and its applications*, edited by: Chui, K., Montefusco, L., and Puccio, L., 1–17, 1994.
- Facchini, U., Sesana, L., Milesi, M., De Saeger, E., and Ottobriani, B.: A year's radon measurements in Milan and at EMEP station in Ispra (Lake Maggiore, Italy), in: *Air pollution Conference (VII)*, edited by: Brebbia, C. A., Jacobson, M., and Power, H., ISBN 1-85312-693-4, 1112, 1999.
- Farge, M.: Wavelet transform and their application to turbulence, *Ann. Rev. Fluid. Mech.*, 24, 395–457, 2000.
- Farge, M. and Schneider, K.: Analysing and computing turbulent flows using wavelets, in: *New trends in turbulence. Les Houches 2000*, edited by: Lesieur, M., Yaglom, A. and David, F., 74, 449–503, Springer, 2002.
- Galmarini, S. and Attié, J.-L.: Turbulent transport at the thermal internal boundary-layer top: wavelet analysis of aircraft measurements, *Bound. Layer Meteorol.*, 94, 175–196, 2000.
- Gao, W. and Li, B. L.: Wavelet analysis of coherent structures at the atmosphere-forest interface, *J. Appl. Meteorol.*, 32, 1717–1725, 1993.
- Gat, J. R., Assaf, G., and Miko, A.: Disequilibrium between short-lived radon daughter products in the lower atmosphere resulting from their washout by rain, *J. Geophys. Res.*, 71, 1525–1535, 1966.
- Gilman, D. L., Fuglister, F. J., and Mitchell Jr., J. M.: On the power spectrum of “red noise.”, *J. Atmos. Sci.*, 20, 182–184, 1963.
- Guedalia, D., Laurent, J.-L., Fontan, J., Blanc, D., and Druilhet, A.: A study of Radon 220 emanation from soil, *J. Geophys. Res.*, 75, 357–369, 1970.
- Guedalia, D., Ntsila, A., Druilhet, A., and Fontan, J.: Monitoring of the atmospheric stability above an urban suburban site using sodar and radon measurements, *J. Appl. Meteorol.*, 19, 839–848, 1980.
- Hatuda, Z.: Radon content and its change in soil air near the ground surface, *Mem.Coll. Sci., Univ. Kyoto, Japan, Ser. B*, vol. XX, 285–306, 1953.

- Ikebe, Y. and Shimo, M.: Estimation of the vertical turbulent diffusivity from thoron profiles, *Tellus*, 24, 29–37, 1972.
- Ishimori, Y., Ito, K., and Furuta, S.: Environmental effect of radon from Uranium Waste rock piles: Part I – Measurements by passive and continuous monitors, in: *Proceedings of the 7th Tohwa University International Symposium on radon and Thoron in the Human Environment*, edited by: Katase, A. and Shimo, M., 23–25 October 1997, Fukouka, Japan, pp. 282–287, 1998.
- Israël, H. and Horbert, M.: Results of continuous measurements of radon and its decay products in the lower atmosphere, *Tellus*, 18(2), 638–641, 1966.
- Kataoka, T., Yunoki, E., Shimizu, M., et al.: A study of the atmospheric boundary layer using radon and air pollutants as tracers, *Bound. Layer Meteorol.*, 101, 131–155, 2001.
- Kataoka, T., Yunoki, E., Shimizu, M., et al.: Concentrations of ^{222}Rn , its short-lived daughters and ^{212}Pb and their ratios under complex atmospheric conditions and topography, *Bound. Layer Meteorol.*, 107, 219–249, 2003.
- Kranner, H. W., Schroeder, G. L., and Evans, R. D.: Measurements of the effects of atmospheric variables on Rn-222 flux and soil gas concentration, in: *The Natural radiation Environment*, pp. 191–215, University of Chicago press, Chicago, Ill, 1964.
- Kristensen, L., Andersen, C. E., Jørgensen, H. E., Kirkegaard, P., and Pilegaard, K.: First-Order Chemistry in the Surface-Flux Layer, *J. Atmos. Chem.*, 27(3), 249–269, 1997.
- Jacob, D. J., Prather, M. J., Rasch, P. J., et al.: Evaluation and inter-comparison of global atmospheric transport models using ^{222}Rn and short-lived tracers, *J. Geophys. Res.*, 102, 5953–5970, 1997.
- Josse, B., Simon, P., and Peuch, V. H.: Radon global simulations with the multiscale chemistry and transport model MOCAGE, *Tellus*, 56B, 339–356, 2004.
- Levin, I., Born, M., Cuntz, M., Langerdörfer, U., Mantsch, S., Naegler, T., Schmidt, M., Varlagin, A., Verclas, S., and Wagenbach, D.: Observations of atmospheric and soil exhalation rate of radon-222 at a Russian forest site. Technical approach and deployment for boundary layer studies, *Tellus B*, 54(5), 642, 2002.
- Leyendecker, W., Brunm, C., Geissm, H., and Rembgesm, D.: Activity of JRC EMEP Station: 1997 annual report, EUR 18084 EN, 1998.
- Nieuwsdtat, F.: A rate equation for the nocturnal boundary layer height, *J. Atmos. Sci.*, 38, 1418–1428, 1981.
- Marcazzan, M. G., Rodella, C. A., and Artesani, R.: Misure della concentrazione di radon in aria esterna come indicatore della diffusione nei bassi strati dell'atmosfera, *Fisica, Istituto Lombardo B B127*, 127–247, 1996.
- Marcazzan, M. G. and Persico, F.: Valutazione dell'altezza dello strato rimescolato a Milano dall'andamento della concentrazione di ^{222}Rn in atmosfera, *Ingegneria Ambientale*, 26, 419–427, 1996.
- Megumi, K. and Mamuro, T.: Radon and Thoron exhalation from the ground, *J. Geoph. Res.*, 78(11), 1804–1808, 1973.
- Moses, H., Lucas Jr., H. F., and Zerbe, G. A.: The effect of meteorological variables upon Radon concentration three feet above ground, *J. Air Poll. Control Assoc.*, 13(1), 13–19, 1963.
- Pearson, J. E. and Moses, H.: Atmospheric Radon-222 concentration variation with height and time, *J. App. Meteorol.*, 5, 175–181, 1966.
- Salmond, J. A.: Wavelet analysis of intermittent turbulence in a very stable nocturnal boundary layer: implications for the vertical mixing of ozone, *Bound. Layer Meteorol.*, 114, 463–488, 2005.
- Schery, S. D., Gaeddert, D. H., and Wilkening, M. H.: Factors affecting exhalation of radon from a gravelly sandy loam, *J. Geophys. Res.*, 89, 7299–7309, 1984.
- Sesana, L., Caprioli, E., and Marcazzan, G. M.: Long period study of outdoor radon concentration in Milan and correlation between its temporal variations and dispersion properties of atmosphere, *J. Environ. Radioact.*, 65, 147–160, 2003.
- Sesana, L., Polla, G., and Facchini, U.: Misure di radon outdoor: turbolenza e stabilita' atmosferica nella citta' di Milano e in un sito della Pianura Padana, *Ing. Amb.*, 34(3/4), 169–179, 2005.
- Torrence, C. and Compo, G. P.: A practical guide to wavelet analysis, *Bull. Amer. Meteorol. Soc.*, 79(1), 61–78, 1998.
- Vinod Kumar, A., Sitaraman, V., Oza, R. B., and Krishamoorthy, T. M.: Application of a numerical model for the planetary boundary layer to the vertical distribution of radon and its daughter products, *Atmos. Environ.*, 33, 4717–4726, 1999.
- Zahorowski, W., Chambers, S. D., and Henderson-Sellers, A.: Ground based radon-222 observations and their application to atmospheric studies, *J. Environ. Radioact.*, 76, 3–33, 2004.