

Evaluation of ECMWF ERA-40 temperature and wind in the lower tropical stratosphere since 1988 from past long-duration balloon measurements

T. Christensen¹, B. M. Knudsen¹, J.-P. Pommereau², G. Letrenne³, A. Hertzog⁴,
F. Vial⁴, J. Ovarlez⁴, and M. Piot⁵

¹Danish Meteorological Institute, Copenhagen, Denmark

²Service d'Aéronomie, IPSL/CNRS, Verrières-Le-Buisson, France

³Centre National d'Études Spatiales, Toulouse, France

⁴Laboratoire de Météorologie Dynamique, IPSL/CNRS, Palaiseau, France

⁵Institut für Umweltphysik, University of Heidelberg, Heidelberg, Germany

Received: 1 February 2007 – Accepted: 27 February 2007 – Published: 5 March 2007

Correspondence to: T. Christensen (tic@dmi.dk)

**Evaluation of ECMWF
ERA-40 temperature
and wind in the
tropical lower
stratosphere**

T. Christensen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Abstract

The temperature and wind of the ECMWF ERA-40 reanalysis in the tropical lower stratosphere during the period 1988–2001 has been evaluated by comparison with independent in situ measurements of 21 IR Montgolfier and superpressure long-duration balloon flights performed by CNES from Pretoria (26°S) in South Africa in 1988–1989, Latacunga (1°S) in Ecuador in 1991–1998 and Bauru (22°S) in Brazil in 2000–2001. The ERA-40 temperature displays a bias varying progressively from +1.16 K in 1988–1989, to +0.26 K in 1994–1996 and –0.46 K after 1998, the latter being fully consistent with recent evaluations of ECMWF operational analysis from radio occultation and in situ long-duration balloon observations. The amplitude of the bias and its evolution are very similar to the results of a previous evaluation from radiosondes in 1991–2003, suggesting that the origin of the drift of ERA-40 might be mainly due to errors in the series of satellite measurements of MSU, replaced by AMSU in 1998, assimilated in the model.

The ERA-40 zonal wind speed in the lower stratosphere appears slightly overestimated by 0.7–1.0 m/s on average in both the tropics and equatorial region, that is by 5–10% compared to the average 10–20 m/s wind speed. This bias, fully consistent with a recent evaluation of ECMWF operational analysis in 2004, is found constant during the whole 1988–2001 period, suggesting that the difference in that case might originate in the model. Finally calculated trajectories using ERA-40, frequently used for analysing field observations, are found in error compared to that of the balloons by ± 500 km after 5 days and ± 1000 km after 10 days.

1 Introduction

Historical reanalyses such as the European Centre for Medium-Range Weather Forecasts 40-year reanalysis (ECMWF ERA-40) are widely used, from research in climate and atmosphere to ecosystems and health.

ACPD

7, 3423–3450, 2007

Evaluation of ECMWF ERA-40 temperature and wind in the tropical lower stratosphere

T. Christensen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Evaluation of ECMWF
ERA-40 temperature
and wind in the
tropical lower
stratosphere**

T. Christensen et al.

In stratospheric research, reanalysis data are used for various purposes, e.g. influence of temperature on photochemical reaction rates or wind data in chemistry transport models, but most important for studying the past evolution of the stratosphere is the long-term change of temperature and winds. However, although reanalyses capture the cooling of the stratosphere, the amplitude of the trend may be quite different from that derived from radiosondes or satellite observations, already different between themselves, for reasons difficult to assess (Karl et al., 2006). As an example, between 100 and 50 hPa between 10°S and 10°N, the trend since 1979 could vary from between -0.69 and -0.64 K/decade in the radiosondes depending on the evaluation, to between -0.37 and -0.29 K/decade for the satellites and -0.83 K/decade for NCEP and only -0.005 K/decade for ERA-40, while the causes for these discrepancies are not clearly identified.

Evaluations of the accuracy of the reanalyses are thus important, but, especially for the stratospheric part, are not easy, since observations are sparse and operational data, like those from radiosondes and certain satellites, are already assimilated in the analyses. Even though reanalyses are designed to be as homogeneous a data set as possible, the atmospheric observation network is constantly changing, and consequently, so is the reanalysis accuracy. A data set which is new to ERA-40, especially a data set covering the data sparse stratosphere of the tropics, as the balloon data from past campaigns presented in this work, can add significant information.

There are very few comparisons of ERA-40 stratospheric temperatures in the tropics and in addition mostly restricted to the 100 hPa or tropopause level. By comparing modelled temperatures to radiosondes at 100 hPa in the equatorial region between 10°N–10°S in 1991–2002, Randel et al. (2004) indicate (their Fig. 11) a drop in deseasonalized temperature anomalies from about $+0.5$ K to -0.5 K of ERA-40 after 1998 and onwards compared to 1991–1997, amplified to 2–3 K after 2001 in NCEP/NCAR and the METO analysis of the UK Met Office, but not in ERA-40. Though the cause of the drops was not clearly identified, the authors note that a potentially important change occurred with the introduction in NCEP and METO of temperature retrievals of

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Evaluation of ECMWF ERA-40 temperature and wind in the tropical lower stratosphere

T. Christensen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ATOVS satellite measurements in July 2001. From comparisons with global radio occultation (RO) temperature measurements during two and half years from March 2002 to August 2004, [Gobiet et al. \(2005\)](#) are concluding at an excellent agreement within ± 1 K on average with ECMWF in the lower stratosphere, but a systematic cold bias of up to 2 K in a narrow layer at the altitude of the tropopause, attributed to the different vertical resolution of the analyses (>1.3 km) and the RO measurements of 1 km at that altitude. Finally, the last piece of information available is the comparison between the HIBISCUS balloons and ECMWF in 2004, displaying an average cold bias of -0.9 ± 1.3 K ([Knudsen et al., 2006](#)).

Similarly, there are few wind comparisons available. From monthly mean comparisons with rawinsondes (assimilated in the model) and rocket sondes (not assimilated), [Baldwin and Gray \(2005\)](#) found very good agreement, especially from the late 1980's and onwards, with an RMS error of less than 1.4 m/s after 1985 and less than 1.0 m/s after 1995. Another comparison available is that of [Knudsen et al. \(2006\)](#) with the data of the HIBISCUS balloon flights in 2004 displaying a positive bias of ECMWF of 0.7 m/s (wind too strong) for the ECMWF zonal wind in the tropics, a negligible difference for the meridional wind and an average trajectory error of 500 km after 5 days.

In the frame of the HIBISCUS project, dedicated in part to an assessment of the quality of meteorological models ([Pommereau et al., 2007](#)), the historical records of long-duration IR Montgolfier and superpressure constant-level balloons launched in the tropics since 1988 were digitized. In this paper the newly digitized past balloon data are used to evaluate the accuracy of the ECMWF ERA-40 reanalysis temperatures, horizontal winds and trajectories in the tropical lower stratosphere and tropopause region. The method is described by [Pommereau et al. \(2002\)](#) and [Knudsen et al. \(2002\)](#) who studied the winter Arctic vortex.

The originality of this study is three-fold: the use of real data which is not assimilated in the re-analysis; data which stems from the sparsely observationally sampled tropical, lower stratosphere, and data which has a resolution sufficient to contain most of the variability of that region.

The focus of this paper is on ECMWF ERA-40 and the lower stratosphere. This work can be seen as part of the general and important study aimed at assessing the accuracy of meteorological analyses and reanalyses.

The paper is outlined as follows: Sect. 2 presents the long-duration balloon flights and describes the data collection and the accuracy of the temperature measurements and the position data for winds and trajectories. Finally that section presents briefly the main characteristics of the ECMWF ERA-40 data used in this paper. Sections 3, 4 and 5 describe the results of the evaluation of ECMWF ERA-40 temperature, horizontal wind and trajectories, respectively, and compare the results to the work of other authors. The last section summarizes the results and conclusions of this study.

2 Data

2.1 Past long-duration balloon flights

A large number of long-duration balloon flights have been carried out by the French Centre National d'Etudes Spatiales (CNES) for a variety of projects: small constant level super-pressure balloons (SPB) at 200 hPa and 100 hPa between 1966–1979, followed by Infra-Red Montgolfier (MIR) since 1981, and more recently 10 m diameter SPB since 1998. But, with the exception of the 480 balloons flown at 200 hPa in the Southern Hemisphere in 1971–1972 whose data were collected by a dedicated EOLE satellite (Hertzog et al., 2006), the limited data transmitted by simple HF systems, moreover handled manually, could not be recovered. Only data after 1988 archived on various media could be available.

For the 30 MIR and 3 SPB flights which were performed in the tropics between 1988 and 2001, reliable data could be recovered for 18 MIR and 3 SPB launches. The whole set of flights is listed in Table 1 as 24 flights, but three of these were separate payloads on the same balloon.

The balloons were launched from three different sites: Pretoria in South Africa

Evaluation of ECMWF ERA-40 temperature and wind in the tropical lower stratosphere

T. Christensen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

(25.73°S, 28.18°E) and Bauru in Brazil (22.36°S, 49.03°W) in the tropics in the stratospheric easterlies of the austral summer season, and at Latacunga in Ecuador (0.91°S, 78.62°W) where the wind direction and speed depend on the phase of the Quasi Biennial Oscillation (QBO). The balloons used were MIR of 36 000 or 45 000 m³ volume and SPB of 10 m diameter. MIR (see e.g. [Pommereau et al. \(2002\)](#)) are hot air balloons heated from below by the Earth thermal emission at night and by solar radiation during the day. Therefore, their altitude varies from 26 km (20 hPa) for the smaller size, 28 km (13 hPa) for the larger during daytime, to 17 km (90 hPa) and 24 km (30 hPa), respectively, at night, depending on the cloud cover, except during the first 3 days of flight before losing the helium when they could fly as high as 34 km (4 hPa). SPB are constant density (isopycnic) balloons flying between 58 hPa (19.9 km) and 65 hPa (19.2 km) depending on the weight of the payload. The SPB flights are described in detail by [Vial et al. \(2001\)](#) and [Hertzog and Vial \(2001\)](#).

Following scientific objectives, balloon capacity and evolution of technology, various payloads are often flown together: always a Samba or Inmarsat service payload of CNES for controlling the flight (cut down if below a prescribed altitude), coupled or not with H₂O or Rumba scientific payloads of LMD (Laboratoire de Météorologie Dynamique) or the SAOZ of Service d'Aéronomie. All of them are carrying an ARGOS satellite transmitter allowing the transmission of 30–50 kB of data per day, which may be affected sometimes by transmission errors of large amplitude. But in addition, Samba is equipped with a HF system at 15 Mhz, replaced in 2000 by a more powerful Inmarsat satellite system allowing the transmission of one format of data every 10–15 min. A major technological step in all the payloads is the replacement after 1994 of the ARGOS Doppler localisation system of ±2–5 km accuracy during the overpass of the satellites 4–5 times per day and the pressure sensor for altitude determination, by a GPS providing 3D location within 100–200 m as fast as required, further improved to 10–20 m after 2000.

Evaluation of ECMWF ERA-40 temperature and wind in the tropical lower stratosphere

T. Christensen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2.2 Temperature data

The measurement of temperature is a difficult task on long-duration balloons because of the solar heating of the sensor and its support, as well as of the gondola, little ventilated at low pressure in the stratosphere. Except on the SAOZ payload where the sensor is a Vaisala radiosonde thermocap, those used on all others are Veco aluminised micro-bead thermistors of $250\ \mu\text{m}$ mounted on one meter long booms 180° apart from the gondola (Samba, H_2O and Inmarsat) or better, but more fragile, hanging 5 m below (Rumba). They are all calibrated in a thermal chamber, for CNES payloads to within $\pm 0.5^\circ\text{C}$, and for LMD payloads to within $\pm 0.2^\circ\text{C}$. As an illustration of the difficulty of the measurements, Fig. 1 shows the difference of temperature between the two sensors mounted 180° apart from the gondola of the 25 day Samba-MIR flight of Dec 2, 1988, plotted against solar zenith angle (upper panel), and pressure (lower panel). The bias of 0.5°C remains constant, but the dispersion increases from $\pm 0.4^\circ\text{C}$ during nighttime at $\text{SZA} > 94^\circ$ at altitude levels below 30 hPa (24 km), to $\pm 7^\circ\text{C}$ during daytime at float altitude at 15 hPa (27 km). Also shown are the drop of dispersion at $\text{SZA} < 20^\circ$ around noontime when the payload is in the shadow of the balloon, and the measurements at high altitude during the first days of flight, the highest being during the first night after launch. Most of the noise is caused by the heating of the sensor passing in the lee of the rotating payload. The noise is significantly reduced in the Rumba mounting 5 m below the payload, but the observations display a systematic day-night offset due to the direct heating of the thermistor.

For all above reasons, daytime measurements will be ignored in the following and only nighttime data at $\text{SZA} > 94^\circ$ will be used, with the exception of two Rumba flights in 2001. The data of each flight have been carefully checked by comparison of measurements of all available temperature sensors. Biases never exceed $\pm 0.8^\circ\text{C}$ between the two Samba sensors and $\pm 1.2^\circ\text{C}$ between those and thermistors of other payloads calibrated independently, with two exceptions: the flight of Jan 14 1994 when one of the booms did not deploy correctly, and that of Jan 28 1994 displaying large biases likely

Evaluation of ECMWF ERA-40 temperature and wind in the tropical lower stratosphere

T. Christensen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

caused by a confusion of calibration files. Fortunately, temperature measurements are also available in both cases from the H₂O payload, showing an average difference of 1.1°C in the first case, but of -5°C in the second, onto which the Samba data has been normalised. Overall, an accuracy of ±1.2°C can be estimated for the full nighttime temperature data set.

2.3 Position data for winds and trajectories

In contrast to temperature, the accuracy of wind measurements improved with time with the evolution of technology. It varies from about ±1.5 m/s on average for the early data set based on ARGOS Doppler localisations of ±2–5 km uncertainty 4–5 times per day until 1994, to 0.15 m/s with a scrambled GPS information of ±100 m accuracy every 15 minutes in 1996 and 1998, to less than 0.01 m/s after 2000. Another parameter contributing to the accuracy of wind comparison with ERA-40 is the uncertainty on pressure/altitude. Until 1994, the altitude is derived from pressure measurements at 0.6 hPa resolution within ±1 hPa accuracy. The corresponding altitude error varies from about ±50 m at 50 hPa, the average MIR flight level during nighttime, to ±450 m at 15 hPa during the day. An average vertical gradient of horizontal wind of 2 m/s/km will thus introduce an additional error of 0.1 m/s at 50 hpa and 1 m/s at 15 hPa. Similarly, the GPS uncertainty ±100 m in 1996–1998 would introduce an average uncertainty of 0.2 m/s dropping to less than 0.04 m/s after 2000. Overall, the accuracy of horizontal wind comparisons with ERA-40 at daytime flight level improves from 2.5 m/s before 1994, to 0.17 m/s in 1996–1998 and 0.05 m/s after 2000. Finally, depending on the sampling of the localisation system, the number of wind measurements per day varies from 5–6 during the ARGOS period, to about a maximum of 96–144/day after 1996 depending on the number of Samba HF ground receiving stations. The number of temperature and localisation data available for each flight is displayed in Table 1. For flights with ARGOS localization (1988–1994), interpolation of the position was done in the digitization process. 23 flights have temperature data, 23 have horizontal wind data and 18 qualify for trajectory comparison: at least 10 days of data with gaps not

Evaluation of ECMWF ERA-40 temperature and wind in the tropical lower stratosphere

T. Christensen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

exceeding 10 h.

2.4 ECMWF ERA-40 reanalysis data

The ECMWF 40 year reanalyses (Uppala et al., 2005) are produced every 6 h with a T159 model using a 3-D variational assimilation (Courtier et al., 1998). The vertical level spacing in the stratosphere is 1.4 km. We have extracted the data in a 1.5×1.5 degree latitude-longitude grid from a T79 truncation. The temperatures and horizontal winds from ECMWF ERA-40 are interpolated to the balloon latitude, longitude and pressure using an interpolation procedure which is linear in time and space and log-linear in pressure.

3 Temperature comparison

Figure 2 shows the temperatures from the reanalysis, T_{E40} , the average of the two simultaneously measured temperatures, $\langle T_{OBS} \rangle$, the difference between the two measured temperatures, $\Delta T_{OBS} = T_1 - T_2$, and the temperature differences between ERA-40 and the balloon average measurement, $\Delta T = T_{E40} - \langle T_{OBS} \rangle$. The many data points where $\Delta T_{OBS} = 0$ are an artifact of setting T_2 equal to T_1 for the nine flights where there was only one temperature sensor.

359 data points with an absolute temperature difference ($|\Delta T|$) greater than 50 K due to ARGOS transmission errors were discarded, leaving 38 623 data points. The mean temperature difference and the standard deviation are: 0.35 ± 2.6 K. Even with the outliers beyond ± 50 K removed, the data set still exhibits outliers, and the use of median statistics is more robust. The median temperature difference and the 68% fractile are 0.37 ± 2.1 K; i.e. 68% of the data points have an absolute deviation from the median (0.37 K) of 2.1 K or less. For the two Rumba flights in 2001, daytime data is included. If only nighttime measurements are considered, the data sample is 36 598 data points and the mean temperature difference and standard deviation are 0.41 ± 2.6 K and the

Evaluation of ECMWF ERA-40 temperature and wind in the tropical lower stratosphere

T. Christensen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

median temperature difference and the 68% fractile are 0.41 ± 2.0 K.

The data above 10 hPa exhibit more noise, but there are not that many data points, and removing them makes very little change to the overall result. Removing data above 10 hPa leaves a sample of 37 527 with median temperature difference and 68% fractile of 0.38 ± 2.0 K. If in addition the daytime data for the 2001 Rumba flights are removed, the sample is 35 621 with median temperature difference and 68% fractile of 0.42 ± 2.0 K.

The dense clusters of data at 58 and 65 hPa in Fig. 2 are from the SPB flights. It can be seen that altitudes in the range 65–15 hPa are well covered. Above 8–10 hPa the data dispersion increases drastically.

The statistics for the 23 individual flights is presented in Fig. 3. The 68% fractiles range from 1.3 to 3.0 K, the largest corresponding to shortest flights. For the longest, there is no significant change of precision between 1998 and 2001, indicating that the improved GPS location/altitude measurements compared to ARGOS/pressure sensor before 1994 has little impact on the results. The high altitude data between 10–30 hPa generally have larger median values than below at pressure larger than 30 hPa, except for two flights in 1994, which have few data points (<100). The increased noise on shorter flights (e.g. the 9 days last flight of 1989) is likely the consequence of the higher altitude of the MIR during the first days when still partly inflated with helium. The average median of the temperature difference for the whole period is 0.37 ± 2.1 K, but there is an indication of a progressive reduction from 1.16 ± 2.0 K in 1988–1991, to 0.26 ± 2.2 K in 1994–1996 and -0.46 ± 1.7 K after 1998, which could translate in an average trend of -0.13 ± 0.03 K/year or 1.9 K during the 14 year period. Since the temperature sensors and their calibration procedures, moreover carried out at the two independent laboratories, remain the same during the whole period, there is no reason to suspect a systematic experimental drift in the measurements. The average -0.46 ± 1.7 K bias observed after 1998 is fully consistent with ECMWF temperature evaluations in the tropical stratosphere of Gobiet et al. (2005) from 2.5 years of global radio occultation measurements in 2002–2004, showing insignificant bias in the lower stratosphere,

**Evaluation of ECMWF
ERA-40 temperature
and wind in the
tropical lower
stratosphere**

T. Christensen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Evaluation of ECMWF
ERA-40 temperature
and wind in the
tropical lower
stratosphere**

T. Christensen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

with the exception of -2 K difference at the tropopause identified by the authors to result from the difference of vertical resolution between the two data sets. -0.46 ± 1.7 K difference between ERA-40 and the balloons is also very consistent with that derived from the comparison of ECMWF with the HIBISCUS balloons in 2004 displaying a cold bias of -0.9 ± 1.3 K (Knudsen et al., 2006). Finally, the trend or step changes since 1988 fully coincide with the observed evolution of the difference at 100 hPa between ERA-40 and radiosondes in the equatorial region from 1991 to 2003 found by Randel et al. (2004). The long-duration balloon data are totally consistent with those of the radiosondes, suggesting that differences with reanalyses, or satellite observations, are to be attributed to models or satellites and not to radiosondes.

The evolution of the difference between balloons and ERA-40 must thus be attributed to ERA-40. Since the reanalysis system has remained constant in time, this result suggests that the model itself is not the cause, but likely the data assimilated. Most of the information used by the model is that of the Microwave Sounding Unit (MSU) onboard the NOAA and TIROS polar orbiting satellites, replaced in May 1998 by the Advanced Microwave Sounding Unit (AMSU), further replaced by ATOVS in July 2001 in NCEP and METO, but not in ERA-40. Intersatellite offsets are known to occur, difficult to fully compensate particularly when changing the instrument type, as well as the method for correcting for drifts in diurnal sampling time because of the drift of the orbit, or the method for correcting calibration drifts associated with the temperature of the hot calibration target (Christy et al., 2003; Karl et al., 2006).

4 Horizontal wind comparison

Exactly as for the temperatures, the ECMWF ERA-40 wind fields are interpolated to the balloon position for comparison with the in situ measurements.

Figures 4 and 5 show observed and reanalysed zonal and meridional winds and the difference plotted respectively against latitude and altitude. The data points are not evenly distributed in latitude: the balloons tend to stay within $\pm 8^\circ$ of their launch latitude,

so that the flights launched from Latacunga from 1991 to 1998 remain in the equatorial region, while those flown from Pretoria in 1988–1989 and Bauru after 2000 stay near the tropic of Capricorn. At the equator, the wind speed varies from 20 m/s to –30 m/s depending on the phase of the QBO which is from the west in 1991 and from the east in 1994, 1996 and 1998. Since the flights in the tropics have always been carried out during the summer, the winds are always from the east there, being maximum (–30 m/s) in mid-summer, i.e. February 2001, and minimum (<10 m/s) immediately after the turn-around in November in 1988–1989 and 2000. As seen in Fig. 5, the wind speed increases with altitude when from the east, at the tropics as well at the equator above 40 hPa during the east phase of the QBO when the lower layers are in the west phase. In contrast, in 1991 the winds are blowing from the east at upper levels and from the west lower down.

The statistics for individual flights are displayed in Fig. 6. Overall, there are 18 794 data points displaying an average difference of 1.05 ± 5.0 m/s with ERA-40 for the zonal wind (0.82 ± 3.9 m/s for median and 68% fractile) and -0.04 ± 3.6 m/s (0.01 ± 3.2 m/s) for the meridional component. The absolute difference and 68% fractile are larger at the equator (1.00 ± 5.1 m/s) than in the tropics (0.71 ± 3.2 m/s) for the u component, as well as for the v component 0.01 ± 4.2 m/s and 0.01 ± 2.6 m/s. Overall, there is an absolute positive bias for the ERA-40 zonal wind at both the equator and the tropics, however relatively small (5–10%) in reference to the average speed.

This picture is also seen in Fig. 6 for the individual flights where all median values for the zonal component are positive except for two flights: 19910403 and 19960328. The first is when the QBO index was west at both 30 and 50 hPa, and the second was when the index was west at 30, but east at 50 hPa, suggesting also a positive bias of ERA-40 in east phase QBO. The largest median values and 68% fractile are observed in the three last equatorial flights of 1994 of relatively short duration (8, 9 and 7 days) and thus at higher average altitude than others; on the 1994 flights where the wind at low latitude during night-time was also from the east; and on the flight of 20001119 launched in light stratospheric wind immediately after the turn-around and

Evaluation of ECMWF ERA-40 temperature and wind in the tropical lower stratosphere

T. Christensen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

thus travelling very slowly in 19 days across the Pacific. Note that this flight is also the one showing the largest dispersion of the meridional component at the tropics. At the equator, the largest deviations of the meridional component are observed on the three constant level balloons around 19–20 km where the trajectories are showing relatively large meridional oscillations associated with waves poorly captured by ERA-40.

The amplitude and deviations reported here are larger than that observed between the rawinsondes/rocketsondes and ERA-40 by Baldwin and Gray (2005) displaying excellent agreement between model and observations in monthly mean amplitude and standard deviations of less than 1.4 m/s after 1985 and less than 1.0 m/s after 1995. The smaller dispersion of monthly averages compared to this work might indicate that most of the discrepancy observed here might be due to atmospheric variations of periods shorter than one month not well resolved by ERA-40. Besides inertia gravity waves, part of the explanation could come from planetary waves trapped in the equatorial wave guide, i.e. Kelvin and Rossby waves like those reported by Hertzog and Vial (2001). Finally, the average high bias of ERA-40 of 0.71 ± 3.2 m/s found here in the tropics between 10° – 30° S is very similar to the average difference of 0.7 m/s between ECMWF and the HIBISCUS balloons in 2004 found by Knudsen et al. (2006) in the same latitude range.

Overall, the comparison of wind speed derived from long-balloon flights since 1988 to the ERA-40 reanalysis suggests a permanent slight systematic high bias in the zonal component in the model of 0.7 m/s in the tropical summer stratospheric easterlies and 1 m/s at the equator during the East phase of the QBO. However, when compared to the absolute wind speed of 10–20 m/s, the relative bias remains limited, not exceeding 5–10%. Finally, no significant bias has been observed in the meridional component either at the tropics or at the equator, but a relatively large dispersion of the order ± 4 m/s likely due to waves of different periods not fully captured by the model.

Evaluation of ECMWF ERA-40 temperature and wind in the tropical lower stratosphere

T. Christensen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

5 Trajectory comparison

Trajectories were calculated with a 2nd order Runge-Kutta integration scheme with a time step of 30 min for the SPB and 10 min for the MIR balloons. Along the balloon flight track new trajectories were started every 2 h. The SPB trajectories were forced to be isopycnic, whereas the MIR trajectories were forced to follow the pressure measured on board the balloon. In Fig. 7 are shown trajectories for three representative, but (for clarity) fairly short flights. The top panel shows the 19980907 flight path at the equator which, as the other two 1998 flights, also at the equator (trajectories not shown), exhibits meridional displacements likely caused by Rossby-gravity waves (see Vial et al. (2001) and Hertzog and Vial (2001) which give detailed analyses of the dynamics of the lower stratosphere during these three SPB flights). The middle panel of Fig. 7 shows the 20001119 MIR flight in the tropics during the spring turn-around period crossing the Pacific in 19 days which exhibits the largest trajectory errors, in contrast to the fast zonal 20010221 flight in the midsummer (shown in the lower panel), crossing the Pacific in 5–6 days. The flights launched from Pretoria earlier in the season in November–December 1988 and 1989 during the growing phase of stratospheric easterlies tend to drift gradually equatorward. The 1991 flight, the only one launched when the QBO was in its westerly phase at 30 hPa, drifted first slowly westward when still at high altitude, then stayed for 12 days near 180° over the central Pacific, drifting westward at 15 hPa during daytime and back eastward at 60 hPa at night.

For each trajectory, the spherical distance between the observed and calculated trajectories was calculated as a function of time since the trajectory start point. For each of the 18 flights for which trajectory comparisons were carried out, the average distance as a function of time is plotted in Fig. 8. The three 1998 SPB flights at 19–20 km stayed within 9° latitude of the equator, resulting in larger errors than that of the MIRs above in the lower stratosphere. The Bauru flight from 20001119, depicted by a blue line in Fig. 8, travelling in the light wind latitude region at the northern border of the stratospheric sub-tropical jet, also encountered waves that were not well described

Evaluation of ECMWF ERA-40 temperature and wind in the tropical lower stratosphere

T. Christensen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

by ECMWF. In Fig. 9 the mean errors after 5, 10 and 12 days are shown for each of the individual flights. Error bars show the 68% confidence intervals which are calculated taking into account the lag one serial correlation by using Jones (1975, Eq. 2.14) For a flight with fairly large error bars, 20001119, the 68% confidence interval is also pictured by the shaded area in Fig. 8. The largest errors of 1200–1500 km after 5 days are observed on the three constant-level balloons in 1998 near the tropopause, the error being significantly smaller on all MIR flights at higher altitude in the stratosphere in the tropics as well as at the equator.

An overall variance for the combined data set of 18 individual results is simply found by summing the variances and dividing by the number of flights, since the flights are considered independent. Combining the results from the 18 independent flights, reduces the variance of the errors considerably: Overall, we find mean ERA-40 trajectory errors for the 18 flights of 512 ± 57 km after 5 days, 995 ± 131 km after 10 days and 1047 ± 157 km after 12 days.

This compares well with the results of Knudsen et al. (2006) who find trajectory errors in SPB and MIR flights at the tropics of about 500 km after 5 days for tropical and southern mid-latitude long-duration balloon flights in 2004 as compared to operational ECMWF analyses.

6 Conclusions

The temperature and wind of the ECMWF ERA-40 reanalysis in the tropical lower stratosphere during the period 1988–2001 have been evaluated by comparison with independent in situ measurements of 21 IR Montgolfier and superpressure long-duration balloon flights performed by CNES from Pretoria (26°S) in South Africa in 1988–1989, Latacunga (1°S) in Ecuador in 1991–1998 and Bauru (22°S) in Brazil in 2000–2001. Compared to that of the balloons, the ERA-40 temperature displays a bias varying progressively from +1.16 K in 1988–1989, to +0.26 K in 1994–1996 and –0.46 K after 1998, the latter being fully consistent with recent evaluations of ECMWF operational

Evaluation of ECMWF ERA-40 temperature and wind in the tropical lower stratosphere

T. Christensen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

analysis from radio occultation and in situ long-duration balloon observations. The amplitude of the bias and its evolution are very similar to the results of a previous evaluation from radiosondes in 1991–2003. The similarity of the difference with radiosondes and long-duration balloons suggests that the origin of the drift of ERA-40 might be mainly due to errors in the series of satellite measurements of MSU and AMSU since 1998 assimilated in the model.

The ERA-40 zonal wind speed in the lower stratosphere appears slightly overestimated by 0.7–1.0 m/s in both the summer tropical easterlies and in the equatorial QBO associated alternative westerlies and easterlies, that is by 5–10% compared to the average 10–20 m/s wind speed. This bias, fully consistent with a recent evaluation of ECMWF operational analysis in 2004, is found constant during the whole 1988–2001 period suggesting that the difference in that case might originate in the model.

Finally the calculated trajectories using ERA-40, frequently used for analysing field observations, are found in error compared to that of the balloons by ± 500 km after 5 days and ± 1000 km after 10 days.

Acknowledgements. We thank ECMWF for providing the ERA-40 reanalysis. This work was supported by the 5th European Commission Framework Programme through the HIBISCUS project, contract no. EVK2-2001-00055, the EC Framework Programme 6 through the SCOUT-O3 project, contract no. GOCE-CT-2004-505390 and The Danish Research Council for Technology and Production Sciences through the Calisto project.

References

- Baldwin, M. P. and Gray, L. J.: Tropical stratospheric zonal winds in ECMWF ERA-40 reanalysis, rocketsonde data, and rawinsonde data, *Geophys. Res. Lett.*, 32, L09 806, doi:10.1029/2004GL022 328, 2005. [3426](#), [3435](#)
- Christy, J. R., Spencer, R. W., Norris, W. B., Braswell, W. D., and Parker, D. E.: Error Estimates of Version 5.0 MSU/AMSU Bulk Atmospheric Temperatures, *J. Atmos. Oceanic Tech.*, 20, 613–629, 2003. [3433](#)

Evaluation of ECMWF ERA-40 temperature and wind in the tropical lower stratosphere

T. Christensen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Courtier, P., Andersson, E., Heckley, W., Pailleux, J., Vasiljevic, D., Hamrud, M., Hollingsworth, A., and Fisher, M.: The ECMWF implementation of three-dimensional variation assimilation (3D-Var), *Q. J. R. Meteorol. Soc.*, 124, 1783–1807, 1998. [3431](#)

Gobiet, A., Foelsche, U., Steiner, A. K., Borsche, M., Kirchengast, G., and Wickert, J.: Climatological validation of stratospheric temperatures in ECMWF operational analyses with CHAMP radio occultation data, *Geophys. Res. Lett.*, 32, doi:10.1029/2005GL022617, 2005. [3426](#), [3432](#)

Hertzog, A. and Vial, F.: A study of the dynamics of the equatorial lower stratosphere by use of ultra-long-duration balloons, 2. Gravity waves, *J. Geophys. Res.*, 106, 22 745–22 762, 2001. [3428](#), [3435](#), [3436](#)

Hertzog, A., Basdevant, C., and Vial, F.: An assessment of ECMWF and NCEP/NCAR reanalyses in the southern hemisphere at the end of the pre-satellite era: Results from the EOLE experiment (1971–1972), *Mon. Wea. Rev.*, 134, 3367–3383, 2006. [3427](#)

Jones, R. H.: Estimating the Variance of Time Averages, *J. Appl. Meteor.*, 14, 159–163, 1975. [3437](#)

Karl, T. R., Hassol, S. J., Miller, C. D., and Murray, W. L.: Temperature Trends in the Lower Atmosphere: Steps for Understanding and Reconciling Differences. A report by the Climate Change Science Program and the Subcommittee on Global Change, Climate Change Science Program and the Subcommittee on Global Change, Washington, DC, 2006. [3433](#)

Knudsen, B. M., Pommereau, J.-P., Garnier, A., Nunes-Pinharanda, M., Denis, L., Newman, P., Letrenne, G., and Durand, M.: Accuracy of analyzed stratospheric temperatures in the winter Arctic vortex from infrared Montgolfier long-duration balloon flights 2. Results, *J. Geophys. Res.*, 107, doi:10.1029/2001JD001329, 2002. [3426](#)

Knudsen, B. M., Christensen, T., Hertzog, A., Deme, A., Vial, F., and Pommereau, J.-P.: Accuracy of analysed temperatures, winds and trajectories in the Southern Hemisphere tropical and midlatitude stratosphere as compared to long-duration balloon flights, *Atmos. Chem. Phys.*, 6, 5391–5397, 2006, <http://www.atmos-chem-phys.net/6/5391/2006/>. [3426](#), [3433](#), [3435](#), [3437](#)

Pommereau, J.-P., Garnier, A., Knudsen, B. M., Letrenne, G., Durand, M., Nunes-Pinharanda, M., Denis, L., Vial, F., Hertzog, A., and Cairo, F.: Accuracy of analyzed stratospheric temperatures in the winter Arctic vortex from infrared Montgolfier long-duration balloon flights 1. Measurements, *J. Geophys. Res.*, 107, doi:10.1029/2001JD001379, 2002. [3426](#), [3428](#)

Pommereau, J.-P., Garnier, A., Held, G., Gomes, A.-M., Goutail, F., Durry, D., Borch, F.,

Evaluation of ECMWF ERA-40 temperature and wind in the tropical lower stratosphere

T. Christensen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Hauchecorne, A., Cocquerez, P., Letrenne, G., Vial, F., Hertzog, A., Legras, B., Pisso, I., Pyle, J. A., Harris, N. R. P., Jones, R. L., Robinson, A., Hansford, G., Eden, L., Gardiner, T., Knudsen, B. M., Larsen, N., Nielsen, J. K., Christensen, T., Cairo, F., Pirre, M., Marécal, V., Huret, N., Rivière, E., Coe, H., Grosvenor, D., Edvardsen, K., Donfrancesco, G. D., Ricaus, P., Longo, K., and Freitas, S.: An overview of the HIBISCUS campaign, *Atm. Phys. Chem. Discuss.*, 7, 2389–2475, 2007. [3426](#)
- Randel, W. J., Wu, F., Oltmans, S. J., Rosenlof, K., and Nedoluha, G. E.: Interannual Changes of Stratospheric Water Vapor and Correlations with Tropical Tropopause Temperatures, *J. Atmos. Sci.*, 61, 2133–2148, 2004. [3425](#), [3433](#)
- Uppala, S. M., Kållberg, P. W., Simmons, A. J., Andrae, U., da Costa Bechtold, V., Fiorino, M., Gibson, J. K., Haseler, J., Hernandez, A., Kelly, G. A., Li, X., Onogi, K., Saarinen, S., Sokka, N., Allan, R. P., Andersson, E., Arpe, K., Balmaseda, M. A., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Caires, S., Chevallier, F., Dethof, A., Dragosavac, M., Fisher, M., Fuentes, M., Hagemann, S., Hólm, E., Hoskins, B. J., Isaksen, L., Janssen, P. A. E. M., Jenne, R., McNally, A. P., Mahfouf, J. F., Morcrette, J. J., Rayner, N. A., Saunders, R. W., Simon, P., Sterl, A., Trenberth, K. E., Untch, A., Vasiljevic, D., Viterbo, P., and Woollen, J.: The ERA-40 re-analysis, *Q. J. Roy. Meteor. Soc.*, 131, 2961–3012, 2005. [3431](#)
- Vial, F., Hertzog, A., Mechoso, C. R., Basdevant, C., Cocquerez, P., Dubourg, V., and Nouel, F.: A study of the dynamics of the equatorial lower stratosphere by use of ultra-long-duration balloons, 1. Planetary scales, *J. Geophys. Res.*, 106, 22 725–22 744, 2001. [3428](#), [3436](#)

Evaluation of ECMWF ERA-40 temperature and wind in the tropical lower stratosphere

T. Christensen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 1. List of long-duration balloon flights. The launch date is given in the format yyyyymmdd (first four digits is the year, the next two digits is the month and the last two is the day of the month). Three flights in 2001 were separate payloads launched on the same balloon.

Site	Launch		Instrumentation		Duration		Number of data points		
	Date	Balloon	Payload	Days	T	u	v	Traj.	Loc.
Pretoria	19881113	MIR 36	Samba	45	2815	280	279	524	176
Pretoria	19881119	MIR 36	Samba	29	2413	0	0	240	62
Pretoria	19881202	MIR 36	Samba	25	2547	1178	1181	255	90
Pretoria	19881208	MIR 36	Samba	53	1557	1556	1548	609	155
Pretoria	19891113	MIR 36	Samba/H ₂ O	12	2746	2444	2444	137	53
Pretoria	19891112	MIR 36	Samba	51	691	827	827	188	59
Pretoria	19891123	MIR 36	Samba/H ₂ O	9	1283	472	472	0	40
Latacunga	19910403	MIR 36	Samba/H ₂ O	20	615	532	557	226	192
Latacunga	19940114	MIR 45	Samba/H ₂ O	41	1727	339	340	486	185
Latacunga	19940118	MIR 45	Samba/H ₂ O	8	159	25	24	0	76
Latacunga	19940123	MIR 45	Samba/H ₂ O	9	253	46	49	0	34
Latacunga	19940128	MIR 45	Samba/H ₂ O	7	158	41	42	0	16
Latacunga	19940203	MIR 45	Samba/H ₂ O	12	0	68	94	0	136
Latacunga	19960328	MIR 45	Samba	14	2437	896	898	0	896
Latacunga	19960405	MIR 45	Samba	24	5977	1948	1958	268	1948
Latacunga	19980825	SPB 10 m	Samba	25	1001	822	810	226	822
Latacunga	19980901	SPB 10 m	Samba	48	2732	2091	2119	557	2091
Latacunga	19980907	SPB 10 m	Samba	24	1305	975	976	279	975
Bauru	20001117	MIR 45	Samba/Rumba	18	439	203	217	159	203
Bauru	20001119	MIR 45	Samba/Inmarsat	19	1601	354	402	225	354
Bauru	20010213	MIR 45	Rumba	39	832	507	686	319	507
Bauru	20010213	MIR 45	Saoz	39	555	511	484	340	511
Bauru	20010213	MIR 45	Inmarsat	39	1062	1038	1048	142	1038
Bauru	20010221	MIR 45	Samba/Rumba	48	1828	1816	1817	234	1816

Evaluation of ECMWF ERA-40 temperature and wind in the tropical lower stratosphere

T. Christensen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

**Evaluation of ECMWF
ERA-40 temperature
and wind in the
tropical lower
stratosphere**

T. Christensen et al.




Fig. 1. Difference of temperature readings between two thermistors mounted on 1 m long booms 180° apart on the Samba payload (MIR flight of 2 December 1988). Upper panel: plotted against solar zenith angle. Lower panel: plotted against pressure.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Evaluation of ECMWF
ERA-40 temperature
and wind in the
tropical lower
stratosphere**

T. Christensen et al.




Fig. 2. Temperature comparisons between ECMWF ERA-40 and long-duration balloon measurements. Panel 1 (from left): T_{E40} . Panel 2: $\langle T_{OBS} \rangle$. Panel 3: $\Delta T_{OBS} = T_1 - T_2$. Panel 4: $\Delta T = T_{E40} - \langle T_{OBS} \rangle$.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 3. Statistics for each of 23 long-duration balloon flights. Temperature comparison between ECMWF ERA-40 and balloon measurements. On the x-axis all flights are grouped together year-by-year. Within one year, data points are equidistant regardless of the date of the flight. Big black dots: all available data points for the flight are used. Small red dots: only data points in the height interval 10–30 hPa are used. Small blue dots: only data points in the height interval below 30 hPa are used. Panel 1 (from top): Median of $\Delta T = T_{E40} - \langle T_{OBS} \rangle$ for each flight. Panel 2: 68% fractile of ΔT for each flight. Panel 3: Number of data points for each flight.

Evaluation of ECMWF ERA-40 temperature and wind in the tropical lower stratosphere

T. Christensen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Evaluation of ECMWF ERA-40 temperature and wind in the tropical lower stratosphere

T. Christensen et al.




Fig. 4. Horizontal wind comparison between ECMWF ERA-40 and long-duration balloon measurements as a function of latitude. From top to bottom panel: u_{OBS} , u_{E40} , v_{OBS} , v_{E40} , $\Delta u = u_{\text{E40}} - u_{\text{OBS}}$ and $\Delta v = v_{\text{E40}} - v_{\text{OBS}}$.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 5. Horizontal wind comparison between ECMWF ERA-40 and long-duration balloon measurements as a function of altitude. From left to right: u_{OBS} , u_{E40} , v_{OBS} , v_{E40} , $\Delta u = u_{\text{E40}} - u_{\text{OBS}}$ and $\Delta v = v_{\text{E40}} - v_{\text{OBS}}$.

Evaluation of ECMWF ERA-40 temperature and wind in the tropical lower stratosphere

T. Christensen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Evaluation of ECMWF ERA-40 temperature and wind in the tropical lower stratosphere

T. Christensen et al.




Fig. 6. Statistics for each of 24 long-duration UTLS balloon flights. Horizontal wind comparison between ECMWF ERA-40 and balloon measurements. On the x-axis all flights are grouped together year-by-year. Within one year, data points are equidistant regardless of the date of the flight. Big black dots: all available data points for the flight are used. Small red dots: only data points in the height interval 10–30 hPa are used. Small blue dots: only data points in the height interval below 30 hPa are used. Panel 1 (from top): Median of Δu for each flight. Panel 2: 68% fractile of Δu for each flight. Panel 3: Number of u data points for each flight. Panel 4: Median of Δv for each flight. Panel 5: 68% fractile of Δv for each flight. Panel 6: Number of v data points for each flight.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close


Full Screen / Esc

Printer-friendly Version


Interactive Discussion

**Evaluation of ECMWF
ERA-40 temperature
and wind in the
tropical lower
stratosphere**


T. Christensen et al.



7 SEP - 29 SEP 1998



20 NOV - 8 DEC 2000



22 FEB - 13 MAR 2001

Fig. 7. Trajectories for three flights: 19980907, 20001119 and 20010221. Balloon positions are shown in red. ECMWF ERA-40 trajectories were started every 2 h; for clarity, only every 12th trajectory (black lines) is shown; the starting points are shown with red plusses. The geographical grids show every 10 degrees in latitude and every 30 degrees in longitude.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Evaluation of ECMWF
ERA-40 temperature
and wind in the
tropical lower
stratosphere**

T. Christensen et al.




Fig. 8. Mean trajectory errors for 18 flights compared to ECMWF ERA-40. The red lines are the three equatorial SPB flights of 1998 (19980907 has the smallest error after 12 days). The yellow line correspond to the longest flight: the 19881208 Pretoria launch. The blue line is the 20001119 flight from Bauru, and for this flight the 68% confidence limits are indicated by the shaded area.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Evaluation of ECMWF ERA-40 temperature and wind in the tropical lower stratosphere

T. Christensen et al.




Fig. 9. Statistics for each of 18 long-duration balloon flights. Trajectory comparison between ECMWF ERA-40 and balloon measurements. On the x-axis all flights are grouped together year-by-year. Within one year, data points are equidistant regardless of the date of the flight. Black dots: Mean spherical distance after 5 days. 68% confidence interval indicated by black error bars. Blue crosses: Mean spherical distance after 10 days. 68% confidence interval indicated by blue error bars. Red crosses: Mean spherical distance after 12 days. 68% confidence interval indicated by red error bars. Panel 1 (from top): Mean error of ERA-40 trajectories. Panel 2: Number of trajectories.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion