

This discussion paper is/has been under review for the journal *Atmospheric Chemistry and Physics (ACP)*. Please refer to the corresponding final paper in *ACP* if available.

**Ice nucleation rate
from contact angle
and activation energy**

J.-P. Chen et al.

Parameterizing ice nucleation rates for cloud modeling using contact angle and activation energy derived from laboratory data

J.-P. Chen¹, A. Hazra¹, and Z. Levin²

¹Department of Atmospheric Sciences, National Taiwan University, Taipei, Taiwan

²Department of Geophysics and Planetary Sciences, Tel Aviv University, Tel Aviv, Israel

Received: 29 May 2008 – Accepted: 1 July 2008 – Published: 29 July 2008

Correspondence to: J.-P. Chen (jpchen@as.ntu.edu.tw)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The rate of ice nucleation in clouds is not easily determined and large discrepancies exist between model predictions and actual ice crystal concentration measured in clouds. In an effort to improve the parameterization of ice nucleating in cloud models, we investigate the rate of heterogeneous ice nucleation under specific ambient conditions by knowing the sizes as well as two thermodynamic parameters of the ice nuclei – contact angle and activation energy. Laboratory data of freezing and deposition nucleation modes were analyzed to derive inversely the two thermodynamic parameters for a variety of ice nuclei, including mineral dusts, bacteria, pollens, and soot particles. The analysis considered the Zeldovich factor for the adjustment of ice germ formation, as well as the solute and curvature effects on surface tension, the latter effects have strong influence on the contact angle. Contact angle turns out to be a more important factor than the activation energy in discriminating the nucleation capabilities of various ice nuclei species. By extracting these thermodynamic parameters, laboratory results can be converted into a formulation that follows classical nucleation theory, which then has the flexibility of incorporating factors such as the solute effect and curvature effect that were not considered in the experiments.

1 Introduction

Ice processes in clouds are important to precipitation formation thus the hydrological cycle. As Lau and Wu (2003) pointed out, more than 50% of the mid-latitude precipitation is produced via cold-cloud (involving ice) processes, whereas in the tropical region the proportion reaches 30%. The initiation of ice formation and cloud glaciation at temperatures above about -35°C usually requires the presence of ice nucleating particles. These so-called ice nuclei (IN) can lower the energy barriers of ice nucleation so that ice particles can be produced heterogeneously at relatively warm environment. They can initiate ice through the deposition mode, condensation-freezing mode, con-

Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tact freezing mode or the immersion mode (Pruppacher and Klett, 1997, p. 309).

Detailed information on atmospheric IN is rather scarce. Earlier studies measured the bulk concentration of atmospheric IN without knowing the physical or chemical properties of individual particles. Some of them characterized the ice nucleating capability of IN with the co-called “threshold temperature” (e.g., Heffernan and Bracewell, 1959; Isono et al. 1959; Carte and Mossop, 1960; Fletcher, 1962; Cooper and Saunders, 1980; Bowdle et al., 1985). However, many observational studies have shown that the number concentration of ice crystals in clouds is not strictly determined by temperature (e.g., Gulpepe et al., 2001; Korolev et al., 2003). In fact, the ice nucleating capability in the deposition mode can be better characterized with the degree of super-saturation with respect to ice (e.g., Bryant et al., 1959; Huffman 1973; Hussain and Saunders 1984; Stein and Georgii 1982). But such notion is not strictly correct either. According to classical theory, nucleation should be a probabilistic rate process, meaning that only a certain proportion of a particular kind of nuclei can be activated into ice in a specific length of time. Those “threshold” formulations are based on the assumption that the nucleation rate varies sharply from negligible to significant over a very narrow range of nucleation conditions (i.e. temperature or supersaturation). Under such a concept, different IN may be treated to nucleate at their specific critical conditions while identical IN must nucleate all together.

The above threshold concept is adopted popularly in many meteorological cloud models. Their formulations present the number of ice particles, generated through deposition or condensation-freezing nucleation, as a function of either temperature or ice supersaturation, or a combination of the two variables (e.g., Fletcher, 1962; Wisner et al., 1972; Cotton et al., 1982; Lin et al., 1983; Myers et al., 1992; Reisner et al., 1998). Although easy to use, such formulations have several deficiencies. First, the formulas are mostly empirical and do not consider the temporal and spatial variations of atmospheric IN. Take the empirical formula of Fletcher (1962) as an example, the amount of active IN is expressed as a function of temperature only. Yet, the concentration of atmospheric IN does vary tremendously in time and space, as well as in their origins.

Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Furthermore, by treating the nucleation process in a diagnostic way, such schemes cannot handle the situations when IN actually have been consumed earlier even in the same cloud, thus they tend to over-predict the number of ice particles. It is also impossible to recount the IN when the ice particles formed on them were evaporated.

Furthermore, these empirical formulas obviously lose track of individual IN, thus are not suitable for evaluating IN of different origins or physical properties such as size.

To simulate accurately the ice-phase processes remains a difficult task for cloud modelers to this date. One of the main complexities is the properties and distribution of IN. A few species of IN have been identified as important ice initiation agent in the atmosphere, such as mineral dust (Kumai 1951; Isono et al., 1959), volcanic ashes (Isono and Komabayashi, 1954; Isono et al., 1959), combustion particles (Hobbs and Locatelli 1969; DeMott 1990), and bio-aerosols which include whole organisms (e.g. bacteria, fungi, and phytoplankton), reproductive materials (e.g. pollens and spores) and plant fragments (Schnell and Vali, 1973; Schnell, 1974; Levin et al., 1987; Diehl et al., 2001).

As will be discussed later, significant progress has been made recently in laboratory measurement of ice nucleation rates of various IN. Combined with increasing model capability of simulating the emission and transport of atmospheric IN species (e.g., Lighthart 1997; Uno et al., 2003; Chen et al., 2004; Kishcha et al., 2005; Pasken and Pietrowicz, 2005), the next generation meteorological models may be able to adequately examine the roles that major IN played in precipitation formation. Such work would be important to the understanding of interactions between land surface (including the ecosystem) and the atmospheric hydrologic cycle (Barth et al., 2005).

Laboratory derived ice nucleation rates can be used adequately in models that resolves cloud microphysics for treatment of single or multiple IN species. However, few attempts were made to extract from these data useful information related to the properties of IN. If fundamental thermodynamic properties of IN can be obtained, they may not only help understanding the physical mechanisms of nucleation but also broaden the application of laboratory results by providing better parameterization for ice nucleation in cloud models. For example, ice nuclei, such as mineral dust, often contain

Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



solute coating that helps to acquire liquid water for freezing. Yet, freezing nucleation experiments often were measured under pure water conditions, and it is difficult to include solute effect into empirical rate equations derived from these experiments. With the help of fundamental analysis, it may be possible to integrate the solute effect or other factors into the nucleation rate formulas. In this study we intend to show this possibility by re-analyzing laboratory data of ice nucleation from various IN, then obtain basic thermodynamic parameters as described by the classical theory so that a more general and accurate parameterization of ice nucleation rate can be obtained for use in cloud microphysical models.

2 Method

In classical nucleation theory, the heterogeneous ice nucleation rates can be calculated by knowing a few key parameters that are specific to each ice nuclei, such as the activation energies and contact angles. If the classical theory holds well, then it is possible to extract these key parameters inversely from laboratory data. Although there have been a number of laboratory studies focused on heterogeneous ice nucleation, few tried to relate the measurement data to these key parameters. Here, we describe the method of analyzing classical formulas of heterogeneous ice nucleation, as well as the laboratory data and apply it to a variety of natural and anthropogenic ice nuclei.

2.1 Laboratory data

We focus on four types of IN that are common and potentially important to ice formation in clouds, namely, mineral dust, bacteria, pollen, and soot particles.

Mineral dust particles have long been identified as a common and efficient IN since Kumai (1951) and others found solid silicate particles in the central portion of snow crystals. Extensive studies have been conducted since to examine the threshold temperatures of dust for ice nucleation. But not until recently did comprehensive mea-

Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 surements of their nucleation rates were conducted. The work of Möhler et al. (2006) provided one of the most detailed information of ice nucleation, including the nucleation rates and corresponding environment conditions of temperature, pressure and saturation ratio. Although they measured ice nucleation on three types of dusts, we selected
10 data (excluding a couple of data points that might be problematic) for the Arizona test dust (ATD) because of their sufficient number of data and wide temperature range. Note that Archuleta et al. (2005) and Field et al. (2006) also measured ice nucleation on dusts, but they only provided the conditions required to activate a certain percentage of dust. These conditions pertain to the aforementioned concept of “threshold super-
15 saturation” or “threshold temperature,” which are not suitable for our purpose. But from Fig. 7 in Field et al. (2006) we were able to get nucleation rates for the Saharan dust (SD) and Asian dust (AD), which are more representative of natural dust than ATD.

The above measurements were performed in the large cylindrical aerosol chamber (AIDA, volume 84 m³) that was used as a moderate expansion cloud chamber at different cloud conditions like temperature, cooling rate, and ice supersaturation. The number concentration of ice particles was measured by OPC-PCS instrument, Small Ice Detector (SID, University of Hertfordshire, UK), and was retrieved also by FTIR extinction spectra. Mean adiabatic cooling rates (close to the time of nucleation) for the experiments were 1–2 K min⁻¹. Crystals mainly formed by deposition nucleation
20 mode. Rate of ice activated at each temperature range was obtained from the concentration ratio of activated particles to total particles divide by the specific time span. Hung et al. (2003) measured freezing nucleation on dust coated with ammonium sulfate solution. This latter data set is used not only to derive the similar thermodynamic parameters but also to examine the closure of surface energy calculation as will be
25 discussed in Sect. 4.

Bacteria are considered a major type of ice-nucleating bio-aerosols because of their ubiquity and efficient ice nucleating capability. Their freezing activity appears associated with the cell structure (Maki et al., 1974), and more precisely due to unique proteins that act as nucleation catalyst (Ruggles et al., 1993). Decaying plant leaves

Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Ice nucleation rate
from contact angle
and activation energy**

J.-P. Chen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

were also found to act as IN, but the ice nucleation was identified to be of bacterial origin. In fact, some plant frost injury has been shown to involve an interaction of certain leaf-surface bacteria (Lindow, 1983). Worldwide availability of such nuclei was established by finding ice-forming nuclei in plant litters collected in different climatic zones (Schnell and Vali, 1973). The subsequent studies of Vali et al. (1976), Yankofsky et al. (1981), Levin et al. (1987), and Hazra et al. (2004) showed that such biogenic IN might be released from the earth's surface to the atmosphere and initiate significant ice formation at temperatures as high as -2°C , even more effective than silver iodide which is an artificial IN commonly used for cloud seeding. In this study we took the nucleation data for two types of bacteria (*Erwinia herbicola* and *Pseudomonas syringae*) from Yankofsky et al. (1981) and one (*Pseudomonas aeruginosa*) from Hazra et al. (2004). *E. herbicola* and *P. syringae* are common plant-pathogenic bacteria that can infect a wide range of plant species, whereas *P. aeruginosa* not only infects plants but also animals and humans. But the above studies considered only immersion freezing nucleation mode, and we were not able to find data for the deposition nucleation on bacteria.

Freezing nucleus activity of *P. syringae* and *E. herbicola* (Yankofski et al., 1981) of drops containing bacteria were determined in the drop freezing spectrometer. Some of these experiments were performed with the drops held for 2 min at fixed temperatures below 0°C . The ice or water saturations were not recorded in these experiments. Nucleation activity of *P. aeruginosa* (Hazra et al., 2004) was conducted in a walk-in cold chamber at water saturation and at each temperature formvar coated slides were placed for 2 min to collect ice crystals.

Pollens are another category of bio-aerosols that may spread in large quantities and over wide ranges. Their surface contains active sites that enable them to take up water effectively. While there have been some investigations on the hygroscopic characteristics of pollens (Durham, 1941; Harrington and Metzger, 1963; Dingle, 1966), much less is known about their role as IN than about bacteria. Not until recently did Diehl et al. (2001, 2002) perform a series of measurements on ice nucleation from various

kinds of pollens. They found that pollens initiate ice nucleation not via the deposition mode but by immersion freezing or condensation-freezing within the investigated temperature and ice supersaturation ranges. Von Blohn et al. (2005) extended the study by including more pollen types. From Diehl et al. (2001) we took nucleation data for 5 four kinds of pollens: birch, oak, grass, and pine. As only the fractions of nucleation were provided, we again obtain their experiment time of 1 min by personal communication. In addition, our own study measured the freezing nucleation rates for the pollens of eucalyptus and China Rose following the procedure of Hazra et al. (2004). These results are also included in our analysis. Freezing nucleation of eucalyptus and 10 China Rose pollen was done in the same way as *Pseudomonas aeruginosa* mentioned earlier. Condensation freezing efficiency of different pollen (Deihl et al., 2001) carried out inside the walk-in cold chamber where glass flask having diameter of 30 cm was installed. At supersaturation with respect to water pollen grains were scattered at equilibrium temperature for a short period (nearly 1 min) and activated grains fell down onto a soap film.

When Kumai (1951) and others analyzed the nucleation center of snow crystals, they also found that some of the cores were composed of combustion products such as soot. Subsequent evidence of soot as ice nuclei was suggested by Hobbs and Locatelli (1969) as well as Langer (1973) who observed a significant increase of IN 20 downwind of burning forest or sugarcane fields. Similar mechanism was proposed by Ström and Ohlsson (1998) who found that cirrus ice crystals in a region of heavy air traffic often contain soot-like particles. In a laboratory experiment, Gorbunov et al. (2001) showed that soot surface contains chemical groups which can form hydrogen bonds with water molecules, thus enhance ice nucleation. However, in contrast to 25 the aforementioned IN species, soot particles can only initiate ice nucleation at rather low temperatures as pointed out by Hallett et al. (1986), DeMott (1990) and Dymarska et al. (2006). In the latter study, ice nucleation never occurred at temperatures above 248 K. Both DeMott (1990) and Dymarska et al. (2006) showed that soot particles initiate ice formation almost exclusively via the freezing nucleation mode. We took the

**Ice nucleation rate
from contact angle
and activation energy**J.-P. Chen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Ice nucleation rate
from contact angle
and activation energy**

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



nucleation fraction data of DeMott (1990), the one that provided sufficient information. For this experimental data, soot particles were generated using simple oxygen-deficient acetylene burner and injected into the CSU dynamic cloud chamber (volume 1.2 m^3) to see its nucleation activity during expansion to -40°C with adiabatic cooling of 1°C min^{-1} . The freezing rate per unit volume was calculated between -26°C and -34°C . Ice crystals were detected by extinction in a laser-based detection device (DeMott and Rogers, 1990). Most crystals formed by immersion freezing mode. Cumulative fraction nucleated as a function of temperature was presented in this experiment. Thus when temperature goes down, soot particles that are not nucleated at a particular temperature have a very little effect on nucleation rate at another temperature.

In addition to the nucleation rates and ambient conditions, the sizes of these IN are also important to our analysis of the thermodynamic parameters for ice nucleation. Some experiments provide exact information of particle sizes and other relevant experimental data, while others lack this type of information. For the missing information we resort to other literatures to obtain estimations. Uncertainties of such estimations will be discussed later. Table 1 gives a summary of the types of IN, as well as their mode of nucleation, representative size and actual size range that will be used for analysis later.

2.2 Classical nucleation theory

Classical nucleation theory is straightforward, but is repeated here for discussion purpose. The rate of heterogeneous nucleation per particle can be expressed as (Mason, 1974, p. 474–478; Hagen et al., 1981; Pruppacher and Klett, 1997, p. 303):

$$J = 4\pi r_N^2 J_s = 4\pi r_N^2 A_1 n_1 \cdot \exp\left(\frac{-\Delta g_g}{kT}\right) \cdot Z, \quad (1)$$

where r_N is the radius of ice nuclei, J_s is the rate of surface nucleation, A_1 is the rate at which the critical embryo gains one molecule through interaction with the parent

Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



phase, n_1 is the number of single molecules in contact with unit area of the substrate, Δg_g is the energy of critical embryo (germ) formation, k is the Boltzmann constant, T is temperature, and Z is the Zeldovich factor which modifies the nucleation rate by accounting for the depletion of embryo population due to germ production (cf. Pruppacher and Klett, 1997, p. 202; Vehkamäki et al., 2007):

$$Z = \frac{1}{n_g} \cdot \sqrt{\frac{\Delta g_g}{3\pi kT}}. \quad (2)$$

Here, n_g is the number of water molecule in the ice germ, defined as:

$$n_g = \frac{4\pi r_g^3}{3v_w}, \quad (3)$$

where v_w is the volume of a water molecule and r_g is the radius of embryo germ (which can be derived according to classical nucleation theory and its definition will be shown later). Equation (1) can be applied to ice nucleation from both the vapor phase (deposition nucleation) and liquid phase (freezing nucleation). Details of the parameters A_1 , n_1 , Δg_g and r_g pertaining to each of these two nucleation modes are given separately below.

2.2.1 Deposition nucleation

For deposition nucleation, the parameters A_1 in Eq. (1) is determined from the kinetic theory (Mason, 1974, p. 474):

$$A_1 = 4\pi r_g^2 \frac{e}{\sqrt{2\pi m_w kT}}, \quad (4)$$

where e is the vapor pressure, m_w is the mass of a water molecule, and the germ size r_g can be expressed as:

$$r_g = \frac{2v_w \sigma_{i/v}}{kT \ln S_i}, \quad (5)$$

in which $\sigma_{i/v}$ is the surface tension between ice and vapor, and S_i is saturation ratio with respect to ice. The second parameter, n_1 , in J_s can be determined under the assumption of a steady state adsorption-desorption fluxes (Pruppacher and Klett, 1997, p. 300):

$$n_1 = \frac{e}{v_s \sqrt{2\pi m_w kT}} \exp\left(\frac{-\Delta g_d}{kT}\right), \quad (6)$$

where v_s is the frequency of vibration of water vapor molecule adsorbed on solid substrate, Δg_d is the energy of desorption. According to classical nucleation theory, the energy of germ (critical embryo) formation for heterogeneous nucleation Δg_g in Eq. (1) is simply the homogeneous nucleation energy of germ formation, Δg_g^o , modified by a geometrical factor f :

$$\Delta g_g = \Delta g_g^o \cdot f \quad (7)$$

where

$$\Delta g_g^o = \frac{16\pi v_w^2 \sigma_{i/v}^3}{3\Delta g_b^2} = \frac{16\pi v_w^2 \sigma_{i/v}^3}{3[kT \ln S_i]^2} = \frac{4\pi}{3} \sigma_{i/v} \cdot r_g^2, \quad (8)$$

and Δg_b is the bulk free energy of phase change. The geometric factor for ice germ formation on a planar surface is a function of the contact angle only:

$$f = f(m) = \frac{(2+m)(1-m)^2}{4}, \quad (9)$$

where $m = \cos(\theta)$ is called the wetting coefficient, θ is the contact angle of ice germ on the substrate. But on a curved surface, it is also a function of the germ size (cf. Fletcher, 1958; Pruppacher and Klett, 1997, p. 302; Liu, 1999):

Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

$$f = f(m, q) = \frac{1}{2} \left\{ 1 + \left(\frac{1 - mq}{\phi} \right)^3 + q^3 \left[2 - 3 \left(\frac{q - m}{\phi} \right) + \left(\frac{q - m}{\phi} \right)^3 \right] + 3mq^2 \left(\frac{q - m}{\phi} - 1 \right) \right\}. \quad (10)$$

where $\phi \equiv \sqrt{1 - 2mq + q^2}$, and $q \equiv r_N / r_g$. When r_N approaches zero or the contact angle is at its extreme ($\theta = \pi$), the geometric factor f approaches unity so the condition reduces to homogeneous nucleation. For r_N approaches infinity, f should reduce to Eq. (9) (i.e. for a planar surface).

2.2.2 Freezing nucleation

For freezing nucleation, Turnbull and Fischer (1949) gave the parameter A_1 as:

$$A_1 = \frac{kT}{h} \exp \left(\frac{-\Delta g_a}{kT} \right), \quad (11)$$

where h is the Plank's constant, and Δg_a is activation energy for the transfer of a water molecule across the water-ice boundary. Similar to Eq. (8), the energy of germ formation for homogeneous freezing is:

$$\Delta g_g^o = \frac{16\pi v_w^2 \sigma_{i/w}^3}{3\Delta g_b^2} = \frac{16\pi v_w^2 \sigma_{i/w}^3}{3[kT \ln(e_{sw}/e_{si})]^2} = \frac{4\pi}{3} \sigma_{i/w} \cdot r_g^2, \quad (12)$$

where $\sigma_{i/w}$ is the surface tension between ice and liquid water, and e_{sw} and e_{si} are the saturation vapor pressure over water and ice, respectively. Note that for the freezing of solution, the parameter e_{sw} in Eq. (12) should be modified by the water activity of the solution (cf. Mason 1974, p. 491; Heymsfield and Sabin 1989). In contrast to that for deposition nucleation, the parameter n_1 for freezing nucleation does not vary significantly. For a water density of 1 g cm^{-3} , one can easily calculate that $n_1 = 1.0 \times 10^{19} \text{ m}^{-2}$, which is the same as that given in Fletcher (1962) but much larger than the $5.3 \times 10^{16} \text{ m}^{-2}$ given by Eadie (1971; also see Pruppacher and Klett, 1997, p. 206). The geometric factor f maintains the same form as in Eqs. (9) and (10) for both

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



nucleation modes. Another parameter that needs to be defined for freezing nucleation is the size of the ice germ:

$$r_g = \frac{2v_w\sigma_{i/w}}{kT \ln(e_{sw}/e_{si})}. \quad (13)$$

3 Linearization analyses of classical theory

5 The nucleation rate given in Eq. (1) consists of two groups of parameters; (A) ambient variables such as air temperature (and temperature dependent variables), vapor pressure and ice supersaturation; (B) particle variables, including particle size r_N , the activation energy (Δg_d for deposition nucleation or Δg_a for freezing nucleation), and the contact angle θ (or the wetting coefficient m). Existing laboratory studies provided
 10 the nucleation rates and the corresponding environment variables of group (A). But the particle parameters of group (B), particularly the activation energy and contact angle, are virtually unknown for common atmospheric IN. Our main task is to derive these parameters from laboratory measurements. This can be done by first rearranging Eq. (1) into:

$$15 \quad J = A' \cdot r_N^2 \cdot \sqrt{f} \cdot \exp\left(\frac{-\Delta g^\# - \Delta g_g}{kT}\right), \quad (14)$$

where A' depends on ambient parameters only; $\Delta g^\#$ is either Δg_d in Eq. (6) or Δg_a in Eq. (11), and they will be generally termed the activation energy for future discussion. One can see that the rate equation for both nucleation modes can be expressed in the same form. For deposition nucleation, $\Delta g^\# = \Delta g_d$ and

$$20 \quad A' = \frac{2\pi(r_g e)^2}{m_w kT v_s n_g \sqrt{3\pi}} \cdot \sqrt{\frac{\Delta g_g^o}{kT}} = \frac{e^2 v_w}{m_w kT v_s} \cdot \sqrt{\frac{\sigma_{i/v}}{kT}}. \quad (15)$$

Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



For freezing nucleation, $\Delta g^\# = \Delta g_a$ and

$$A' = 4\pi \cdot \frac{1}{n_g} \cdot \sqrt{\frac{\Delta g_g^o}{3\pi kT}} \cdot \frac{kT}{h} \cdot n_1 \quad (16)$$

The rate Eq. (14) contains three parameters that are properties of the ice nuclei: particle size r_N , activation energy $\Delta g^\#$, and wetting coefficient m (or contact angle). With a few sets of nucleation rate and ambient condition data that measured in the laboratory, it is possible to derive these particle parameters from Eq. (14) particularly when the particle size is known. In the following, we intent to linearize Eq. (14) and perform data fitting to derive the two parameters $\Delta g^\#$ and m .

3.1 Initial analysis

Recall from Eq. (10) that the geometric factor f is a complicated function of nuclei size, contact angle, and air properties. But under typical conditions, its primary dependent variable is the contact angle. So we shall first assume it is a constant, taking the form of Eq. (9), and extend the analysis later for an exact f . By taking the logarithmic of Eq. (14), we have

$$\underbrace{kT \ln \left(J/A' / r_N^2 \right)}_y = \underbrace{kT \ln \sqrt{f}}_a - \underbrace{\Delta g^\#}_b - \underbrace{f \cdot \Delta g_g^o}_x \quad (17a)$$

This formula is arranged in such a way that x and y can be directly determined from the experimental data, whereas the particle-related parameters are kept in the coefficients a and b . Therefore, by fitting the experimental data with the linear equation $y = a - bx$, we are able to derive the coefficients a and b , and from which we can obtain the particle parameters. For example, using Eq. (17a), the geometric factor f can be derived immediately as it is exactly the coefficient b . However, the coefficient a contains air temperature and thus is not truly a constant, and the uncertainties contained in it will

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



inevitably pass to the value of b . This also means that we are not able to obtain directly the second parameter, the activation energy Δg^\ddagger , from the coefficient a .

To derive the value of the activation energy, we can take a variation of Eq. (17a) as:

$$\underbrace{kT \ln (J/A') + f \cdot \Delta g_g^o - kT \ln \sqrt{f}}_y = \underbrace{-\Delta g^\ddagger}_a + \underbrace{\ln r_N^2}_{-b} \cdot \underbrace{kT}_x. \quad (17b)$$

5 In this formula, we applied into y the previously derived coefficient f as an initial guess. This gives us not only the value of Δg^\ddagger but also the nuclei size which we use to indicate the robustness of our method. Of course, all values derived so far still contain the previously mentioned uncertainties. Some of the uncertainties can be minimized by introducing another variation of Eq. (17a):

$$10 \underbrace{\ln (J/A') + \Delta g^\ddagger / kT}_y = \underbrace{\ln (r_N^2 \sqrt{f})}_a - \underbrace{f}_b \cdot \underbrace{\Delta g_g^o / kT}_x. \quad (17c)$$

Here, we can apply the Δg^\ddagger obtained from Eq. (17b) into y to re-derive f . Further calculation by iteration between the two procedures (Eqs. 17b and 17c) has been tried, but the results do not converge. But this is not a problem because we only want to obtain an initial estimation and get a sense of the order of magnitude of the desired parameters. More detailed analysis will be performed in the next section.

15 Table 2 shows the results obtained by a simple iteration from Eqs. (17a) to (17c) for the species shown in Table 1. One can see that the two values of f are quite close to each other, but the calculated r_N may differ significantly from the originally assigned values for some IN species. The discrepancy in r_N arises either from the uncertainties in the formulation procedure or due to the assumption of constant f . In the next section we will try to minimize both.

Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



3.2 Refined geometric factor

In the above analysis, we have assumed that f takes the form of Eq. (9) and is independent of the ambient parameters. But in fact f is also a function of particle size r_N and the germ size r_g as given in Eq. (10). For a more accurate analysis we need to consider these dependences. But the original form of Eq. (10) is too complicated to be decomposed for the analysis done in the previous section. This problem can be circumvented by fitting Eq. (10) into either of the following two equations, which separate out particle size (contained in q) and contact angle:

$$\ln f \approx c_1 + c_2 \cdot \ln(1 - m) + c_3/q. \quad (18a)$$

$$\ln f \approx c_4 + c_5 \cdot \ln(1 - m) + c_6 \cdot \ln q, \quad (18b)$$

The coefficients of the above equations are given in Table 3 for the species of IN considered in this study under appropriate ranges of q and θ . The coefficients of determination (R^2) for these fittings are all greater than 0.98, and their maximum errors are quite small except for ATD.

By applying Eq. (18a) into Eqs. (17a) and (17b), we obtained two variations of fitting on the nucleation rate formula as done in the previous section but without assuming f as a constant:

$$\underbrace{kT \left\{ \ln \left(J/A' / r_N^2 \right) - 0.5 \cdot (c_1 + c_3 \cdot r_g / r_N) - 0.5 \cdot c_2 \cdot \ln(1 - m) \right\}}_y \quad (19a)$$

$$= \underbrace{-\Delta g^\#}_a - \underbrace{(1 - m)^{c_2}}_b \cdot \underbrace{\Delta g_g^o \cdot \exp(c_1 + c_3 \cdot r_g / r_N)}_x$$

$$\underbrace{\ln \left(J/A' / r_N^2 \right) + \Delta g^\# / (kT) - 0.5 \cdot (c_1 + c_3 \cdot r_g / r_N)}_y \quad (19b)$$

$$= \underbrace{0.5 \cdot c_2 \cdot \ln(1 - m)}_a - \underbrace{(1 - m)^{c_2}}_b \cdot \underbrace{\Delta g_g^o / (kT) \cdot \exp(c_1 + c_3 \cdot r_g / r_N)}_x$$

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

In both equations, the coefficients a and b no longer contain ambient parameters but are purely functions of the particle parameters Δg^\ddagger (activation energy) and m (cosine contact angle). However, the independent variable y also contains these supposedly unknown parameters. So, we need an initial guess of either m or Δg^\ddagger in order to solve Eqs. (19a) and (19b) iteratively. But now it is not necessary to go through a similar procedure as Eq. (17a) to get the initial guess because the results in Table 2 already suffice the purpose. The calculations converged quickly within three iterations after including more accurate treatment of the factor f . Note that we also applied Eq. (18b) into Eq. (17b) to obtain a fitting formula in the form of:

$$\underbrace{\ln \left(\frac{J/A'}{y} + \frac{\Delta g^\ddagger}{kT} \right) - 0.5 \cdot (c_4 - c_5 \ln r_g)}_y$$

$$= \underbrace{\ln r_N^2 + 0.5 \cdot [c_5 \cdot \ln(1 - m) + c_6 \cdot \ln r_N]}_a - \underbrace{(1 - m)^{c_5} \cdot r_N^{c_6}}_b \cdot \underbrace{\frac{\Delta g^o}{kT} \cdot \exp(c_4) \cdot r_g^{-c_6}}_x \quad (19c)$$

This fitting formula, when used in conjunction with either Eq. (19a) or Eq. (19b), gives very similar results as before. Furthermore, it also gives the nuclei size which we use to check the self-consistency. The refined results are given in Table 4. We can see that the values of Δg^\ddagger differ slightly from those in Table 2 for all species except dusts, for the latter the deviation may reach 90% (Saharan dust). The differences in m behaved similarly, with the largest discrepancy of about 40% occurred in the species ATD. These large errors signify the importance of a more detailed treatment of the factor f . Note that the calculated r_N are now very close to their assigned values, and this is a good indication of self-consistency of our method.

3.3 Curvature and solute effects on surface tension

When extending our analysis on the data of Hung et al. (2003) we realized that our previous treatment was not comprehensive enough. Their experiment considered the freezing of ammonium sulfate solution with mineral dust as the nucleation center. So

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



we included the solute effect on e_{sw} in Eq. (12) by multiplying it with the water activity using the empirical formula in Chen (1994). Note that in Hung et al. (2003) the measurements for hematite (Fe_2O_3) were done in a rather narrow temperature range, but their different dust sizes give extra variability that benefits our analysis. However, for corundum (Al_2O_3) there is only one measurement temperature, which makes the fitting analysis difficult and no reasonable result was obtained. Our calculation shows that the contact angle of ice germ on hematite ranges from 109.2° at low solute concentration (mole fraction $\chi=0.03$) to 74.4° at high solute concentration ($\chi=0.13$). Similar to our approach, Hung et al. (2003) applied measurement data into classical theory to find an optimized value of contact angle, except that they apparently neglected the Zeldovich factor in Eq. (1) as well as the activation energy term and other details that will be discussed later. The contact angles they obtained were 90° or 97° , depending on the different criteria they set. Although the two calculations seem to be reasonably close to each other, these contact angles are significantly larger than for other types of freezing IN shown in Table 4. As will be discussed later, contact angle is the most important factor in determining the nucleation capability of IN. The large angles derived above would imply a poor nucleating capability of hematite that is even inferior to soot. Such a result seems to be at odds with the perceptions that mineral dusts typically have higher threshold temperature of nucleation than soot, and this prompted us to go back over our analysis.

The first missing factor that we found is the effect of solute on the surface tension of water, which strongly influences the germ formation energy and germ size. Unfortunately, no data exist for the surface tension between ice and water solution. We overcome this by applying the Antonoff's rule, which states that the surface tension between a solid and liquid ($\sigma_{i/l}$) can be approximated by the difference between the surface tension of solid-air interface ($\sigma_{i/a}$) and that of liquid-air interface ($\sigma_{l/a}$):

$$\sigma_{i/l} \approx \sigma_{i/a} - \sigma_{l/a}, \quad (20)$$

where the subscript “ l ” represents a solution to contrast with the subscript “ w ” that we used to for pure water. The values of $\sigma_{i/a}$ can be found in various literatures, whereas

Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

the value of $\sigma_{i/a}$ for the solutions of ammonia sulfate and several other solute species can be obtained from Chen (1994). As $\sigma_{i/a}$ increases with increasing ammonium sulfate concentration, surface tension between ice and solution ($\sigma_{i/a}$) becomes lower under the presence of solute. Table 5 listed the calculated thermodynamic parameters with and without the consideration of solute effect on surface tension. Unfortunately, under a solute mole fraction of $\chi=0.03$, the value of contact angle did not become smaller as hoped; whereas for the high solute concentration the wetting parameter m even exceeded its lower bound of -1 , thus no contact angle can be obtained. So, the inclusion of solute effect leads us in the wrong direction.

Another important factor that may influence contact angle is the curvature adjustment of surface tension on ice germs. Tolman (1949) suggested that surface tension deviates from its bulk value when the particle is small, and the size adjustment follows $\sigma'_{A/B} = \sigma_{A/B}^{\infty} \cdot \phi$, where σ^{∞} is the bulk surface tension, and subscripts A and B represent the particle phase (here, ice germ) and the medium phase (here, water or air), respectively. The proportionality coefficient can be expressed as (also see Dufour and Defay 1963, p. 61):

$$\phi = \frac{1}{1 + 2\Gamma/r_A(\rho_A - \rho_B)} \equiv \frac{1}{1 + 2\delta/r_A}, \quad (21)$$

where r_A is the particle size, ρ is the density of the phase that indicated by its subscript, $\Gamma = 1.55 \times 10^{-7} \text{ kg m}^{-2}$ is the Gibbs surface adsorption, and

$$\delta \equiv \Gamma / (\rho_A - \rho_B). \quad (22)$$

Note that this equation is equally applicable when the particle is of phase B , for which case a negative radius (curvature) is used for r_A (i.e. treat it as a “bubble”). However, Eq. (21) gives unrealistic (negative) value of ϕ when $(\rho_A - \rho_B)$ is a small negative value (for the case of ice in water) and the particle size is small. Lu and Jiang (2004) provided

a more general formula of curvature adjustment for nanocrystals in the gas phase:

$$\phi = \left(1 - \frac{1}{4r/\delta - 1}\right) \cdot \exp\left(-\frac{2E}{3RT} \cdot \frac{1}{4r/\delta - 1}\right), \quad (23)$$

where E is the bulk energy of phase transformation and R is the gas constant. We extended the use of their formula to other particle-medium systems by applying Eq. (22) for the factor δ , and found that it can be safely applied to all our situations. Besides influencing contact angle, curvature adjustment may also affect the germ size and energy of germ formation. From Eqs. (5) and (13) we see that the germ size is directly proportional to the surface tension, so it should be modified accordingly with

$$r'_g = r_g^\infty \cdot \phi, \quad (24)$$

where r_g^∞ is the bulk germ size given by Eqs. (5) or (13). However, the factor ϕ itself is a function of the germ size as can be seen in Eq. (23). Thus, r'_g and ϕ need to be solved simultaneously (e.g. by iteration) from Eqs. (23) and (24).

From Eq. (23) one can see that the curvature adjustment depresses both the surface tension and size of ice germ that is embedded in air (e.g. deposition nucleation), but enlarge them if the surrounding medium is water (e.g. freezing nucleation). As shown in Table 5, the resulting contact angle of 69.4° at $\chi=0.03$ has shifted toward the desirable direction. Furthermore, the originally invalid value at $\chi=0.13$ now becomes a much more reasonable 63.4° . But these contact angles are still larger than that of soot. In the next section we will further elaborate the inconsistency of these values with other data.

In Table 6 we recalculated the contact angle and activation energy with the consideration of curvature adjustment for all species, and compare them with the old values shown in Table 4. One can see that the curvature adjustment generally results in smaller contact angles for freezing nucleation and inversely so for deposition nucleation. As for the activation energies the influence is less prominent. Those values calculated with curvature adjustment can be regarded as the final result of this study.

Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Ice nucleation rate
from contact angle
and activation energy**

J.-P. Chen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Take Eqs. (17a) and (19a) as examples, we show the effects of linearization on the distribution of experimental data in Fig. 1. Figure 1a uses Eq. (17a) but neglects curvature adjustment; Fig. 1b and 1c are both from Eq. (19a) but without and with the curvature adjustment, respectively. From Fig. 1a one can see that Eq. (17a) already produced fairly good linearization. The curves are grouped naturally according to the general species types (i.e. bacteria, pollens or soot), and the distinct slope reflects the wetting parameter f that is unique for each species. Results from Eq. (19a) are not much different in terms of the degree of linearization. But more importantly, these curves give more accurate values of contact angle and activation energy. The effect of curvature adjustment is shown in Fig. 1c as oppose to Fig. 1b. One can see significant changes in the values of x and y , resulting in substantial changes in the slope (mainly representing the contact angle) and intercept (mainly representing the activation energy). Note that the inclusion of curvature adjustment also slightly improved the R^2 of linear regression.

4 Discussions

4.1 Relative significance of the activation energies and contact angles

Table 6 can tell us about the ice nucleating ability of various species. We can see that activation energy Δg_a for freezing nucleation is the highest for pollens and slightly lower for bacteria and soot, but the differences between species and, particularly, between sub-species are not significant except for hematite. However, all the differences in $\Delta g^\#$ are too small to account for the vastly different nucleation capability among species. By examining Eq. (14) one may realize that $\Delta g^\#$ (either Δg_a or Δg_d) is not a major factor in determining the nucleation rate because there is another stronger energy requirement for nucleation – the energy of germ formation Δg_g . Typically, Δg_g are of the order of 10^{-16} to $10^{-18} J$, about two to three orders of magnitude larger than the activation energies. Only under very low temperature or high supersaturation the

activation energy becomes important. Another interesting contrast is that changes in the activation energy are relatively smaller for the freezing mode than for the deposition mode nucleation. This implicates that the adsorption energy of vapor (Δg_d) is sensitive to the composition or surface structure of the substrate, whereas the activation energy for the transfer of a water molecule across the water-ice boundary (Δg_a) is not.

Let us see if the difference in contact angles may reflect the nucleation capability. Bacteria have significantly smaller contact angles (see Table 6), and this indeed reflects the fact that bacteria are better IN than the other freezing mode species as shown by the nucleation rate data or the so-called threshold temperature. *Pseudomonas aeruginosa* obviously is a more effective IN because of its smallest contact angle. Soot seems to have a lower (but only slightly) activation energy, yet it has a rather large contact angle. Exacerbated by its small size, soot apparently is not a very effective IN as can be seen from Eq. (1) for the size dependence of the nucleation rate. Pollens have the advantages of large sizes and relatively small contact angles, and thus they are relatively efficient ice nucleating agents. Among the six types of pollens, pine pollen has the smallest contact angle and largest size thus it should have better nucleation ability than other pollens, and this is confirmed by the experimental data. Hematite has the largest contact angle thus its freezing nucleation capability is even lower than soot.

For the three deposition-mode dust species, the contact angles are rather small, particularly for the Arizona test dust and next to it the Asian dust, indicating mineral dusts are good nucleation agent, at least by deposition nucleation. We are not able to explain why the contact angle for freezing on a seemingly similar mineral dust – hematite – is much larger, except that hematite might not represent common mineral dust in the atmosphere.

4.2 Verification of results

There are very little direct measurements on the activation energies for the types of IN and nucleation conditions considered in this study, and essentially none for the contact angles. Therefore, it is difficult to get a direct comparison and verification for the results

Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



shown in Table 6. However, by comparing with similar data one may get a sense of the correctness of our findings. We first look at Δg_a , the activation energy for the transfer of a water molecule across the water-ice boundary.

Hagen et al. (1981) applied an approach similar to ours to analyze measurement data of nucleation rate, but for homogeneous freezing, to derive Δg_a . They obtained 8×10^{-20} J (per H₂O molecule) for Δg_a at -40°C , and this value roughly increases by 0.3×10^{-20} per degree K of temperature rise. If their calculation can be extrapolated to the temperature range of the experimental data, it would be similar to the values calculated here, which lie between 10×10^{-20} and 16×10^{-20} J. Pruppacher and Klett (1997, Table 7.3) also calculated Δg_a but found somewhat lower values: 3×10^{-20} J at -36°C and increases to 5.4×10^{-20} J at -29°C . Note that the activation energy Δg_a might have different controlling mechanisms at different temperatures. Hagen et al. (1981) suggested that at temperatures warmer than -29°C , the transfer of water molecules across the water-ice interface is limited by the self-diffusion through the bulk water, while at temperatures lower than -32°C the transfer is by large water clusters.

The values of $\Delta g^\#$ that we derived should be considered as averages over the temperature range of each experimental data set. In principle, one should be able to get the temperature dependence by inserting the parameters r_N and contact angle from Table 6 along with the measured nucleation rate into Eq. (14) to get $\Delta g^\#$ at each measured temperature. Of course, this has to be done under the assumption that m is a constant with respect to temperature. Figure 2 shows the activation energies that were calculated this way. Within each species, the variation of Δg_a with temperature (Fig. 2a) is not significant comparing to the theoretical estimations of Hagen et al. (1981) and Pruppacher and Klett (1997) for homogeneous nucleation. This suggests either that our method and data are not sufficient to resolve it or that there exist other factors that diminish the temperature dependence. One possible cause for the difference is that the factor m also depends on temperature, and neglecting this somehow offset the temperature dependence of Δg_a . Thus, we tried to fix $\Delta g^\#$ as an alternative to calculate m for each data point but found that m calculated this way is nearly independent of

Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



temperature. It is also possible that the presence of IN substrate regulated the value of Δg_a so its behavior deviates from that for homogeneous nucleation. As can be seen from Fig. 2a, the differences in Δg_a among species are greater than for single species at different temperatures.

5 For the deposition nucleation mode, the activation energy is the energy of adsorption of water vapor onto the IN substrate but represented by the energy of desorption Δg_d . Since the adsorption process strongly depends on the properties of the substrate, one would expect the value of Δg_d to change more significantly among species as compared to the changes in Δg_a . Indeed, our values of Δg_d vary by more than one fold,
10 even though they are all for mineral dusts, whereas those for freezing nucleation vary within 40% among different species. Some direct measurements of Δg_d can be found in the literature for comparison. Seisel et al. (2004) found the adsorption energy of water vapor on mineral dust to be 6.6×10^{-20} J; Hu and Michaelides (2007) obtained
15 1×10^{-19} J for water adsorption on Kaolinite; whereas Gustafsson et al. (2005) obtained 7.6×10^{-20} J for calcite and 8.7×10^{-20} J in the case of Arizona Test Dust. Although our values are of the right orders of magnitude, they are somewhat lower than those found in the literature. Note that the adsorption energy is controlled by not only the chemical composition but also the physical properties such as roughness and defects of the substrate. This implies that the way the dust samples are processed in the laboratory may influence the value of Δg_d . Also note that, as shown in Fig. 2b, we did not find an obvious temperature trend for Δg_d . The following are a few measurement data on adsorption energy which might be useful for relevant atmospheric applications:
20 4.9×10^{-20} J on grey soot and 6.3×10^{-20} J on black soot (Alcala-Jornod et al., 2002), and 1.5×10^{-19} J for Agl (Corrin and Nelson, 1968). One can see that the difference between species is pretty much the same as that between sub-species of dusts.
25

Contact angle is more important in determining the heterogeneous nucleation ability. Unfortunately, most of the past efforts were done for the contact angle of liquid water as enclosed by air and the substrate, which is conventionally written as $\theta_{w/a}$. Yet, the contact angles that we are considering here are for ice, as enclosed either by air and

Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



substrate ($\theta_{i/a}$; for deposition nucleation) or by liquid water and substrate ($\theta_{i/w}$; for freezing nucleation). The only attempt that we are aware of to find the contact angle of ice is the study of Hung et al. (2003), but those values were not measured directly but were obtained by a similar analysis as ours. Here we introduce an indirect way of verifying the results on contact angle. According to the Young-Dupré relation, the contact angles can be expressed as:

$$\cos \theta_{w/a} = \frac{\sigma_{s/a} - \sigma_{s/w}}{\sigma_{w/a}}, \quad (25)$$

$$\cos \theta_{i/a} = \frac{\sigma_{s/a} - \sigma_{s/i}}{\sigma_{i/a}}, \quad (26)$$

$$\cos \theta_{i/w} = \frac{\sigma_{s/w} - \sigma_{s/i}}{\sigma_{i/w}}, \quad (27)$$

where the subscripts represent the interface between any two of the following phases: liquid water (w), ice (i), air (a) or substrate (s). A closure of the equation set Eqs. (25) to (27) can be obtained by using Eqs. (26) and (27) to replace $\sigma_{s/a}$ and $\sigma_{s/w}$ in Eq. (25) and obtain:

$$\cos \theta_{w/a} = \frac{\cos \theta_{i/a} \cdot \sigma_{i/a} - \cos \theta_{i/w} \cdot \sigma_{i/w}}{\sigma_{w/a}}. \quad (28)$$

In this equation the values of $\sigma_{w/a}$, $\sigma_{i/w}$ and $\sigma_{i/a}$ (with temperature dependence) can be found in many textbooks (e.g., Pruppacher and Klett, 1997). Then, using values of $\theta_{i/a}$ from the literature and values of $\theta_{i/w}$ obtained here, one may derive $\theta_{w/a}$ from Eq. (28).

If one assumes that hematite is similar to the other mineral dusts, then the freezing contact angle $\theta_{i/w}$ of hematite and the deposition contact angle $\theta_{i/a}$ of ATD, AD or SD given in Table 6 can be used to calculate $\theta_{w/a}$ according to Eq. (28). There are quite a few measurements on $\theta_{w/a}$ that can be used for comparison. For instance,

Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Ice nucleation rate
from contact angle
and activation energy**

J.-P. Chen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Bergh (1965) observed a contact angle of less than 6° on SiO_2 , but it increases to 40° after heat treatment of the substrate; Chen et al. (1993) measured 33° on SiO_2 and 57° on Al_2O_3 ; Janczuk and Zdziennicka (1994) measured 26.8° on quartz; Pruppacher and Klett (1997, Table 5.2) reported $43\sim 52^\circ$ on quartz (beach sand); whereas Gence (2005) reported 10.5° on magnesite and 6.7° on dolomite. One can see that there is a large spread in the values of water contact angle on different minerals. In any case, to derive from Eq. (28) a reasonable value of $\theta_{w/a}$ within the range reported above, $\theta_{i/w}$ must be smaller than $\theta_{i/a}$. Yet, the results in Table 6 show just the opposite. Note that $\theta_{i/w}$ would be even larger when extrapolated to pure water situation. The above inconsistencies indicate that either we missed some aspects of the ice germ calculation or the hematite particle used by Hung et al. (2003) behave very differently from the other dust species considered here. If the latter is true, then it is still feasible to apply the above method to derive one contact angle with the knowledge of the other two. Of course, experimental data on both the deposition mode and freezing mode nucleation for the same IN species would be necessary for verification. It is also possible that the data of Hung et al. (2003) were measured over a temperature range that is too narrow for an accurate fitting with our method, or the formula for curvature adjustment (Eq. 23) is not accurate enough.

4.3 Application to other nucleation modes

In Sect. 4.2 we discussed the possibility of using the contact angles in Table 6 to derive various surface tension parameters so that the results for freezing nucleation (or commonly called immersion freezing) can be applied to deposition nucleation or vice versa. It would also be interesting to know whether the results obtained here can be applied to the other two modes of heterogeneous nucleation: condensation-freezing nucleation and contact nucleation. Condensation-freezing nucleation can be treated as a two-step process: condensation nucleation followed by the immersion freezing nucleation. Without the presence of salt (solute), the occurrence of condensation nucleation requires at least some supersaturation with respect to water. Its rates can be

**Ice nucleation rate
from contact angle
and activation energy**J.-P. Chen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

calculated easily following formulations similar to those given in Sect. 2 and by know-
ing the adsorption energy and contact angle $\theta_{w/a}$, so the details will not be elaborated
here. Due to the extra step of condensation nucleation, IN that have good freezing ca-
pability might not be good condensation-freezing nuclei. In this regard, pollens could
5 be good IN for condensation-freezing nucleation because they are also efficient in con-
densation nucleation, partly due to their large sizes and partly because the capillary
effect on pollens allows rather easy water uptake even under sub-saturated environ-
ment (Diehl et al., 2001). The presence of solute could also help the uptake of liquid
water thus enhances the condensation nucleation but, in the mean time, depresses the
10 probability of freezing nucleation.

The remaining heterogeneous ice nucleation is the contact nucleation, whose rate
is difficult to measure in the laboratory. Cooper (1974) suggested that the contact
nucleation could be calculated in a manner analogous to that for immersion freezing
and deposition freezing. Of course, the probability of making contact also needs to
15 be considered. Cooper's theoretical approach seems to be a viable way of dealing
with this potentially important nucleation mechanism. Therefore, the thermodynamic
parameters obtained in this study in effect can be used on all four ice nucleation modes.

4.4 Error sources

Even though it has been proven successful against laboratory studies, the classical
theory has several apparent deficiencies. For instance, we used many bulk thermo-
dynamic properties to describe the behavior of microscopic ice germs. Caution must
20 be made that the contact angle is a macroscopic concept, which might not hold for the
minuscule nucleation germs. Most of the contact angles mentioned in the literature
were measured on a macroscopic scale. As Evans and Lane (1973) pointed out, there
is a marked difference between the macroscopic contact angle and that of the ice germ
due to the line tension effect. Thus, the contact angles derived here probably should
25 be viewed as an equivalent thermodynamic quantity. Similar concerns also exist in the
specific volume (density) and surface tension terms that appeared in Eqs. (5), (8) and

(12).

Inherent uncertainties also arise from some of the basic assumptions of the classical theory, such as the Boltzmann distribution and quasi-steady state assumption of germ concentration. The latter assumption can be corrected with the Zeldovich factor, but the factor itself was derived with mathematical approximations. Fortunately, the error involved is about 1% (Pruppacher and Klett, 1997, p. 202).

Some error of estimation may also arise from the assumptions that we made. In Eq. (8) the bulk free energy of phase change Δg_b in the denominator may be affected by elastic strain ε between the lattices across the interface (Turnbull and Vonnegut, 1952):

$$\Delta g_b = kT \ln S_i - v_w \cdot C \cdot \varepsilon^2, \quad (29)$$

where C is about $1.7 \times 10^{10} \text{ J m}^{-3}$. In previous discussions, we have ignored this term because very little is known about the values of ε . However, one may wonder how significantly the elastic strain affects our results. The strain is typically less than a few percents between ice and most other materials. Assume it is on the order of 1%, this would decrease the bulk free energy Δg_g by less than 15% when the ice saturation ratio S_i is above 1.1. The same estimation can be made on Eq. (12) for the freezing nucleation mode. The error from neglecting the strain term would cause a similar error in the factor f but no obvious effect on the activation energies. However, as the contact angle and m vary slowly with f for the conditions considered here, we expect the error associated with the elastic strain effect to be less than 15%. Other physical properties such as defects or dislocations that may enhance the nucleation capability also contribute to similar uncertainties.

A more complicated problem is the uncertainties in r_N . In natural environment, bacteria, pollens, soot and dust particles all have large variations in r_N . The variations might be smaller for the laboratory particles mentioned in Sect. 2.1, which were pre-processed before applying to the measurement of nucleation rates. However, the particles will never reach a mono-dispersed size distribution as we have implicitly assumed.

**Ice nucleation rate
from contact angle
and activation energy**

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Ice nucleation rate
from contact angle
and activation energy**

J.-P. Chen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Some of the measurements did provide particle size distribution, for which we took the modal size as the assigned r_N . But for those that did not provide size information, we have to make an educated guess according to the information provided in the published literature. Either way, the assigned r_N must contain uncertainties. We estimated the uncertainties associated with the assumed particle size by the following analysis. For each IN species, we made two more sets of calculations, one assuming the mean particle size is one-third of the assigned r_N , and the other is three times of the assigned r_N . The results are given in Table 7. One can see that, within this nearly one order of magnitude variation, the uncertainties in the wetting parameter m and contact angles are rather limited, except for the contact angle of ATD, which might change by as much as 18% for the smaller r_N . The activation energies $\Delta g^\#$, however, are more sensitive to the assigned value of r_N , particularly for the three dust species the values may change by over 100%. Fortunately, for the pollen species, which we are most unsure of the sizes used in the laboratory experiments, the results are quite consistent. Furthermore, the nucleation rate is not very sensitive to the variations in activation energy as discussed earlier.

Our analysis also included several idealizations that might have introduced uncertainties. For example, all IN particles were assumed spherical but in reality, few really are. In addition, particles of each IN species were assumed to have the same surface physical properties (roughness, defects, impurities, active sites, etc.), but in the real world or even in the laboratory experiments variations must exist from particle to particle. Variations in the surface chemical properties may also exist among IN of the same species. Deactivation of IN may result when they have been in contact with foreign gases or particles (Georgii and Kleinjung, 1967). Gorbunov et al. (2001) also suggested that oxidation might affect the concentration of soot's surface chemical groups thus change their ice nucleating capability, implying that not all soot should be treated the same. Unfortunately, the errors associated with these factors are difficult to evaluate.

With many of the above assumptions or uncertainties, it might be more appropriate

to call the results derived from the classical theory the apparent parameters, such as the “apparent activation energy” and “apparent contact angle.” Thus, one may regard the neglected factors, such as the elastic strain effect, to have been embedded into the contact angle in our analysis.

5 Besides evaluating basic thermodynamic properties for various IN, our analysis also provide a way to broaden the application of laboratory results. For use in cloud microphysical models, laboratory results on ice nucleation are often converted into rate formulas by simple curve fitting techniques. These formulas are empirical in nature and not easy to modify to include additional factors that were not considered in the
10 experiments. By extracting the contact angle and activation energy, the empirical rate formulas can be transformed into the classical nucleation rate equation given in Eq. (1), which allows the flexibility of revising various basic parameters. For example, one can include solute effect by modifying the energy of germ formation given in Eq. (8), so that the original experiment done for the heterogeneous freezing of pure water may be
15 extended for application to solution freezing. A similar treatment can also be done for the curvature effect. In laboratory measurements, such as that of Möhler et al. (2006) and Field et al. (2006), the size range of IN that applied might deviate from that of naturally occurring IN. As discussed in Sect. 3.3, curvature effect may have strong influence on the nucleation rate. Therefore, rates measured for one size might not be
20 adequately applied to other sizes. The present approach could be easily used to solve such problems.

5 Conclusions

Comprehensive representations of heterogeneous ice nucleation rate were achieved by a combination of classical theory and experimental data, from which we derived
25 the activation energy and contact angle for various ice nuclei (including soot, bacteria, pollen, and mineral dust) that are important to the formation of ice in clouds. When these thermodynamic parameters are established, one will be able to describe ice

Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



nucleating behavior in clouds pertaining to a particular species of ice nuclei with the knowledge of their size distributions.

The activation energy for freezing nucleation does not vary much among different IN species, whereas larger variability exists in the activation energy for deposition nucleation even among similar types of IN (e.g. three types of mineral dusts). Furthermore, the latter are about an order of magnitude lower than the former. It is important to note that under most circumstances, the activation energy is not critical to the calculation of nucleation rate.

Contact angle is a key parameter in determining the nucleation capability of IN. Among the IN species that are analyzed here, bacteria have the smallest contact angle of ice germ and are thus the most effective in initiating freezing nucleation. Pollens also have relatively small contact angle of ice-germ formation. In addition, they also have the advantage of large sizes and water absorbing surface for acquiring liquid water, thus are efficient in ice nucleation via both freezing and condensation-freezing. Soot has relatively large contact angle thus is less efficient in freezing nucleation. One type of mineral dust (hematite) analyzed here show even larger contact angles for freezing nucleation. However, for deposition nucleation, three other types of mineral dusts show rather small contact angle thus are effective IN.

Our detailed analyses considered the Zeldovich factor for modification of ice germ formation, the particle-size dependent geometric factor for the calculation of wetting coefficient, as well as the solute and curvature effects on surface tension, thus the value of contact angle and energy of ice germ formation. All these factors to various degrees influence contact angle, and should be taken into consideration in the theoretical calculation of nucleation rates. We also discussed a possible closure calculation connecting the three surface tensions in the ice-substrate-medium system. With these calculations, the contact angle for deposition nucleation can be used to obtain the contact angle for freezing nucleation or vice versa. Unfortunately, current data are not sufficient for verification of these calculations.

Considering the microscopic nature of the ice germ that may deviate from that mea-

Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



sured in the bulk, as well as various uncertainties involved in our analysis, it might be more appropriate to view the contact angle and the activation energy derived here as the “apparent” parameters. When applied to theoretical formulations, these apparent parameters should nevertheless give nucleation rates close to that from laboratory measurements.

Our approach makes laboratory data more useful because factors (e.g. solute effect or curvature effect) that were not considered in the experiment can still be incorporated into the rate equations. Furthermore, the same equation can be easily applied to different IN and different nucleation modes by simply changing the relevant thermodynamic parameters (e.g. contact angle and activation energy). It is also desirable to apply our method in future laboratory work to analyze the measurement data so as to gain more thorough understanding of the physical mechanisms of ice nucleation process. Additional asset may be gained if the laboratory measurements could be performed on both the deposition and freezing modes of the same IN species so that a closure analysis of the contact angles and surface tensions between different phases may be obtained. This approach can not only provide more physical insight but also serve for result verification.

Acknowledgements. This research was supported under Grants NSC 96-2475-M-002-009 and 97-2111-M-002, as well as Grant NSC 96-2752-M-002-013-PAE.

References

- Archuleta, C. M., DeMott, P. J., and Kreidenweis, S. M.: Ice nucleation by surrogates for atmospheric mineral dust and mineral dust/sulfate particles at cirrus temperatures, *Atmos. Chem. Phys.*, 5, 2617–2634, 2005, <http://www.atmos-chem-phys.net/5/2617/2005/>.
- Alcala-Jornod, C., van den Bergh, H., and Rossi, M. J.: Can soot particles emitted by airplane exhaust contribute to the formation of aviation contrails and cirrus clouds?, *Geophys. Res. Lett.*, 29, pp. 1–1, CitelID 1820, doi:10.1029/2001GL014115, 2002.
- Barth, M., McFadden, J. P., Sun, J., et al.: Coupling between Land Ecosystems and the At-

Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ospheric Hydrologic Cycle through Biogenic Aerosol Pathways, B. Am. Meteorol. Soc., 86, 1738–1742, 2005.

Bergh, A. A.: The correlation between water contact angle and KPR adherence on SiO₂ surfaces, J. Electrochem. Soc., 112, 457–458, 1965.

5 Bowdle, D. A., Hobbs, P. V., and Radke, L. F.: Particles in the lower troposphere over the High Plains of the United States. Part III: Ice nuclei, J. Appl. Meteorol., 24, 1370–1376, 1985.

Bragg, L. H.: Pollen size variation in selected grass taxa, Ecology, 50, 124–127, 1969.

Brandl, M. T., Quiñones, B., and Lindow, S. E.: Heterogeneous transcription of an indoleacetic acid biosynthetic gene in *Erwinia herbicola* on plant surfaces, Proc. Natl. Acad. Sci., 98, 3454–3459, doi: 10.1073/pnas.061014498, 2001.

10 Bryant, G. W., Hallett, J., and Mason, B. J.: The epitaxial growth of ice on single-crystalline substrates, J. Phys. Chem. Solids, 12, 189–195, 1959.

Carte, A. E. and Mossop, S. C.: Measurements of the concentration of atmospheric ice nuclei in Southern Africa, Bulletin de l'Observatoire du Puy de Dome, 4, 137–149, 1960.

15 Chen, C.-C., Hung, L.-C., and Hsu, H.-K.: Heterogeneous nucleation of water vapor on particles of SiO₂, Al₂O₃, TiO₂, and carbon black, J. Colloid Interface Sci., 157, 465–477, 1993.

Chen, J.-P.: Theory of deliquescence and modified Köhler curves, J. Atmos. Sci., 51, 3505–3516, 1994.

20 Chen, J.-P., Wang, Z., Young, C.-Y., Tsai, F., Tsai, I.-C., Wang, G.-J., Shieh, W.-C., Lin, H. W., Huang, J.-Y., and Lu, M.-J.: Simulations of Asian yellow dust incursion over Taiwan for the spring of 2002 and 2003, Terr. Atmos. Oceanic Sci., 15, 949–981, 2004.

Cooper, W.: A possible mechanism for contact nucleation, J. Atmos. Sci., 31, 1832–1837, 1974.

25 Cooper, W. A. and Saunders, C. P. R.: Winter storms over the San Juan Mountains. Part II: Microphysical processes, J. Appl. Meteorol., 19, 927–941, 1980.

Corrin, M. L. and Nelson, J. A.: Energetics of the Adsorption of Water Vapor on “Pure” Silver Iodide, J. Phys. Chem., 72, 643–645, 1968.

30 Cotton, W. R., Stephens, M. A., Nehr Korn, T., and Tripoli, G. J.: The Colorado State University three-dimensional cloud/mesoscale model–1982. Part II: An ice phase parameterization, J. Rech. Atmos., 16, 295–320, 1982.

DeMott, P. J.: An Exploratory Study of Ice Nucleation by Soot Aerosols, J. Appl. Meteorol., 29, 1072–1079, 1990.

DeMott, P. J. and Rogers, D. C.: Freezing nucleation rates of dilute solution droplets measured

**Ice nucleation rate
from contact angle
and activation energy**

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- between -30 and -40°C in laboratory simulations of natural clouds, *J. Atmos. Sci.*, 47, 1056–1064, 1990.
- Diehl, K., Quick, C., Matthias-Maser, S., Mitra, S. K., and Jaenicke, R.: The ice nucleating ability of pollen, Part I: Laboratory studies in deposition and condensation freezing modes, *Atmos. Res.*, 58, 75–87, 2001.
- Diehl, K., Matthias-Maser, S., Jaenicke, R., and Mitra, S. K.: The ice nucleating ability of pollen: Part II. Laboratory studies in immersion and contact freezing modes, *Atmos. Res.*, 61, 125–133, 2002.
- Dingle, A. N.: Pollen as condensation nuclei, *J. Rech. Atmos.*, 2, 232–237, 1966.
- Dufour, L. and Defay, R.: *Thermodynamics of Clouds*. International Geophysics Series, vol. 6, Academic Press, New York and London, 255 pp., 1963.
- Durham, O. C.: *Methods in aerobiology*, *J. Aviation Med.*, 12, 153–162, 1941.
- Dymarska, M., Murray, B. J., Sun, L., Eastwood, M. L., Knopf, D. A., and Bertram, A. K.: Deposition ice nucleation on soot at temperatures relevant for the lower troposphere, *J. Geophys. Res.*, 111, D04204, doi:10.1029/2005JD006627, 2006.
- Eadie, W. J.: A molecular Theory of the Homogeneous Nucleation of Ice from Supercooled Water. Ph.D. thesis, Department of Geophysical Sciences, University of Chicago, 1971.
- Evans, L. F. and Lane, J. E.: Line tension and ice nucleation theory, *J. Atmos. Sci.*, 30, 326–331, 1973.
- Field, P. R., Möhler, O., Connolly, P., Krämer, M., Cotton, R., Heymsfield, A. J., Saathoff, H., and Schnaiter, M.: Some ice nucleation characteristics of Asian and Saharan desert dust, *Atmos. Chem. Phys.*, 6, 2991–3006, 2006, <http://www.atmos-chem-phys.net/6/2991/2006/>.
- Fletcher, N. H.: Size Effect in Heterogeneous Nucleation, *J. Chem. Phys.*, 29, 572–576, 1958.
- Fletcher, N. H.: *The Physics of Rainclouds*, Cambridge University Press, 286 pp., 1962.
- Gence, N.: Wetting behavior of magnesite and dolomite surfaces, *Appl. Surface Sci.*, 252, 3744–3750, 2006.
- Georgii, H. W. and Kleinjung, E.: Relations between the chemical composition of atmospheric aerosol particles and the concentration of natural ice nuclei, *J. Rech. Atmos.*, 3, 145–156, 1967.
- Gorbunov, B., Baklanov, A., Kakutkina, N., Windsor, H., and Toumi, R.: Ice nucleation on soot particles, *J. Aerosol Sci.*, 32, 199–215, 2001.
- Gultepe, I., Isaac, G. A., and Cober, S. G.: Ice crystal concentration versus temperature, *Int. J. Climatol.*, 21, 1281–1302, 2001.

**Ice nucleation rate
from contact angle
and activation energy**

J.-P. Chen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Ice nucleation rate
from contact angle
and activation energy**

J.-P. Chen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Gustafsson, R. J., Orlov, A., Badger, C. L., Griffiths, P. T., Cox, R. A., and Lambert, R. M.: A comprehensive evaluation of water uptake on atmospherically relevant mineral surfaces: DRIFT spectroscopy, thermogravimetric analysis and aerosol growth measurements, *Atmos. Chem. Phys.*, 5, 3415–3421, 2005, <http://www.atmos-chem-phys.net/5/3415/2005/>.
- 5 Hagen, D. E., Anderson, R. J. and Kassner, J. L., Jr.: Homogeneous condensation-freezing nucleation rate measurements for small water droplets in an expansion cloud chamber, *J. Atmos. Sci.*, 38, 1236–1243, 1981.
- Hallett, J. B., Gardiner, B., Hudson, J., and Rogers, F.: Cloud condensation and ice nucleation of a range of carbonaceous aerosols. Preprints, Amer. Meteor. Soc. Conf. on Cloud Physics, Vol. 2, Snowmass, Amer. Meteorol. Soc., 9–12, 1986.
- 10 Harrington, J. B. and Metzger, K.: Ragweed pollen density, *Am. J. Bot.*, 50, 532–539, 1963.
- Hazra, A., Saha, M., De, U. K., Mukherjee, J., and Goswami, K.: Study of ice nucleating characteristics of *Pseudomonas aeruginosa*, *J. Aero. Sci.*, 35, 1405–1414, 2004.
- Heffernan, K. J. and Bracewell, R. N.: Comparison of Florida and California freezing-nucleus measurements, January 1957, *J. Meteorol.*, 16, 337–339, 1959.
- 15 Heymsfield, A. J. and Sabin, R. M.: Cirrus crystal nucleation by homogeneous freezing of solution droplets, *J. Atmos. Sci.*, 46, 2252–2264, 1989.
- Hobbs, P. V. and Locatelli, J. D.: Ice Nucleation from a natural forest fire, *J. Appl. Meteorol.*, 8, 833–834, 1969.
- 20 Hu, X. L. and Michaelides, A.: Ice formation on kaolinite: Lattice match or amphoterism?, *Surf. Sci.*, 601, 5378–5381, 2007.
- Huffman, P. J.: Supersaturation spectra of Agl and natural ice nuclei, *J. Appl. Meteorol.*, 12, 1080–1082, 1973.
- Hung, H. M., Malinowski, A., and Martin, S. T.: Kinetics of heterogeneous ice nucleation on the surfaces of mineral dust cores inserted into aqueous ammonium sulfate particles, *J. Phys. Chem. A*, 107, 1296–1306, 2003.
- 25 Hussain, K. and Saunders, C. P. R.: Ice nucleus measurement with a continuous flow chamber, *Q. J. Roy. Meteorol. Soc.*, 110, 75–84, 1984.
- Isono, K. and Komabayasi, M.: Volcanoes as a source of atmospheric ice nuclei, *J. Meteorol. Soc. Jpn.*, 32, 345–353, 1954.
- 30 Isono, K., Komabayasi, M., and Ono, A.: The nature and origin of ice nuclei in the atmosphere, *J. Meteorol. Soc. Jpn.*, 37, 211–233, 1959.
- Janczuk, B. and Zdziennicka, A.: A study on the components of surface free energy of quartz

- from contact angle measurements, *J. Materials Sci.*, 29, 3559–3564, 1994.
- Kishcha, P., Barnaba, F., Gobbi, G. P., Alpert, P., Shtivelman, A., Krichak, S. O., and Joseph, J. H.: Vertical distribution of Saharan dust over Rome (Italy): Comparison between 3-year model predictions and lidar soundings, *J. Geophys. Res.*, 110, D06208, doi:10.1029/2004JD005480, 2005.
- Korolev, A. V., Isaac, G. A., Cober, S. G., Strapp, J. W., and Hallett, J.: Microphysical characterization of mixed-phase clouds, *Q. J. Roy. Meteorol. Soc.*, 129, 39–65, 2003.
- Kumai, M.: Electron microscope study of snow crystal nuclei, *J. Meteorol.*, 18, 151–156, 1951.
- Langer, G.: Evaluation of NCAR ice nucleus counter. Part I: Basic operation, *J. Appl. Meteorol.*, 12, 1000–1011, 1973.
- Lau, K.-M. and Wu, H.-T.: Warm rain processes over tropical oceans and climate implications, *Geophys. Res. Lett.*, 30, 2290–2294, 2003.
- Levin, Z., Yankofsky, S. A., Pardes, D., and Magal, N.: Possible application of bacterial condensation freezing to artificial rainfall enhancement, *J. Climate Appl. Meteorol.*, 26, 1188–1197, 1987.
- Lighthart, B.: The ecology of bacteria in the al fresco atmosphere, *FEMS Microbiol. Ecol.*, 23, 263–274, 1997.
- Lindow, S. E.: The importance of bacterial ice nuclei in plant frost injury, *Curr. Top. Plant Biochem. Physiol.*, 2, 119–128, 1983.
- Lin, Y.-L., Farley, R. D., and Orville, H. D.: Bulk parameterization of the snow field in a cloud model, *J. Climate Appl. Meteor.*, 22, 1065–1089, 1983.
- Liu, X. Y.: A new kinetic model for three-dimensional heterogeneous nucleation, *J. Chem. Phys.*, 111, 1628–1635, 1999.
- Lu, H. M. and Jiang, Q.: Size-dependent surface energies of nanocrystals, *J. Phys. Chem.*, 108, 5617–5619, 2004.
- Maki, L. R., Galyan, E. L., Chang-Chien, M. M., and Caldwell, D. R.: Ice Nucleation induced by *Pseudomonas syringae*, *Appl. Environ. Microbiol.*, 28, 456–459, 1974.
- Mansvelt, E. L. and Hattingh, M. J.: Scanning electron microscopy of invasion of apple leaves and blossoms by *Pseudomonas syringae* pv. *Syringae*, *Appl. Environ. Microbiol.*, 55, 533–538, 1989.
- Mason, B. J.: *The Physics of Clouds*, Oxford, Clarendon Press, 544 pp., 1971.
- Meyers, M. P., DeMott, P. J., and Cotton, W. R.: New primary ice-nucleation parameterizations in an explicit cloud model, *J. Appl. Meteorol.*, 31, 708–721, 1992.

**Ice nucleation rate
from contact angle
and activation energy**J.-P. Chen et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Ice nucleation rate
from contact angle
and activation energy**

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Miller, H. J., Quinn, C. E., and Graham, D. C.: A strain of *Erwinia herbicola* pathogenic on *Gypsophila paniculata*, *Neth. J. Pl. Path.*, 87, 167–172, 1981.
- Möhler, O., Field, P. R., Connolly, P., Benz, S., Saathoff, H., Schnaiter, M., Wagner, R., Cotton, R., Krämer, M., Mangold, A., and Heymsfield, A. J.: Efficiency of the deposition mode ice nucleation on mineral dust particles, *Atmos. Chem. Phys.*, 6, 3007–3021, 2006, <http://www.atmos-chem-phys.net/6/3007/2006/>.
- Pasken, R. and Pietrowicz, J. A.: Using dispersion and mesoscale meteorological models to forecast pollen concentrations, *Atmos. Environ.*, 39, 7689–7701, 2005.
- Reisner, J., Rasmussen, R. M., and Bruintjes, R. T.: Explicit forecasting of supercooled liquid water in winter storms using the MM5 mesoscale model, *Q. J. Roy. Meteor. Soc.*, 124, 1071–1107, 1998.
- Plotkowski, M. C., Chevillard, M., Pierrot, D., Altenmayer, D., Zahm, J. M., Colliot, G., and Puchelle, E.: Differential adhesion of *Pseudomonas aeruginosa* to human respiratory epithelial cells in primary culture, *J. Clin. Invest*, 87, 2018–2028, 1991.
- Pruppacher, H. R. and Klett, J. D.: *Microphysics of clouds and precipitation*, Kluwer, Netherlands, 160 pp., 1997.
- Ruggles, J. A., Nemecek-Marshall, M., and Fall, R.: Kinetics of appearance and disappearance of classes of bacterial ice nuclei support an aggregation model for ice nucleus assembly, *J. Bacteriology*, 175, 7216–7221, 1993.
- Schnell, R. C.: Biogenic sources of atmospheric ice nuclei, Univ. of Wyoming., Report No. AR111., 45 pp., 1974.
- Schnell, R. C. and Vali, G.: World wide source of leaf derived freezing nuclei, *Nature*, 246, 212–213, 1973.
- Seisel, S., Pashkova, A. Lian, Y., and Zellner, R.: Water uptake on mineral dust and soot: A fundamental view of the hydrophilicity of atmospheric particles?, *Faraday Discuss.*, 130, 437–451, doi:10.1039/b417449f, 2005.
- Stein, D. and Georgii, H. W.: Investigation on the saturation spectra of ice nuclei, *Idojaras, J. Hungarian Meteorological Service*, 86, 124, 1982.
- Ström, J. and Ohlsson, S.: In situ measurements of enhanced crystal number densities in cirrus clouds caused by aircraft exhaust, *J. Geophys. Res.*, 103, 11 355–11 361, 1998.
- Tolman, R. C.: The effect of droplet size on surface tension, *J. Chem. Phys.*, 17, 333–337, 1949.
- Turnbull, D. and Fisher, J. C.: Rate of nucleation in condensed systems, *J. Chem. Phys.*, 17,

71–73, 1949.

Turnbull, D. and Vonnegut, B.: Nucleation catalysis, *Ind. Eng. Chem.*, 44, 1292–1298, 1952.

Uno, I., Carmichael, G. R., Streets, D., Satake, S., Takemura, T., Woo, J.-H., Uematsu, M., and Ohta, S.: Analysis of surface black carbon distributions during ACE-Asia using a regional-scale aerosol model, *J. Geophys. Res.*, 108, D23, 8636, doi:10.1029/2002JD003252, 2003.

Vali, G., Cristensen, M., Fresh, R. W., Galyan, E. L., Maki, L. R., and Schnell, R. C.: Biogenic ice nuclei. II. Bacterial sources, *J. Atmos. Sci.*, 33, 1565–1570, 1976.

Vehkamäki, H., Määttänen, A., Lauri, A., Napari, I., and Kulmala, M.: Technical Note: The heterogeneous Zeldovich factor, *Atmos. Chem. Phys.*, 7, 309–313, 2007, <http://www.atmos-chem-phys.net/7/309/2007/>.

von Blohn, N., Mitra, S. K., Diehl, K., and Borrmann, S.: The ice nucleating ability of pollen, Part III: New laboratory studies in immersion and contact freezing modes including more pollen types, *Atmos. Res.*, 78, 182–189, 2005.

Wisner, C., Orville, H. D., and Meyers, C.: A numerical model of a hail-bearing cloud, *J. Atmos. Sci.*, 29, 1160–1181, 1972.

Yankofsky, S. A., Levin, Z., Bertold, T., and Sandlerman, N.: Some basic characteristics of bacterial freezing nuclei, *J. Appl. Meteorol.*, 20, 1013–1019, 1981.

ACPD

8, 14419–14465, 2008

Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Information regarding to the species of ice nuclei, including the mode of nucleation, the assigned sizes, the size range, as well as the data sources of nucleation rate and particle size.

category	sub-species	nucleation mode	diameter in μm		reference	
			assigned	range	nucleation data	size
dust	Asian dust	deposition	0.4	0.1~2	a	a,b
	Saharan dust	deposition	0.35	0.1~2	a	a,b
	Arizona test dust	deposition	0.35	0.1~2	b	a,b
bacteria	Hematite/corundum	freezing	–	0.05~0.25	c	c
	<i>E. herbicola</i>	freezing	1.	0.5~ 2.0	d	d, i*, j*, m
	<i>P. syringae</i>	freezing	1.	0.5~2	d	d, k*, m
	<i>P. aeruginosa</i>	freezing	1.	0.5~3.0	e	e, l*, m
	birch	freezing	25.	–	f	f
pollen	oak	freezing	25.	20~35	f	f
	grass	freezing	35.	30~45	f	f,n
	pine	freezing	50.	35~70	f	f
	eucalyptus sp.	freezing	45.	10~80	f	g*
soot	China rose	freezing	45.	10~80	g	g*
	–	freezing	0.1	0.08~0.12	h	h

* electron microscopy images

- a. Field et al. (2006); b. Möhler et al. (2006); c. Hung et al. (2003);
 d. Yankofsky et al. (1981); e. Hazra et al. (2004); f. Diehl et al. (2001);
 g. this study; h. DeMott (1990); i. Brandl et al. (2001);
 j. Miller et al. (1981); k. Mansvelt and Hattingh (1989); l. Plotkowski et al. (1991);
 m. Levin et al., 1987; n. Bragg (1969).

Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Table 2. Results of IN properties derived from the initial analysis. Units are 10^{-20} J for $\Delta g^\#$, and μm for r_N . The coefficient of determination, R^2 , for the final fitting is also listed.

Species	Assigned r_N	Eq. 17a 'f'	Eq. 17b		Eq. 17c		R^2
			$\Delta g^\#$	r_N	f	r_N	
Soot	0.05	0.0693	12.75	0.06	0.0686	0.06	0.994
<i>E. herbicola</i>	0.5	0.0025	13.16	0.82	0.00249	0.82	0.990
<i>P. syringae</i>	0.5	0.000719	13.42	0.90	0.000706	0.88	0.996
<i>P. aeruginosa</i>	0.5	0.00024	13.32	1.11	0.000225	1.11	0.940
Grass	17.5	0.0065	15.71	17.66	0.0063	17.34	0.951
Oak	12.5	0.0075	15.21	12.5	0.0073	12.21	0.989
Pine	25	0.0029	15.71	25.07	0.0028	24.39	0.985
Birch	12.5	0.0052	15.10	12.21	0.0050	11.79	0.934
Eucalyptus	22.5	0.0058	15.46	23.1	0.0057	22.9	0.990
China rose	22.5	0.0072	15.41	20.1	0.0071	19.87	0.992
Asian dust	0.2	0.00142	0.6317	0.29	0.00137	0.27	0.958
Saharan dust	0.175	0.00085	1.685	0.68	0.00084	0.67	0.972
ATD	0.175	0.000089	2.763	0.41	0.000087	0.4	0.995

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Table 3. Coefficients of the fitting formulas in Eq. (18) for various ice nuclei and under different ranges of germ size ratio q , contact angle θ and the corresponding wetting coefficient. Also given is the maximum error from each approximation formula.

	bacteria	soot	pollens	Asian dust	Saharan dust	ATD	Hung et al.
q	10~363	9~140	>250	10~200	5~110	2.4~44	40~400
m	0.940~0.997	0.602~0.875	0.866~0.971	0.940~0.970	0.961~0.990	0.985~0.996	-0.866~0.450
θ	4~20	29~53	14~30	14~20	8~16	5~10	63~150
c_1	-0.3541568	-0.5411955	-0.3962579	-0.3598818	-0.3353619	-0.3497709	-0.7352132
c_2	1.983928	1.879918	1.968181	1.982032	1.990979	1.994605	1.315406
c_3	2.019456	1.607947	1.858932	2.025390	2.175539	2.527253	0.5413986
r^2	0.9999	0.9999	0.9999	0.9999	0.9999	0.9994	0.9853
max. error	3.1%	3.9%	1.46%	0.24%	0.8%	3.1%	12%
c_4	-0.08853648	-0.27167433	-0.3748873	-0.06841440	0.1725698	0.7220012	-0.7064756
c_5	1.983878	1.879725	1.968181	1.983733	1.990979	1.994605	1.3154064
c_6	-0.05019220	-0.05747428	-0.002749135	-0.05734672	-0.1147096	-0.3046048	-0.00478509
r^2	0.9990	0.9987	0.9999	0.9976	0.9963	0.9794	0.9853
max. error	6.2%	5.4%	1.43%	2.6%	6.2%	15%	11%

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Table 4. Results of IN properties derived from the refined analysis. Units are 10^{-20} J for $\Delta g^\#$, degree for θ , and μm for r_N . Note that f^* is calculated from m using Eq. (9) for comparison with the rough estimates given in Table 2. The coefficient of determination, R^2 , for each fitting is also listed.

Species	$\Delta g^\#$	m	f^*	θ	Assigned r_N	Derived r_N	R^2
Soot	12.6	0.691	6.42E-2	46.3	0.05	0.05	0.993
<i>E. herbicola</i>	12.8	0.943	2.39E-3	19.4	0.5	0.5	0.990
<i>P. syringae</i>	13.0	0.969	7.04E-4	14.3	0.5	0.5	0.996
<i>P. aeruginosa</i>	12.7	0.983	2.10E-4	10.5	0.5	0.5	0.941
Grass	15.7	0.907	6.27E-3	24.9	17.5	17.6	0.951
Oak	15.2	0.899	7.41E-3	26.0	12.5	12.5	0.989
Pine	15.7	0.939	2.75E-3	20.2	25.0	24.8	0.985
Birch	15.1	0.917	5.05E-3	23.5	12.5	12.4	0.934
Eucalyptus	15.9	0.912	5.66E-3	24.24	22.5	22.57	0.990
China rose	16.0	0.901	7.08E-3	25.7	22.5	22.49	0.992
Asian Dust	1.38	0.959	1.26E-3	14.2	0.20	0.200	0.956
Saharan Dust	1.00	0.969	7.18E-4	14.3	0.175	0.171	0.971
ATD	2.36	0.991	6.19E-5	7.8	0.175	0.173	0.991

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Table 5. Contact angle and activation energy calculated from the freezing nucleation data of Hung et al. (2003) with consideration of the solute effect and curvature effect on the surface tension. The conditions for surface tension include solute mole fraction χ and ice-germ size r , so $\sigma(0,\infty)$ means both effects are ignored, $\sigma(\chi,\infty)$ includes solute effect only, and $\sigma(\chi,r)$ considers both effects. Values were calculated from data for hematite at concentrations of $\chi=0.03$ and $\chi=0.13$. Units are 10^{-20} J for Δg_a and degree for θ .

Solute mole fraction χ		0.03			0.13		
parameters	conditions	m	θ	Δg_a	m	θ	Δg_a
$\sigma(0,\infty)$		-0.279	109.2	13.3	0.269	74.4	14.1
$\sigma(\chi,\infty)$		-0.825	145.6	14.1	-1.609	–	15.7
$\sigma(\chi,r)$		0.351	69.4	9.9	0.448	63.4	14.3

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Table 6. Contact angle and activation energy for various IN species calculated with and without the curvature adjustment on surface tension. The values are for freezing nucleation except for the lowest three rows (AD, SD, and ATD), which pertain to deposition nucleation. Units are 10^{-20} J for $\Delta g^\#$ and degree for θ .

Curvature adjustment	no		yes	
	θ	$\Delta g^\#$	$\theta(r)$	$\Delta g^\#(r)$
parameter species				
soot	46.3	12.7	33.2	13.8
<i>E. herbicola</i>	19.4	12.8	16.0	12.7
<i>P. syringae</i>	14.3	13.0	12.5	12.8
<i>P. aeruginosa</i>	10.5	12.7	5.30	12.2
Grass	24.9	15.7	18.4	15.5
Oak	26.0	15.3	20.6	15.1
Pine	20.2	15.7	17.8	15.2
Birch	23.5	15.1	18.8	15.0
Eucalyptus	24.2	15.9	18.4	15.7
China rose	25.7	16.0	20.3	15.8
Hematite (0.03)	145.6	14.1	69.4	9.90
Hematite (0.13)	–	15.7	63.4	14.3
AD	14.2	1.38	15.2	1.17
SD	14.3	1.00	14.7	1.31
ATD	7.78	2.36	7.85	2.32

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Ice nucleation rate from contact angle and activation energy

J.-P. Chen et al.

Table 7. Examination of the effect of assigned IN size on the results of activation energy and contact angle parameters. The values shown are percentage changes for each parameter when the originally assigned r_N is altered by one third or three times.

	change in $\Delta g^\#$		change in θ	
	$1/3r_N$	$3 r_N$	$1/3r_N$	$3 r_N$
Soot	-4.8%	5.6%	-3.9%	2.1%
<i>E. herbicola</i>	-6.2%	6.3%	-1.9%	0.7%
<i>P. syringae</i>	-6.2%	6.2%	-2.7%	0.0%
<i>P. aeruginosa</i>	-5.5%	6.3%	0.0%	0.0%
Grass pollen	-5.1%	5.1%	-0.9%	0.8%
Oak pollen	-4.6%	5.3%	-0.7%	0.2%
Pine pollen	-4.5%	5.1%	-0.6%	1.2%
Birch pollen	-4.6%	5.3%	-0.3%	0.7%
Eucalyptus pollen	-4.4%	5.0%	-0.6%	0.6%
China rose pollen	-5.0%	5.0%	-0.5%	0.5%
Asia dust	-79.6%	131.9%	-4.5%	2.8%
Saharan dust	-68.4%	76.3%	-5.5%	2.5%
Arizona test dust	-30.1%	31.0%	-18.0%	3.7%

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Ice nucleation rate
from contact angle
and activation energy

J.-P. Chen et al.




Fig. 1. Effects of various degrees of linearization. Panels (a) and (b) applies linearization from Eqs. (17a) and (19a), respectively, both without curvature adjustment; panel (c) is the same as panel (b) but with curvature adjustment. Not all species studied here are included in order to avoid cluttering of curves.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Ice nucleation rate
from contact angle
and activation energy

J.-P. Chen et al.




Fig. 2. Temperature dependence of the activation energy for various IN species, calculated by assuming constant m according to the values given in Table 6: **(a)** Δg_a for the freezing nucleation on other IN species; **(b)** Δg_d for the deposition nucleation on dust. Note that the vertical scale of panel (a) does not originate from zero so the range of values is relatively small.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)