

**N₂O enhancement
during solar proton
events**

B. Funke et al.

Enhancement of N₂O during the October–November 2003 solar proton events

**B. Funke¹, M. García-Comas¹, M. López-Puertas¹, N. Glatthor², G. P. Stiller²,
T. von Clarmann², K. Semeniuk³, and J. C. McConnell³**

¹Instituto de Astrofísica de Andalucía, CSIC, Granada, Spain

²Forschungszentrum Karlsruhe und Universität Karlsruhe, Institut für Meteorologie und
Klimaforschung, Karlsruhe, Germany

³Department of Earth and Space Science and Engineering, York University, Toronto, Ontario,
Canada

Received: 4 January 2008 – Accepted: 31 January 2008 – Published: 4 March 2008

Correspondence to: M. López-Puertas (puertas@iaa.es)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

In this paper we present evidence of enhanced N_2O concentrations in the upper stratosphere/lower mesosphere polar regions after the solar proton events that occurred during October–November 2003. The observations were performed by the MIPAS instrument on the Envisat satellite. Simulations performed using the Canadian Middle Atmospheric Model (CMAM) show that such enhancements are most likely produced by the reaction of $\text{N}(^4\text{S})$ with NO_2 , both of which species are largely enhanced just after the solar proton events in the winter polar night.

1 Introduction

Nitrous oxide is the main precursor of active nitrogen in the middle atmosphere. Its major sources, both natural and man-made, originate at the surface, and it is transported into the stratosphere, where photo-dissociation by solar UV is its major sink and where its reaction with $\text{O}(^1\text{D})$ leads to the formation of chemically active nitrogen species like NO_2 and NO . Hence it indirectly plays a major role in controlling atmospheric ozone abundance through the NO_x ($\text{NO}_x = \text{NO} + \text{NO}_2$) ozone-destroying catalytic cycle (Brasseur and Solomon, 2005).

It has been widely reported that solar proton events (SPEs) have significant effects on the composition of the stratosphere and mesosphere in the polar regions (e.g., Jackman and McPeters, 2004, for a recent review). The major effects have been found to be significant enhancements in HO_x ($\text{H} + \text{OH} + \text{HO}_2$) and NO_x ($\text{NO} + \text{NO}_2$), followed by large depletions of O_3 in these atmospheric regions.

In recent years, there have been two large solar proton events (October–November 2003 and January 2005) (Jackman et al., 2008) which have been intensively observed by several instruments on different satellite platforms, including, for example, NOAA 16 SBUV/2 and HALOE data (Jackman et al., 2005a,b; Randall et al., 2005); MIPAS, GOMOS and SCIAMACHY on Envisat (Seppälä et al., 2004; López-Puertas et al.,

N_2O enhancement during solar proton events

B. Funke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2005a,b; von Clarmann et al., 2005; Orsolini et al., 2005; Rohen et al., 2005); and MLS on AURA (Verronen et al., 2006). In particular, during late October and early November 2003, three active solar regions produced solar flares and solar energetic particles of extremely large intensity, the fourth largest event observed in the past forty years (Jackman et al., 2005a, 2008). Some of the Geostationary Operational Environmental Satellite (GOES)-11 instruments measured very large fluxes of highly energetic protons (e.g., López-Puertas et al., 2005a). The protons are guided by the Earth's magnetic field to both polar regions (geomagnetic latitudes $>60^\circ$), where they penetrate down to ~ 87 km, if their energy is >1 MeV, or even down to ~ 30 km, if their energy is >100 MeV (Jackman et al., 2005a).

During and after this solar proton event, the MIPAS instrument observed global changes (e.g. in both the Northern and Southern polar regions, during day and nighttime) in the stratospheric and lower mesospheric composition. This includes enormous enhancements in NO_x , e.g., in NO and NO_2 , and large depletions in O_3 (López-Puertas et al., 2005a) as well as significant changes in other NO_y species, such as HNO_3 , N_2O_5 , ClONO_2 (López-Puertas et al., 2005b). In addition, there also have been observed changes in ClO and HOCl, as evidence of perturbations by solar protons on the HO_x and chlorine species abundances (von Clarmann et al., 2005).

However, to our knowledge, no solar-proton-event induced changes in N_2O concentration have so far been reported. A paper on the production and transport of N_2O by auroral electron precipitation seen by the Fourier transform spectrometer on SCISAT-1 has just been submitted by Semeniuk et al. (2008).¹ One should expect this species being enhanced also if sufficient atomic nitrogen and NO_2 is available after the solar proton event in the polar night upper stratosphere and lower mesosphere. The aim of this paper is to report the enhanced N_2O concentration observed by MIPAS during the October–November 2003 solar proton event and to explain the reason for it using

¹Semeniuk, K., McConnell, J. C., Jin, J. J., Jarosz, J. R., Boone, C. D., and Bernath, P. F.: N_2O production by high energy auroral electron precipitation, J. Geophys. Res., submitted, 2008.

N_2O enhancement during solar proton events

B. Funke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



modeling calculations by the Canadian Middle Atmospheric Model (CMAM).

2 MIPAS measurements

MIPAS is a limb emission Fourier transform spectrometer designed for the measurement of trace species from space (Fischer and Oelhaf, 1996; ESA, 2000; Fischer et al., 2007). It is part of the instrumentation of the Environmental Satellite (ENVISAT) which was launched into its sun-synchronous polar orbit of 98.55° inclination at about 800 km altitude on 1 March 2002. MIPAS operated from July 2002 to March 2004 at full spectral resolution of 0.035 cm^{-1} (unapodized) in terms of full width at half maximum and with reduced resolution since August 2004. MIPAS observes the atmosphere during day and night with global coverage from pole to pole. Within its standard observation mode at full spectral resolution, MIPAS covers the altitude range from 68 down to 6 km with tangent altitudes at 68, 60, 52, 47, and then at 3 km steps from 42 to 6 km. Occasionally, MIPAS also operates in several upper atmospheric modes scanning up to 170 km. The field of view of MIPAS is 30 km in horizontal and approximately 3 km in vertical direction. MIPAS passes the equator in southerly direction at 10.00 am local time 14.3 times a day. During each orbit up to 72 limb scans are recorded. The Level-1b processing of the data (version 4.61/62 was used here), including processing from raw data to calibrated phase-corrected and geolocated radiance spectra, is performed by the European Space Agency (ESA) (Nett et al., 1999, 2002).

The retrieval of N_2O abundances was performed with the IMK-IAA data processor (von Clarmann et al., 2003a), which is based on a constrained non-linear least squares algorithm with Levenberg-Marquardt damping and line by line radiative transfer calculations with the Karlsruhe Optimized and Precise Radiative Transfer Algorithm (KOPRA) (Stiller et al., 2000). The first step in the L2 processing was the determination of the spectral shift, followed by the retrieval of temperature and elevation pointing (von Clarmann et al., 2003b), where pressure is implicitly determined by means of hydrostatic equilibrium. The retrieval of volume mixing ratio (vmr) profiles of species was carried

N_2O enhancement during solar proton events

B. Funke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



out in the following order: O_3 , H_2O , HNO_3 , and then CH_4 and N_2O simultaneously. The results of the species retrieved first are then used in the retrievals of the subsequent species.

The N_2O vmr was retrieved from the MIPAS spectra around 1284.9 cm^{-1} , where the ν_1 band of N_2O is located (Glatthor et al., 2005). The retrievals were performed from selected spectral regions (micro-windows) which vary with observation geometries in order to optimize computation time and minimize systematic errors (von Clarmann and Echle, 1998). Thus, height dependent combinations of micro-windows were selected with a trade-off between computation time and total retrieval error. The retrieval noise error in the N_2O vmr for unperturbed conditions (i.e., not during solar proton events) is typically 3% at 10–44 km and 22% at 50 km. The total error varies between 10 and 20% at 10–35 km and is about 30% between 35–50 km (Glatthor et al., 2005).

The resulting vertical resolution was typically about 4 km in the altitude range 15–40 km and decreased to more than 10 km below and above this region. More details on the N_2O retrieval strategy can be found in Glatthor et al. (2005). For a profile of the enhanced N_2O during the SPE (typical of the data shown here) the noise error is about 0.2 ppbv (smaller than 5%) at 50–75 km. However, the vertical resolution is rather sparse with values of 8–15 km above 52 km. This study is focussed mainly on MIPAS data from 25 October to 22 November 2003, which were retrieved by the IMK-IAA processor (data version V3O_N2O_12) from Level-1b data version 4.61/62.

In addition to N_2O we also use MIPAS data for NO_2 for that period and for 1 November for NO . These data correspond to the IMK/IAA processed versions V3O_NO2_9 and V3O_NO2_11 and V3O_NO_9, which have been significantly improved from previously published data in López-Puertas et al. (2005a).

2.1 N_2O Enhancement in polar regions

Figure 1 shows the temporal evolution of the N_2O distribution at an approximate altitude of 58 km in the Northern polar Hemisphere (40°N – 90°N) for the period of 26 October to 11 November 2003, i.e., from two days before the major solar proton events (SPEs)

N_2O enhancement during solar proton events

B. Funke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



until about ten days after them (note that some days are not shown). Since we are comparing these data with the corresponding NO₂ maps (see below), and at this height nighttime–NO₂ is a good proxy of total NO_x, we have plotted the N₂O maps considering only nighttime conditions.

5 A large increase in the N₂O abundance is observed at polar latitudes reaching values of about 9 ppbv at 58 km after the SPEs. This coincides with the latitudes where solar protons penetrate into the atmosphere, that is, in the polar cap regions (approximately at >60° geomagnetic latitude). Note the contrast between the distribution on 26 October, before the SPEs, and on 29 October and following days, during and after the
10 SPEs. Maximum N₂O abundances are observed during the first few days of the SPEs (on 30 October until 2 November), when the larger proton fluxes took place (see Fig. 1 in López-Puertas et al., 2005a). We also observe that the enhancement in N₂O concentration lasts until 22 November (see Fig. 5a) when the enhancement is still above 2 ppbv and moved to lower altitudes.


15 The energetic charged particles (protons and associated secondary electrons) collide with and dissociate N₂, thus producing atomic nitrogen and, subsequently, odd nitrogen in the forms of NO and NO₂ through reactions:



and the further oxidation of nitric oxide through



The generation of N(⁴S) can also lead to a reduction in odd nitrogen (NO_y) via



This NO_y loss mechanism thus limits the buildup of NO_x and is important especially during large SPEs, when a huge amount of NO_y is produced in a short time (Rusch et al., 1981). However, as a net effect, SPEs will result in an increase in NO_y constituents.
25

N₂O enhancement during solar proton events

B. Funke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



For reasons discussed below, we think that the N₂O enhancement shown in Fig. 1 is produced from the enhancements in N(⁴S) and in NO₂ followed by the reaction



which is effective only during nighttime conditions since NO₂ is rapidly dissociated in the sunlight, e.g.,



Thus, the condition required for producing N₂O in the upper stratosphere is to have sufficient amounts of atomic nitrogen N and NO_x under dark conditions, which is well fulfilled when solar proton events occur in the dark polar winter region.

To verify this hypothesis we have investigated the spatial correlation between N₂O and NO₂ which indeed is very pronounced for the whole period under investigation (Figs. 1 and 2) and thus supports our explanation. Figure 3 shows the mean distribution of both species at the same altitude averaged over the first eight days after the first SPE, from 29 October to 5 November 2003. In this case we have included all data, accounting both day and nighttime conditions. The good spatial correlation between the enhancements in both species is again evident, being both confined to the polar cap (>60° geomagnetic latitude). Note the larger abundances in NO₂ close to the North pole (≳80°), which reflects the dark polar night where NO₂ is not photodissociated.

The altitude/latitude distribution of the enhancement is shown in Fig. 4 for four days, one just before the SPEs and three days after these events. The sudden appearance of the N₂O enhancement after the SPE on 29 October, delimited to latitudes northward of 60° and at altitudes above around 40 km, is clearly evident in the top right panel. The other panels of the figure show that the enhancement persists for the following few days, being slightly diluted and descended downwards. There seems to be also an indication of aurorally enhanced N₂O (e.g., produced by energetic electron precipitation EEP), in the top left panel of Fig. 4 (26 October). This small enhancement, as that appearing in late November (see next section) is not the focus of this work.

N₂O enhancement during solar proton events

B. Funke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.2 Temporal evolution of the N₂O enhancement

Figure 5a shows the temporal evolution of zonal mean N₂O in the Northern polar cap (latitudes polewards of 70° geographic) for nighttime conditions. The maximum enhancement took place at 50–55 km on 30–31 October (day of year (doy) 303–304) just after the first major SPE that came about on 29 October. Another enhancement occurred on 3–4 November (doy 307–308), following another major increase in the high-energy solar proton flux (see Fig. 1 in López-Puertas et al., 2005a). A weaker enhancement is also observed around 60 km near the end of this period (18–22 November, doy 324–326). This enhancement seems not to be related to solar proton events. Although a smaller SPE took place on 22–25 November the ionization rates due to protons calculated by Jackman (2007b) for these days were about three orders of magnitude lower than for 30–31 October. Also, this N₂O enhancement already started on 18 November, a few days before this secondary SPE occurred. This weaker enhancement, however, seems to be correlated with the onset of the polar winter descent of NO_x from the mesosphere-lower thermosphere (MLT) region which took place in the second half of November and made available sufficient NO₂ during nighttime below approximately 70 km (see Fig. 6a). Atomic nitrogen, also required for N₂O formation, has a too short chemical lifetime to be transported downward from the MLT region, but it could be produced by the high-energy electron precipitation (EEP) taking place at that time. Indeed, fluxes of electrons with energies >100 keV (able to penetrate down to 70 km) measured by the MEPED instrument on NOAA 16 (<http://poes.ngdc.noaa.gov/data/avg/>) were significantly increased on 16 and 20 November.

The temporal evolution of N₂O in the Northern Hemisphere (Fig. 5a) shows also that N₂O-rich air masses were descending from around 50 km to ~43 km during November. This descent is also observed in the temporal evolution of NO₂ (Fig. 6a) and is related to the polar winter descent of the meridional circulation.

Smaller N₂O enhancements have also been observed in the Southern Hemisphere (not shown here) but with significantly lower mixing ratios (just ~1–1.2 ppbv). These

N₂O enhancement during solar proton events

B. Funke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion



smaller enhancements in the sunlit Southern Hemisphere polar cap are expected due to less available NO₂ during daytime conditions (see Fig. 4a in López-Puertas et al., 2005a). The temporal evolution of N₂O in the Southern Hemisphere shows, in contrast to the Northern Hemisphere, a small upward motion which can also be explained by the meridional circulation pattern in this period.

3 Modelling

We have also investigated N₂O production by the 2003 SPEs using the Canadian Middle Atmosphere Model (CMAM). CMAM is a chemistry climate model with a comprehensive set of physical parameterizations and chemistry package. The model chemistry takes into account 102 gas phase, 37 photolysis and 12 heterogeneous reactions on type Ib and II polar stratospheric clouds (de Grandpré et al., 2000). For the simulations conducted here the model chemistry was extended to include Reaction (R4) and the additional branches:



Based on multiple simulations, it was decided to ascribe 50% of the reaction rate to the primary branch (R4) and 25% to each of the secondary branches (R6 and R7). This split is not excluded by the laboratory work used to determine the reaction rate (Wennberg et al., 1994) and without it the N₂O production is too high for a given amount of NO₂.

The October–November 2003 SPEs production of NO_x, HO_x and atomic oxygen based on vertical profiles of hourly ion pair production rates derived from NOAA GOES-11 observations of proton fluxes was implemented as in Semeniuk et al. (2005). However, here we use a new chemistry solver that is fully implicit without any approximation

N₂O enhancement during solar proton events

B. Funke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



to the Jacobian matrix. Ground state atomic nitrogen is not assumed to be in photochemical steady state. We carried out one simulation where Reaction (R4) and its branches were turned off and one simulation where they were active.

The run performed without including Reaction (R4) shows very little N₂O above ~40 km (due to descent during this time of the year). When this reaction is included, however, the model predicts a significant enhancement for N₂O (see Figs. 5b and 5c). The values computed by the model (Fig. 5b) look, in a first instance, significantly larger than those measured by MIPAS (Fig. 5a) and with the peak values situated at about 5–10 km higher. We mentioned before that MIPAS measurements have a rather sparse vertical resolution at these altitudes. Hence, for meaningful comparisons, the MIPAS averaging kernels have to be applied to model results. This is shown in Fig. 5c, where we show the model predictions as would have been measured by MIPAS. There is excellent agreement with the measurements (panel a), both in the magnitude of the enhancement (slightly overestimated by the model) and in the altitude of the peak of the enhancement. The model also simulates well the double peak in the temporal evolution of the enhancement, corresponding to the two major SPEs on 29 October and 3–4 November, and the dilution and slow descent of the N₂O increase during the following days. This then confirms our hypothesis about the origin of the N₂O enhancement. The model does not show the weaker enhancement on 18–20 November at about 60 km, since this is likely produced by EEPs which are not modeled in the results shown here.

As a further check of the mechanism, we compared the measured and modeled values for NO₂, one of the precursors of the N₂O, during this period. The results are shown in Figs. 6a and 6b for MIPAS and the model, respectively. In absence of reprocessed MIPAS NO data to compare the total NO_x abundance, we compared only nighttime NO₂ data, which, below about 70 km, is a good proxy for NO_x.

As has been discussed in the introduction, and shown in Fig. 6, NO₂ increased significantly during these SPEs. Focusing on the first few days after the SPEs, the figure shows that measured and modeled NO₂ are in rather good agreement. This is more clearly seen in Fig. 7 where we show the profiles for 1 November for NO₂ and NO_x. We

N₂O enhancement during solar proton events

B. Funke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



see that below ~ 50 km the agreement is very good although between 55 to 65 km modeled NO_2 and NO_x are overpredicted. This model overestimation is consistent with the slightly larger values predicted by the model for the N_2O enhancement in this region (see Fig. 5).

Large values of NO_2 seen by MIPAS above 60 km are noticeable in Fig. 6a starting on 8 November (doy 312) and mainly after 16 November (doy 320). This enhancement is attributed to a strong descent of mesospheric air, very rich in NO_x , which was produced by energetic electron precipitation in the mesosphere and with some possible local contribution by EEPs (see, e.g., López-Puertas et al., 2006). This NO_x -production mechanism is not included in the current model runs. Note that MIPAS N_2O also exhibits a weak enhancement in this region and time (see Fig. 5a), which is not shown in the model run because of insufficient NO_2 and lack of high-energy electron precipitation. The detailed study of these N_2O and NO_2 enhancements are, however, beyond the scope of this paper.

As a final remark, we should note that the small enhancement of 1–1.2 ppbv in the Southern Hemisphere is also consistent with the N_2O production mechanism proposed here. Peak averaged NO_2 values in the Northern Hemisphere are about 80 ppbv (Fig. 6a) while the enhancement in the SH (not shown here) is about 15 ppbv. The ratio of both is about a factor of 5–6, very similar to the NH/SH ratio for N_2O of (6–7)/(1–1.2). In that sense, CMAM predictions for the Southern Hemisphere are also very similar to those measured with peak averaged enhancements ranging from 1 to 1.5 ppbv.

4 Summary and conclusions

We have shown in this paper the first evidence of enhanced N_2O concentrations in the upper stratosphere/lower mesosphere polar regions after solar proton events. In particular we show the enhanced N_2O in the Northern Hemisphere polar cap after the SPEs that occurred during October–November 2003. The observations were performed with the MIPAS instrument on the Envisat satellite and show that N_2O was enhanced by

N_2O enhancement during solar proton events

B. Funke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



about 5 ppbv in the upper stratosphere/lower mesosphere Northern polar cap. Simulations performed using the Canadian Middle Atmospheric Model (CMAM) show that such enhancements can be produced by the reaction of $N(^4S)$ with NO_2 , both of which species are largely augmented just after the solar proton events in the polar night.

Such N_2O enhancement requires a condition which is rarely present in the upper stratosphere: the simultaneous presence of atomic nitrogen, N, and high amounts of NO_2 , a condition which is met when solar proton events come up under polar night conditions. High-energy electron precipitation is also capable of producing atomic nitrogen in the middle mesosphere, and might then represent a continuous source of atmospheric N_2O in the polar night regions in the upper mesosphere, which can occasionally descend to the stratopause region (Semeniuk et al., 2008¹)

N_2O is almost completely produced at the surface by both natural and anthropogenic sources. The mechanism presented here represents an additional natural atmospheric source of N_2O . Its investigation in the future is therefore of high importance.

Acknowledgements. The IAA team was supported by the Spanish project ESP2004-01556 and EC FEDER funds. The authors acknowledge ESA for providing MIPAS spectra. The IMK team was supported by the Priority Program CAWSES of the German science foundation (DFG) under the project MANOXUVA. J. C. McConnell and K. Semeniuk would like to thank the Natural Sciences and Engineering Research Council of Canada, the Canadian Foundation for Climate and Atmospheric Sciences and the Canadian Foundation for Innovation.

N_2O enhancement during solar proton events

B. Funke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



References

- Brasseur, G. and Solomon, S.: Aeronomy of the middle Atmosphere, 3rd revised and enlarged ed., Springer, Dordrecht, 644 pp., 327–358, 2005. [4670](#)
- de Grandpré, J., Beagley, S. R., Griffioen, E., McConnell, J. C., and Medvedev, A. S.: Ozone climatology using interactive chemistry: Results from the Canadian Middle Atmosphere Model, *J. Geophys. Res.*, 105, 26 475–26 491, 2000. [4677](#)
- European Space Agency, Envisat, MIPAS An instrument for atmospheric chemistry and climate research, ESA Publications Division, ESTEC, The Netherlands, SP-1229, 1–123, 2000. [4672](#)
- Fischer, H. and Oelhaf, H.: Remote sensing of vertical profiles of atmospheric trace constituents with MIPAS limb-emission spectrometers, *Appl. Optics*, 35, 16, 2787–2796, 1996. [4672](#)
- Fischer, H., Birk, M., Blom, C., Carli, B., Carlotti, M., von Clarmann, T., Delbouille, L., Dudhia, A., Ehhalt, D., Endemann, M., Flaud, J. M., Gessner, R., Kleinert, A., Koopmann, R., Langen, J., López-Puertas, M., Mosner, P., Nett, H., Oelhaf, H., Perron, G., Remedios, J., Ridolfi, M., Stiller, G., and Zander, R.: MIPAS: An Instrument for Atmospheric and Climate Research, *Atmos. Chem. Phys. Discuss.*, 7, 8795–8893, 2007, <http://www.atmos-chem-phys-discuss.net/7/8795/2007/>. [4672](#)
- Funke, B., López-Puertas, M., Gil-López, S., Clarmann, T., Stiller, G., Kellmann, S., and Fischer, H.: Downward transport of upper atmospheric NO_x into the polar stratosphere and lower mesosphere during the Antarctic 2003 and Arctic 2002/2003 winters, *J. Geophys. Res.*, 110, D24308, doi:10.1029/2005JD006463, 2005.
- Glatthor, N., von Clarmann, T., Fischer, H., Funke, B., Grabowski, U., Höpfner, M., Kellmann, S., Kiefer, M., Linden, A., Milz, M., Steck, T., Stiller, G. P., Mengistu Tsidu, G., and Wang, D.-Y.: Mixing processes during the Antarctic vortex split in September–October 2002 as inferred from source gas and ozone distributions from ENVISAT-MIPAS, *J. Atmos. Sci.*, 62, 787–800, 2005. [4673](#)
- Jackman, C. H. and McPeters, R. D.: The effects of solar proton events on ozone and other constituents, *Solar variability and its Effects on climate*, *Geophys. Mon.*, 141, 305–319, 2004. [4670](#)
- Jackman, C. H., DeLand, M. T., Labow, G. J., Fleming, E. L., Weisenstein, D. K., Ko, M., Sinnhuber, M. K. W., and Russell, J. M.: Neutral atmospheric influences of the solar proton events

ACPD

8, 4669–4691, 2008

N₂O enhancement during solar proton events

B. Funke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



in October–November 2003, *J. Geophys. Res.*, 110, A09S27, doi:10.1029/2004JA010888, 2005a. [4670](#), [4671](#)

Jackman, C. H., DeLand, M. T., Labow, G. J., Fleming, E. L., Weisenstein, D. K., Ko, M., Sinnhuber, M. K. W., Anderson, J., and Russell, J. M.: The influence of the several very large solar proton events in Years 2000–2003 on the neutral middle atmosphere, *Adv. Space Res.*, 35, 445–450, 2005b. [4670](#)

Jackman, C. H., Marsh, D. R., Vitt, F. M., Garcia, R., Fleming, E. L., Labow, G. J., Randall, C., López-Puertas, M., and Funke, B.: Short- and medium-term atmospheric effects of very large solar proton events, *Atmos. Chem. Phys.*, 8, 765–785, 2008, <http://www.atmos-chem-phys.net/8/765/2008/>. [4670](#), [4671](#)

Jackman, C. H.: Ionization rates for 1963–2007 from Solar Proton events, http://strat-www.met.fu-berlin.de/~matthes/sparc/SOLARIS_Jackman_SPEs.pdf, 2007. [4676](#)

López-Puertas, M., Funke, B., Gil-López, S., von Clarmann, T., Stiller, G. P., Höpfner, M., Kellmann, S., Fischer, H., and Jackman, C. H.: Observation of NO_x enhancement and ozone depletion in the Northern and Southern Hemispheres after the October–November 2003 solar proton events, *J. Geophys. Res.*, 110, A09S43, doi:10.1029/2005JA011050, 2005a. [4670](#), [4671](#), [4673](#), [4674](#), [4676](#), [4677](#)

López-Puertas, M., Funke, B., Gil-López, S., Mengistu Tsidu, G., Fischer, H., and Jackman, C. H.: HNO₃, N₂O₅, and ClONO₂ enhancements after the October–November 2003 solar proton events, *J. Geophys. Res.*, 110, A09S44, doi:10.1029/2005JA011051, 2005b. [4671](#)

López-Puertas, M., Funke, B., Clarmann, T., Fischer, H., and Stiller, G.: The Variability of Stratospheric and Mesospheric NO_y in the Arctic and Antarctic 2002–2004 Polar Winters, *Space Sci. Rev.*, 125, 403–416, doi:10.1007/s11214-006-9073-2, 2006. [4679](#)

Nett, H., Carli, B., Carlotti, M., et al.: MIPAS ground processor and data products, *Proc. IEEE 1999 Inter. Geosci. and Remote Sensing Symp.*, Hamburg, Germany, 1692–1696, 1999. [4672](#)

Nett, H., Perron, G., Sanchez, M., Burgess, A., and Mosner, P.: MIPAS in-flight calibration and processor verification, in: *ENVISAT Calibration Review–Proc. of the European Workshop, 9–13 September 2002*, ESTEC, Noordwijk, The Netherlands, CD-ROM, SP–520, edited by: H. Sawaya-Lacoste, ESA Publications Division, ESTEC, Postbus 299, 2200 AG Noordwijk, The Netherlands, 1–10, 2002. [4672](#)

Orsolini, Y. J., Manney, G. L., Santee, M. L., and Randall, C. E.: An upper stratospheric layer of enhanced HNO₃ following exceptional solar storms, *Geophys. Res. Lett.*, 32, L12S01,

N₂O enhancement during solar proton events

B. Funke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



doi:10.1029/2004GL021588, 2005. [4671](#)

Randall, C. E., Harvey, V. L., Manney, G. L., et al.: Stratospheric effects of energetic particle precipitation in 2003–2004, *Geophys. Res. Lett.*, 32, L05802, doi:10.1029/2004GL022003, 2005. [4670](#)

5 Rohen, G., von Savigny, C., Sinnhuber, M., Llewellyn, E. J., Kaiser, J. W., Jackman, C. H., Kallenrode, M.-B., Schroter, J., Eichmann, K.-U., Bovensmann, H., and Burrows, J. P.: Ozone depletion during the solar proton events of Oct./Nov. 2003 as seen by SCIAMACHY, *J. Geophys. Res.*, 110, A09S39, doi:10.1029/2004JA010984, 2005. [4671](#)

10 Rusch, D. W., Gerard, J.-C., Solomon, S., Crutzen, P. J., and Reid, G. C.: The effect of particle precipitation events on the neutral and ion chemistry of the middle atmosphere, 1, Odd nitrogen, *Planet. Space Sci.*, 29, 767–774, 1981. [4674](#)

Semeniuk, K., McConnell, J. C., and Jackman, C. H.: Simulation of the October–November 2003 solar proton events in the CMAM GCM: Comparison with observations, *Geophys. Res. Lett.*, 32, L15S02, doi:10.1029/2004GL022392, 2005. [4677](#)

15 Seppälä, A., Verronen, P. T., Kyrölä, E., Hassinen, S., Backman, L., Hauchecorne, A., Bertaux, J. L., and Fussen, D.: Solar proton events of October–November 2003: Ozone depletion in the Northern Hemisphere polar winter as seen by GOMOS/Envisat, *Geophys. Res. Lett.*, 31, L19107, doi:10.1029/2004GL021042, 2004. [4670](#)

20 Stiller, G. P.: The Karlsruhe Optimized and Precise Radiative Transfer Algorithm (KOPRA), *Wiss. Ber. FZKA 6487*, Forschungszentrum, Karlsruhe, Germany, 521–540, 2000. [4672](#)

Verronen, P. T., Seppälä, A., Kyrölä, E., Tamminen, J., Pickett, H. M., and Turunen, E.: Production of odd hydrogen in the mesosphere during the January 2005 solar proton event, *Geophys. Res. Lett.*, 33, L24811, doi:10.1029/2006GL028115, 2006. [4671](#)

25 von Clarmann, T. and Echle, G.: Selection of optimized microwindows for atmospheric spectroscopy, *Appl. Optics*, 37, 7661–7669, 1998. [4673](#)

von Clarmann, T., Chidziezie Chineke, T., Fischer, H., et al.: Remote sensing of the middle atmosphere with MIPAS, in: *Remote Sensing of Clouds and the Atmosphere VII*, 4882, edited by: Schäfer, K., Lado-Bordowsky, O., Comerón, A., and Picard, R. H., SPIE, Bellingham, WA, USA, 172–183, 2003a. [4672](#)

30 von Clarmann, T., Glatthor, N., Grabowski, U., et al.: Retrieval of temperature and tangent altitude pointing from limb emission spectra recorded from space by the Michelson Interferometer for Passive Atmospheric Sounding (MIPAS), *J. Geophys. Res.*, 108, D23, 4736, doi:10.1029/2003JD003602, 2003b. [4672](#)

**N₂O enhancement
during solar proton
events**

B. Funke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



von Clarmann, T., Glatthor, N., Höpfner, M., et al.: Stratospheric HOCl measurements provide evidence of increased stratospheric odd hydrogen abundances due to solar proton events 2003, J. Geophys. Res., 110, A09S45, doi:10.1029/2005JA011053, 2005. [4671](#)

5 Wennberg, P. O., Anderson, J. G., and Weisenstein, D. K.: Kinetics of reactions of ground state nitrogen atoms ($^4S_{3/2}$) with NO and NO₂, J. Geophys. Res., 99, 18 839–18 846, 1994. [4677](#)

ACPD

8, 4669–4691, 2008

N₂O enhancement during solar proton events

B. Funke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



N₂O enhancement during solar proton events

B. Funke et al.




Fig. 1. Northern Hemisphere distributions of N₂O (in ppbv, parts per billion by volume) for days from 26 October to 11 November 2003 at an altitude of 58 km. Only nighttime data is included. Contours are zonally smoothed within 700 km. Individual measurements are represented by white symbols.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)




Fig. 2. As Fig. 1 but for NO₂.

N₂O enhancement during solar proton events

B. Funke et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



N₂O enhancement during solar proton events

B. Funke et al.




Fig. 3. Northern Hemisphere mean distributions of N₂O (**a**) and NO₂ (**b**) for days from 29 October to 5 November 2003 at an altitude of 58 km. Both day and nighttime data are included. Contours are zonally smoothed within 700 km. Individual measurements are represented by white symbols.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

N₂O enhancement during solar proton events

B. Funke et al.




Fig. 4. Zonal mean cross sections of N₂O in the Northern Hemisphere for the days 26 and 30 October (**a, b**) and 5 and 11 November (**c, d**) 2003 measured in nighttime conditions. The enhancement of N₂O at high latitudes above 40 km is evident. White areas at highest altitudes represent MIPAS measurements with no information and those at the lowest altitudes near the tropics denote values outside of the color scale range.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



N₂O enhancement during solar proton events

B. Funke et al.




Fig. 5. Time series of N₂O abundance after the solar proton events of October–November 2003 for the Northern Hemisphere polar cap (70–90° N) during nighttime conditions. **(a)** MIPAS measurements; where white areas at highest altitudes represent MIPAS measurements with no information and the vertical white bands represent days with no processed data. Lower panels: Simulations by the Canadian Middle Atmosphere Model without **(b)** and with **(c)** application of the averaging kernels of the MIPAS retrievals.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

N₂O enhancement during solar proton events

B. Funke et al.




Fig. 6. Time series of NO₂ enhancement after the solar proton events of October–November 2003 for the Northern Hemisphere polar cap (70–90° N) at nighttime conditions as measured by MIPAS **(a)** and modeled by the Canadian Middle Atmosphere Model (CMAM) **(b)**. The vertical white bands in panel (a) represent days with no processed data.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

N₂O enhancement during solar proton events

B. Funke et al.




Fig. 7. Mean NO, NO₂, and NO_x profiles for the Northern Hemisphere polar cap (70–90° N) at nighttime conditions for the 1 November 2003 measured by MIPAS and modeled by the CMAM model.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

