

This discussion paper is/has been under review for the journal *Atmospheric Chemistry and Physics (ACP)*. Please refer to the corresponding final paper in *ACP* if available.

**Nocturnal isoprene
oxidation**

S. S. Brown et al.

Nocturnal isoprene oxidation over the Northeast United States in summer and its impact on reactive nitrogen partitioning and secondary organic aerosol

S. S. Brown¹, J. A. deGouw^{1,2}, C. Warneke^{1,2}, T. B. Ryerson¹, W. P. Dubé^{1,2}, E. Atlas³, R. J. Weber⁴, R. E. Peltier^{4,*}, J. A. Neuman^{1,2}, J. M. Roberts¹, A. Swanson^{2,5,**}, F. Flocke⁵, S. A. McKeen^{1,2}, J. Brioude^{1,2}, R. Sommariva^{1,2,***}, M. Trainer¹, F. C. Fehsenfeld^{1,2}, and A. R. Ravishankara¹

¹NOAA Earth System Research Laboratory, 325 Broadway, Boulder, CO 80305, USA

²Cooperative Institute for Research in Environmental Sciences, University of Colorado, Boulder, CO 80309, USA

³RSMAS/MAC, University of Miami, 4600 Rickenbacker Causeway, Miami, FL 33149, USA

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Nocturnal isoprene
oxidation**

S. S. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

I◀

▶I

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



⁴School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA 30332, USA

⁵National Center for Atmospheric Research, 1850 Table Mesa Dr., Boulder, CO 80305, USA

*now at: Department of Environmental Medicine, NYU School of Medicine, Tuxedo, NY 10987, USA

**now at: Chemical Technology Department, Northrup Grumman Space Technology, Redondo Beach, CA 90278, USA

***now at: School of Environmental Sciences, University of East Anglia, Norwich, NR4 7TJ, UK

Received: 20 October 2008 – Accepted: 15 November 2008 – Published: 6 January 2009

Correspondence to: S. S. Brown (steven.s.brown@noaa.gov)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Abstract

Isoprene is the largest single VOC emission to the atmosphere. Although it is primarily oxidized photochemically during daylight hours, late-day emissions that remain in the atmosphere at sunset undergo oxidation by NO_3 in regionally polluted areas with large NO_x levels. A recent aircraft study examined isoprene and its nocturnal oxidants in a series of night flights across the Northeast US, a region with large emissions of both isoprene and NO_x . Substantial amounts of isoprene that were observed after dark were strongly anticorrelated with measured NO_3 and were the most important factor determining the lifetime of this radical. The products of photochemical oxidation of isoprene, methyl vinyl ketone and methacrolein, were more uniformly distributed, and served as tracers for the presence of isoprene at sunset, prior to its oxidation by NO_3 . Comparison of a determination of the mass of isoprene oxidized in darkness by NO_3 to a calculation of integrated isoprene emissions showed that large amounts (>20%) of emitted isoprene may undergo nocturnal oxidation in this region. Organic nitrates produced from the NO_3 +isoprene reaction, though not directly measured, were estimated to account for 2–9% of total reactive nitrogen and 7–31% of other long-lived organic nitrates such as PAN. The mass of isoprene oxidized by NO_3 was comparable to and correlated with the organic aerosol loading for flights with relatively low organic aerosol background. The contribution of nocturnal isoprene oxidation to secondary organic aerosol was determined in the range 1–17%, and isoprene SOA mass derived from NO_3 was calculated to exceed that due to OH by approximately 50%.

1 Introduction

Isoprene is the most important biogenically emitted VOC (440–660 Tg C yr⁻¹ globally, Guenther et al., 2006) and is important in regulating levels of tropospheric oxidants such as ozone and OH (Lelieveld et al., 2008; Poisson et al., 2000; Trainer et al., 1987) and in the formation of secondary organic aerosol (Claeys et al., 2004; Henze and

Nocturnal isoprene oxidation

S. S. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Nocturnal isoprene
oxidation**

S. S. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Seinfeld, 2006). Emissions of isoprene from the terrestrial biosphere are dependent on season, temperature and sunlight (Fuentes et al., 2000), such that the majority of isoprene emissions occur at warm temperatures and during daylight hours. Because isoprene emissions are approximately coincident in time with the photochemical generation of OH radicals, and because of the reactivity of isoprene with OH (1.4 h lifetime for $[\text{OH}] = 2 \times 10^6 \text{ molecules cm}^{-3}$, Atkinson and Arey, 2003), the majority of emitted isoprene undergoes rapid photochemical oxidation.

Isoprene that is emitted late in the day, particularly during the late afternoon as OH concentrations begin to fall, will not be fully oxidized photochemically and can remain in the nighttime atmosphere. Ozonolysis typically consumes less than 50%, of this isoprene (Atkinson and Arey, 2003), allowing for overnight transport and oxidation on the following day. In polluted areas however, the presence of NO_x leads to the formation of the nitrate radical, NO_3 , which is also a strong oxidant toward isoprene (Winer et al., 1984). This radical is unstable in sunlight, but at night its presence leads to an estimated lifetime for isoprene that is similar to that predicted for photochemical oxidation (Atkinson and Arey, 2003). If the nocturnal production rate of NO_3 is sufficient, and if the level of NO_x in a nighttime air mass exceeds that of isoprene itself, then reaction with NO_3 will lead to complete consumption of isoprene overnight. The presence of NO_x pollution thus represents an anthropogenic perturbation on the isoprene oxidation cycle.

The size of this NO_x perturbation depends on the fraction of the diurnal isoprene emission that remains in the atmosphere at sunset and the spatial overlap of this isoprene with anthropogenic NO_x emissions. Recent model estimates that consider global or continental averages place the fraction of total isoprene emissions that are oxidized by NO_3 at 6% of North American (Horowitz et al., 2007) and 6–7% of global isoprene emissions (Ng et al., 2008). Even though these figures are a small fraction of total isoprene emissions, they nevertheless represent a large mass on a global scale (e.g. 22–29 Tg of isoprene consumed by this reaction annually, Ng et al., 2008). If such estimates are correct, they represent a significant fraction of oxidized nitrogen con-

sumption as well (4.5–6.0 Tg N annually, or 12–16% of year 2000 global anthropogenic NO_x emissions, Olivier and Berdowski, 2001).

Despite the large isoprene mass oxidized via NO₃ reaction, the resulting products have not been widely studied. The reaction is known to produce unsaturated organic nitrates with a yield of approximately 80% as first-generation products (Barnes et al., 1990). These nitrates consist of unsaturated hydroxy-, hydroperoxy- and carbonyl nitrates (Kwok et al., 1996; Skov et al., 1992, 1994; Tuazon et al., 1999; Berndt and Boge, 1997), with a recent study also showing evidence for formation of dinitrates and oligomers (Ng et al., 2008). The subsequent fates of such compounds are poorly understood and are an important uncertainty in isoprene oxidation models (Fan and Zhang, 2004; Paulson and Seinfeld, 1992). These uncertainties may reduce the accuracy of modeled transport and/or loss of reactive nitrogen or of modeled surface level ozone (Horowitz et al., 2007; von Kuhlmann et al., 2004). Recently, Ng et al. (2008) measured mass yields of secondary organic aerosol from the oxidation of isoprene by NO₃ to be in the range 5–25%, much larger than that from OH (1–3%, Kroll et al., 2005, or O₃<1%, Kleindienst et al., 2007) reactions, and estimated the contribution of this reaction to global secondary organic aerosol production at 2–3 Tg yr⁻¹.

Several recent surface-level field studies have examined nocturnal isoprene oxidation in both polluted and remote forested environments. Although nighttime decays of surface-level isoprene are well-documented (Hurst et al., 2001; Trainer et al., 1991), their interpretation is often complicated by the competing effects of chemistry and transport, particularly within shallow nocturnal boundary layers (Apel et al., 2002). For example, measurements at forested sites impacted by urban NO_x emissions have in several instances found evening isoprene loss rates consistent with consumption by NO₃ radicals (Starn et al., 1998; Steinbacher et al., 2005; Stroud et al., 2002), while analysis of data from more remote locations have shown the importance of either transport or, possibly, additional oxidants such as nighttime OH (Biesenthal et al., 1998; Sillman et al., 2002). Analyses of surface level NO₃ measurements in regions downwind of forested areas have shown that biogenic VOCs, including isoprene and monoter-

**Nocturnal isoprene
oxidation**

S. S. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



penes, consume a significant fraction of NO₃ radicals (Aldener et al., 2006; Ambrose et al., 2007; Geyer et al., 2001; McLaren et al., 2004). A ship-based study in coastal New England also found that reaction with NO₃ was responsible for some 40% of the loss of isoprene advected over water from mixed urban/forested areas (Warneke et al., 2004).

Here, we report nocturnal observations of isoprene, its oxidation products and its major nighttime oxidants from the NOAA P-3 aircraft. Table 1 lists the trace gas and aerosol measurements relevant for this analysis. This study differs from surface level measurements at a fixed location in that it provides a regional analysis of the degree to which isoprene is oxidized at night. Furthermore, the nighttime aircraft observations sample primarily above the nocturnal boundary layer and are therefore decoupled from surface emissions, nocturnal boundary layer dynamics and deposition. This study examines the fate of isoprene that is transported or oxidized aloft within the residual, daytime boundary layer, where the majority of the daytime-emitted isoprene remaining at sunset resides. These measurements were part of the 2004 New England Air Quality Study (NEAQS), part of the larger Intercontinental Consortium for Atmospheric Research on Transport and Transformation (ICARTT) (Fehsenfeld et al., 2006), which took place in July and August in the Northeast US. Figure 1 shows the nighttime flight tracks of the P-3 superimposed on an isoprene emissions inventory map for this region (Guenther et al., 2006). These nighttime flights sampled in urban and forested regions across the Northeast US, an area with both large isoprene and NO_x emissions, where NO₃-driven isoprene oxidation is likely to be particularly important.

2 NO_x enhancement of nocturnal alkene ozonolysis

Ozone and NO₃ are both highly reactive toward unsaturated VOCs and are the most important nocturnal oxidants for isoprene. While ozone is more uniformly distributed in the troposphere and is thought to be a more important oxidant for isoprene globally, reaction with NO₃ can easily dominate the nighttime chemical loss for isoprene on a

Nocturnal isoprene oxidation

S. S. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



regional scale. Because NO_3 also requires ozone for its production, its further reaction with isoprene is equivalent to an enhancement to the background rate of nocturnal isoprene ozonolysis.



$$- \frac{d[\text{Isop}]}{dt} = (k_1[\text{O}_3] + k_3[\text{NO}_3])[\text{Isop}] \quad (1)$$

Quantities in square brackets indicate concentrations in number density units. In the early evening, when isoprene concentrations over a forested areas are large, isoprene will be the dominant reaction partner for NO_3 radicals produced in Reaction (2). Under these conditions, NO_3 will rapidly achieve a steady state between Reactions (2) and (3) such that its concentration can be approximately written as the ratio of its production rate and first-order loss rate coefficient (Platt et al., 1984).

$$[\text{NO}_3] \approx \Phi_{\text{Isop}} \frac{k_2[\text{O}_3][\text{NO}_2]}{k_3[\text{Isop}]} \quad (2)$$

Here Φ_{Isop} is the fraction of NO_3 radicals produced in Reaction (2) that react with isoprene (i.e., the ratio of the first order loss rate coefficient of NO_3 with isoprene to the sum of all first order loss rate coefficients of NO_3 radicals, including indirect loss by hydrolysis of N_2O_5). Substitution of Eq. (2) into Eq. (1) yields a simple result that demonstrates the enhancement of the background isoprene ozonolysis rate in the presence of NO_x .

$$- \frac{d[\text{Isop}]}{dt} \approx (k_1[\text{Isop}] + \Phi_{\text{Isop}}k_2[\text{NO}_2])[\text{O}_3] \quad (3)$$

Nocturnal isoprene oxidation

S. S. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Nocturnal isoprene oxidation

S. S. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



In order for the NO_3 reaction to consume a large fraction of the available isoprene, the concentration of NO_2 must exceed that of isoprene such that NO_x is not the limiting reagent. For $[\text{NO}_2] \geq [\text{Isop}]$, the second term in Eq. (3) exceeds the first by at least a factor of 2.8 at 298 K (for $\Phi_{\text{Isop}} \approx 1$) (Atkinson and Arey, 2003), such that the NO_3 reaction always dominates if there is sufficient NO_2 available (e.g., Sommariva et al., 2008).

Figure 2 shows the eastern part of the P-3 track after sunset on the night of 9–10 August 2004, color and size coded by the mixing ratios of both NO_3 and isoprene. (The western side of the track over western Pennsylvania and Ohio was characterized by lower isoprene and large sulfate aerosol loading, such that reactivity of NO_3 and N_2O_5 was dominated by N_2O_5 uptake to particles rather than by NO_3 isoprene chemistry, Brown et al., 2006b). Comparison of the two maps shows that the spatial distribution of NO_3 and isoprene were strongly anticorrelated, with NO_3 present at large mixing ratios (up to 350 pptv) downwind of major urban areas, and isoprene present in all areas not impacted by rapid production of NO_3 . The scatter plot in the lower panel of Fig. 2 shows this anticorrelation, which was a persistent feature of all night flights in which measurable isoprene was present. As the next section demonstrates, the anticorrelation is not simply the result of the spatial distribution of isoprene and NO_x emissions, but rather due to oxidation of a diffusely distributed background of isoprene in areas where the NO_x levels were sufficient to rapidly consume isoprene in the hours just after sunset.

Figure 3 shows the time series of isoprene and the first-order loss rate coefficients for NO_3 radicals for the portion of the flight tracks shown in Fig. 2. The figure compares two different calculations of this rate coefficient. The first is the ratio of the production rate for NO_3 from Reaction (2) to the observed NO_3 concentration.

$$\tau(\text{NO}_3)^{-1} = \frac{k_2[\text{O}_3][\text{NO}_2]}{[\text{NO}_3]} \quad (4)$$

This quantity, referred to as the inverse steady state lifetime (Platt et al., 1984) or the loss frequency (Geyer and Platt, 2002) for NO_3 , is a measure of the total first-order loss

Nocturnal isoprene oxidation

S. S. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



rate coefficient for NO_3 radicals under conditions where the NO_3 and N_2O_5 system has achieved a steady state with respect to production from Reaction (2) and losses to reactions of both NO_3 and N_2O_5 (generally valid for conditions of rapid sinks for NO_3 and N_2O_5 , warmer temperatures, and lower NO_2 concentrations, Allan et al., 2000; Brown et al., 2003). At steady state, the following approximate relationship between Eq. (4) and NO_3 sinks holds.

$$\tau(\text{NO}_3)^{-1} \approx \sum_i k_i[\text{VOC}_i] + K_{eq}[\text{NO}_2]k_{\text{hydrolysis}} \quad (5)$$

The first term on the right hand side above is a summation of the products of bimolecular rate coefficients and VOC concentrations for all VOCs that are reactive with NO_3 . The second term is the indirect loss rate coefficient for NO_3 radicals due to hydrolysis of N_2O_5 , with a first-order loss rate coefficient to aerosols $k_{\text{hydrolysis}}$. K_{eq} is the temperature dependent equilibrium constant for the reversible reaction of NO_3 with NO_2 to form N_2O_5 . This expression neglects daytime reactions, such as photolysis of NO_3 and reaction of NO_3 with NO , and heterogeneous uptake of NO_3 (although the latter could easily be incorporated into the summation).

The second calculation of the NO_3 first order loss rate coefficient in Fig. 3 is the loss of NO_3 due only to reaction with isoprene.

$$k_l(\text{Isoprene}) = k_3[\text{Isop}] \quad (6)$$

Here, k_3 is the bimolecular rate coefficient for Reaction (3). Although Eq. (6) represents only a single term from the summation in Eq. (5), the correspondence of the time series between the two independently measured quantities in Fig. 3 is remarkable, and was a consistent feature of all NEAQS night flights. The plot of $\tau(\text{NO}_3)^{-1}$ vs. $k_l(\text{Isoprene})$ in Fig. 4 shows that for isoprene levels greater than 50 pptv, loss to isoprene accounted for $73 \pm 8\%$ of the NO_3 loss determined from $\tau(\text{NO}_3)^{-1}$. This observation suggests not only that the approximate isoprene loss rate in Eq. (3) is valid, but that the value of Φ_{Isop} can be taken as ~ 0.73 . The remaining consumption of NO_3 radicals is not

defined here, although possibilities include reactions with other biogenic VOC, second generation isoprene oxidation products, peroxy radicals and heterogeneous uptake of either NO_3 or N_2O_5 (although the latter was negligible for the eastern leg of the 9–10 August flight, Brown et al., 2006b).

5 This analysis shows that isoprene, when present within the residual daytime boundary layer (observed here to be approximately 0.1–1.5 km, Brown et al., 2007), controls the loss of NO_3 and therefore the consumption of NO_x at night. Furthermore, a large fraction of the isoprene present at sunset within NO_x – containing air masses is lost to reaction with NO_3 rather than reacting with ozone or being transported overnight. The
10 next section outlines a method by which isoprene and NO_2 at sunset may be estimated for nocturnally sampled air masses. Sunset NO_2 was nearly always in excess over isoprene, such that the second term in Eq. (3) was dominant. In fact, comparison of the two terms in Eq. (3) at sunset shows that well over 90% of the isoprene oxidized in the dark was lost to reaction with NO_3 rather than ozone. Thus, in an area with large
15 regional NO_x and isoprene emissions such as the Northeast US, NO_x emissions are a determining factor for isoprene loss budgets overnight.

3 Relationship between isoprene and MVK+MACR at night

The products of the photochemical oxidation of isoprene, methyl vinyl ketone and methacrolein (MVK+MACR), serve as tracers for the presence of isoprene within the residual daytime boundary layer at sunset. These compounds are produced in high
20 yields from the atmospheric oxidation of isoprene by either OH or O_3 (Atkinson and Arey, 2003) but at low yields (3.5%) in the NO_3 oxidation of isoprene (Kwok et al., 1996). Nighttime oxidation of isoprene by ozone produces MVK+MACR, but much more slowly than does OH oxidation, and their production is nearly negligible if most of
25 the nighttime oxidation occurs via NO_3 . Furthermore, these compounds are much less reactive with either O_3 or NO_3 than isoprene itself. Surface level measurements have shown that MVK+MACR change only slowly at night while isoprene decays rapidly in

Nocturnal isoprene oxidation

S. S. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Nocturnal isoprene
oxidation**

S. S. Brown et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

air masses with a large NO_3 production rate (Stroud et al., 2002). Because methyl vinyl ketone and methacrolein have the same mass, they are measured as a single sum by the PTRMS instrument. Canister samples analyzed by gas chromatography provide a speciation of the two compounds that can be useful, for example, in analyzing their individual production and loss. Canister samples also showed agreement with the PTRMS measurement for the MVK+MACR sum. The analysis in this paper uses the sum derived from the PTRMS at higher time resolution.

Figure 5 shows maps of the night flight on 3 August 2004, color and size coded by NO_3 , isoprene and MVK+MACR. The spatial distribution of isoprene and NO_3 is similar to that for the 9–10 August flight shown in Fig. 2, and the scatter plot of isoprene against NO_3 again shows the same anticorrelation observed between these compounds that was characteristic of the night flights. The spatial distribution of MVK+MACR is considerably different, however. This tracer is more uniformly distributed than isoprene and appears in areas where isoprene is absent but NO_3 is present at large mixing ratios. The observed anticorrelation between NO_3 and isoprene at night was therefore not simply a result of spatial separation between NO_x and isoprene emissions, but rather due to oxidation of isoprene in regions where NO_x emissions provided a sufficient NO_3 source.

4 Isoprene oxidized after dark

The amount of isoprene that had been present at sunset in air masses sampled after dark may be calculated directly from integration of Eq. (3) backwards in time. This analysis is only possible for air samples in which measurable isoprene remains in an air mass at night because the isoprene concentration itself appears in this equation and because the approximation relies on the assumption that isoprene is the major reaction partner for NO_3 . It further relies on the assumption that air masses sampled aloft at night were decoupled from surface emissions trapped within the shallow nocturnal boundary layer after sunset. The integral has an analytical solution under the

Nocturnal isoprene oxidation

S. S. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



assumption of constant ozone over the period since sunset; the validity of this approximation depends on the amount of ozone consumed in Reaction (2) and is appropriate for ozone in large excess over NO_2 , which was generally the case for the analysis presented here ($\text{O}_3 \sim 20\text{--}100 \times \text{NO}_2$). Because N_2O_5 hydrolysis is unimportant in comparison to the reaction of NO_3 with isoprene for large isoprene levels, the rate of NO_2 consumption is equal to the rate of Reaction (2) (rather than being twice this rate, as would be the case if N_2O_5 hydrolysis were dominant, Brown et al., 2004). However, the NO_3 +isoprene reaction regenerates a fraction of NO_2 with a yield, Φ_{NO_2} , of 20%; the remainder of the nitrogen goes to organic nitrates (Barnes et al., 1990).



With these assumptions, the integral of Eq. (3) over the time interval to sunset (taken here as solar zenith angle of 90° at the sampling location) has an analytical solution for the initial isoprene.

$$15 \quad [\text{Isop}]_0 = [\text{Isop}] e^{k_1[\text{O}_3]\Delta t} + \frac{\Phi_{\text{Isop}} k_2 [\text{NO}_2] e^{\Phi_S k_2 [\text{O}_3] \Delta t}}{\Phi_S k_2 - k_1} (1 - e^{(k_1 - \Phi_S k_2)[\text{O}_3] \Delta t}) \quad (7)$$

$$\Phi_S = 1 - \Phi_{\text{Isop}} \Phi_{\text{NO}_2} \quad (8)$$

Here Δt is the time since sunset, $[\text{Isop}]$, $[\text{NO}_2]$ and $[\text{O}_3]$ are the observed isoprene, NO_2 and O_3 concentrations, k_1 and k_2 are the bimolecular rate coefficients for the reaction of O_3 with isoprene and NO_2 , respectively, and Φ_{Isop} is the fraction of NO_3 radicals produced in Reaction (2) that react with isoprene (taken as a constant $\Phi_{\text{Isop}} = 0.73$, see above).

Figure 6 shows initial isoprene calculated from Eq. (7) for the 3 August flight as a time series along with observed isoprene and MVK+MACR. The lower panel of this figure gives the correlation between calculated initial isoprene and observed MVK+MACR.

**Nocturnal isoprene
oxidation**

S. S. Brown et al.

The correlation between derived sunset isoprene and observed MVK+MACR on the 3 August flight (correlation coefficient=0.72, see Table 2) was higher than for other flights. For this flight, MVK+MACR has been used to estimate sunset isoprene even in air masses where the isoprene had already been completely lost to oxidation. The slope of the correlation plot in Fig. 6 is 0.6 ± 0.3 , where the $\pm 50\%$ error bars, shown as dashed lines on the plot represent limits that encompass the majority of the data. The ratio of isoprene to MVK+MACR at sunset is expected to be variable since it depends on the time of emission and the photochemical processing time prior to sunset. Measured values of this ratio for late day measurements (within 3 h prior to 0.5 h after sunset) from NEAQS 2004 gave an average value of 0.4 ± 0.3 ($\pm 1\sigma$). Values from surface level measurements, on the other hand, tend to be somewhat larger, approximately 1:1 (e.g., Stroud et al., 2002). The correlation between derived sunset isoprene and MVK+MACR for the 3 August flight, and the large late-day isoprene emission on this flight (see below), indicate that the use of MVK+MACR as a semi-quantitative isoprene tracer is appropriate for this flight.

Figure 7 shows the estimate for isoprene oxidized after dark for the 3 August flight. As described above, this estimate is taken as a constant ratio of 0.6 ± 0.3 times the MVK+MACR data, minus the measured isoprene still present in the air mass. Maximum values for this estimate are in the range of 1–2 ppbv. Also shown is the integrated isoprene emission into each sampled air mass on the preceding day, calculated using a Lagrangian particle dispersion model (FLEXPART) (Stohl et al., 2005). FLEXPART was driven by model-level data from the European Centre for Medium-Range Weather Forecasts (ECMWF) with a temporal resolution of 3 h (analyses at 00:00, 06:00, 12:00, 18:00 UTC ; 3-h forecasts at 03:00, 09:00, 15:00, 21:00 UTC) and 91 vertical levels and a horizontal resolution of 0.360×0.360 . Isoprene emissions were taken from BEIS3.12. The temperature and light dependence of isoprene was calculated hourly for each isoprene emission grid with the EPA recommended canopy environment model by using the ECMWF 2 m temperature and net solar radiation, interpolated linearly in space and time using the two nearest ECMWF fields.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Nocturnal isoprene
oxidation**

S. S. Brown et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

One calculation shows isoprene emissions for the preceding daylight hours, and a second shows emissions for only the 6 h preceding the time of sampling. These late day emissions were calculated to be a large fraction of the total, reflecting the air mass history. In the early part of the flight, backward trajectories had passed over the isoprene-rich areas of western Massachusetts, as illustrated in the map in the lower panel of the figure. For air masses sampled below the top of the residual daytime boundary layer on this flight (<1.5 km), the average ratio of the isoprene oxidized after dark to the total emissions was $22\pm 11\%$, although values over Massachusetts were larger. The estimated isoprene oxidized after dark was a large fraction of the isoprene emitted within the preceding 6 h of the sample time in the early part of the flight. The flight began approximately two hours after local sunset, so that in the early part of the flight, the calculated 6 h isoprene emissions, which are sunlight dependent, would have occurred within 2–4 h of sunset.

The fraction of isoprene oxidized after dark, determined from the 3 August flight can be compared to models of North American isoprene oxidation. For example, a recent calculation of the fraction of total North American isoprene emissions that are oxidized by NO_3 puts this figure at 6% (Horowitz et al., 2007). Within a domain that includes only the Northeast US, encompassing the land areas sampled by the flight tracks shown in Fig. 1, a WRF-CHEM (Weather Research Forecast) model simulation for a single summer day predicts 8% of total isoprene emissions oxidized by NO_3 for clear sky conditions, and 20% for cloudy conditions. These WRF-CHEM runs use the same domain, photochemistry, updated 2004 anthropogenic emissions and physics options as those documented in Kim et al. (2006), however they are based on a newer release of WRF and WRF-CHEM (ARW version 3.0, Skamarock et al., 2008). The value of $22\pm 11\%$ is larger than these domain-wide average estimates, likely because of sampling in an air mass with large late-day isoprene emissions in close proximity to NO_x pollution sources. This relatively larger fraction of isoprene emissions that are oxidized by NO_3 inferred from some of the NEAQS flights has consequences for ozone, reactive nitrogen and aerosol growth, as outlined in the next sections.

5 Reactive nitrogen partitioning

The products of the NO_3 +isoprene reaction are primarily organic nitrates that may contribute measurably to total reactive nitrogen (NO_y). Starn et al. (1998) estimated NO_3 -derived isoprene nitrates to be 8% of total reactive nitrogen at an NO_x impacted forested site. Direct measurements of the organic nitrates from nocturnal isoprene oxidation, however, are sparse. Ballschmitter and coworkers measured these compounds by chromatographic methods in remote areas (e.g., marine environments) where there were only very small concentrations (Fischer et al., 2000; Werner et al., 1999). Shepson and coworkers reported diurnal patterns for these compounds in forested regions in Michigan and Tennessee (Giacopelli et al., 2005; Grossenbacher et al., 2001, 2004) but focused mainly on the photochemical production from reactions of isoprene peroxy radicals with NO . For the P-3 flights during NEAQS 2004, there were no specific measurements of isoprene nitrates.

Figure 8 shows a time series of different components of reactive nitrogen, including HNO_3 , PAN (shown here as the sum of speciated PANs measured with a chemical ionization mass spectrometer, see Table 1), NO_x , N_2O_5 and isoprene nitrates from isoprene+ NO_3 . The isoprene nitrate trace is the product of the estimate for total isoprene reacted after dark and the laboratory organic nitrate yield of 80% (Barnes et al., 1990). It does not include an estimate for isoprene nitrates derived from the photochemical reactions of isoprene peroxy radicals with NO , which are thought to be between approximately 4–10% of the isoprene oxidized during daytime under high NO_x conditions (Chen et al., 1998; Sprengnether et al., 2002). Although regional models predict approximately equal production of isoprene nitrates from daytime and nighttime mechanisms (Horowitz et al., 2007), for the air masses sampled on the 3 August flight, where a large fraction of the emitted isoprene were shown to have been oxidized after dark, isoprene nitrates derived from NO_3 should dominate over their photochemical analogs. Also shown in Fig. 8 is the sum of reactive nitrogen, denoted here as NO_y Sum. This sum is not a direct measurement from an NO_y instrument (not available

Nocturnal isoprene oxidation

S. S. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



for these flights) but from the sum of all of the compounds shown in the time series, including $2 \times \text{N}_2\text{O}_5$. The middle panel of Fig. 8 shows the partitioning among these compounds, and the two pie charts at the bottom show the average partitioning for the early part of the flight, over the isoprene rich areas of New Hampshire, Massachusetts and Rhode Island, and for the entire flight.

Nitric acid was the dominant component of reactive nitrogen on this and other night flights during this campaign. Because the aircraft samples from the residual daytime boundary layer, which decouples from the surface at night, soluble gases such as HNO_3 do not undergo efficient loss to dry deposition, and are longer-lived than on daytime flights in well mixed boundary layers (Neuman, 2006). Analysis of HNO_3 - O_3 correlations (Brown et al., 2006a) and steady state lifetimes of NO_3 and N_2O_5 (Brown et al., 2006b) (not shown) showed that much of the HNO_3 in the early part of the flight had been formed photochemically from reaction of NO_2 with OH , while the plume over New York City sampled later in the flight had a large contribution from N_2O_5 hydrolysis. The dominance of HNO_3 resulting from N_2O_5 hydrolysis within the New York City plume reduced the relative contribution of isoprene nitrates to NO_y , although the total mass of these compounds was still similar to the earlier part of the flight.

The calculated isoprene nitrates from nocturnal oxidation represent $9 \pm 5\%$ and $4 \pm 2\%$ of total reactive nitrogen in the isoprene-rich areas and in the New York plume, respectively. In the isoprene-rich areas, this contribution was slightly less than half of PAN, the longest-lived component of NO_y . Therefore, if the NO_3 -derived isoprene nitrates were as long-lived as PAN compounds, they could represent a large fraction of NO_y as the air mass ages and as NO_x and HNO_3 are lost to oxidation and deposition.

The importance of isoprene nitrates to reactive nitrogen partitioning at night can be demonstrated by examining only the nighttime NO_x oxidation products. Figure 9 is a plot of calculated isoprene nitrates against the calculated amount of NO_2 oxidized after dark (i.e., $\Delta[\text{NO}_2] = [\text{NO}_2](\exp(k_1[\text{O}_3]\Delta t_{\text{sunset}}) - 1)$). The slope is 0.56, indicating that about half of the NO_2 consumed in darkness was likely present as isoprene nitrates. Although the NO_3 -derived isoprene nitrates were generally less than 10% of total NO_y ,

**Nocturnal isoprene
oxidation**

S. S. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



they were approximately half of the NO_x consumption after dark in this air mass. Similar analyses hold for the other night flights analyzed for this campaign, as shown in Table 3. These observations are in part a consequence of measurements from aircraft in the residual daytime boundary layer, which are isolated from NO_x emissions occurring below the nocturnal boundary layer and are thus likely to show evidence for a larger fraction of photochemical NO_y oxidation products, such as PAN and photochemically generated HNO_3 . At lower altitudes, particularly the upper portion of the nocturnal boundary layer, the oxidation products of the nighttime reactions may represent a larger fraction of NO_y (Brown et al., 2007; Stutz et al., 2004).

6 Secondary organic aerosol

The production of secondary organic aerosol (SOA) is a topic of considerable current interest. For example, analyses of VOC and aerosol data in the same region of the Northeast US have shown that organic aerosol is well-correlated with tracers for anthropogenic emissions, such as acetylene, but that budgets of organic aerosol production are poorly represented by current understanding based on yields for individual VOCs measured in smog chambers (de Gouw et al., 2005, 2008; Quinn et al., 2006; Sullivan et al., 2006). The contribution of biogenic VOCs to organic aerosol production remains uncertain, although recent analysis of radiocarbon in organic aerosol has suggested that a large fraction comes from biogenic sources, even in areas with large anthropogenic VOC emissions (Bench et al., 2007; Schichtel et al., 2008; Weber et al., 2007). Because the yield of SOA from the NO_3 +isoprene reaction (Ng et al., 2008) is some 5–8× larger than from photochemical isoprene oxidation (Kroll et al., 2005), its nighttime oxidation is a mechanism by which biogenic VOCs can more efficiently form SOA in polluted atmospheres. Furthermore, since NO_3 is derived from NO_x and is an anthropogenic oxidant, aerosol produced via this oxidation would correlate with anthropogenic tracers, but maintain a biogenic carbon signature.

Figure 10 shows the time series for organic aerosol, measured as water soluble

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Nocturnal isoprene
oxidation**

S. S. Brown et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

organic carbon (WSOC) with a particle into liquid sampler (PiLS) (Table 1), for the 3 August flight. Shown together on the same scale is the calculated amount of isoprene oxidized after dark, converted to the same units ($\mu\text{g C m}^{-3}$). The two quantities are similar in magnitude (oxidized isoprene was 65% of WSOC) and reasonably well correlated, as the lower left plot shows. It is important to note, however, that WSOC was also well correlated with other anthropogenic traces (e.g., CO) such that the correlation in the lower left plot of Fig. 10 does not suggest that the majority of SOA was attributable to nighttime isoprene oxidation. Indeed, the smaller, shaded region in the upper graph of Fig. 10 shows nocturnally oxidized isoprene multiplied by the range of SOA yields for NO_3 +isoprene from the recent chamber study (Ng et al., 2008). The range from the chamber study is given as aerosol mass produced per unit mass of parent hydrocarbon oxidized. Conversion from total mass to carbon mass requires the molecular weight of the condensing species. Under the assumption that the condensable products are carbonyl nitrates, with a formula $\text{C}_5\text{H}_7\text{O}_4\text{N}$, the organic mass yield of 5–25% from the smog chamber is equivalent to an organic carbon yield of 2.5–12.5%. The predicted nocturnal isoprene SOA is plotted against observed WSOC in the lower right hand plot. The slopes of this graph give a range of 2–8% of measured WSOC attributable to nocturnally oxidized isoprene.

Although this analysis shows only a small contribution (typically <10%) of nighttime isoprene oxidation to total observed organic aerosol, the SOA mass derived from nighttime isoprene oxidation was large in comparison to that predicted from daytime isoprene oxidation. Figure 11 shows this calculation, again using the 3 August flight as an example. Although the amount of isoprene oxidized after dark was smaller than that oxidized photochemically, the laboratory mass yields for NO_3 -isoprene reactions are larger than those for OH- or O_3 -isoprene reaction, such that in this regionally polluted environment NO_3 was calculated to have made a larger contribution. Figure 11 shows the isoprene SOA mass (as total rather than carbon mass) from each oxidant averaged for each of the sampled air masses on 3 August. The NO_3 -isoprene SOA mass is the product of the isoprene mass oxidized after dark by NO_3 and the labora-

**Nocturnal isoprene
oxidation**

S. S. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tory SOA mass yield, taken here as $15\pm 10\%$ (for the 5–25% range given) (Ng et al., 2008). Similarly, the OH-isoprene SOA mass is the difference between the FLEXPART integrated isoprene emission for each air mass and the amount oxidized after dark times the laboratory yield of $2\pm 1\%$ (range of 0.9–3%) (Kroll et al., 2005). Finally, the O_3 -isoprene SOA mass is taken as 5% of the total isoprene emission times its laboratory SOA yield (1%) (Kleindienst et al., 2007). The contribution of NO_3 to the total isoprene SOA is approximately 50% greater than OH in this regionally polluted area. However, this does not consider other possible routes for SOA formation from isoprene, such as heterogeneous reactions (Altieri et al., 2006; Ervens et al., 2008; Liggio et al., 2007).

Analysis of several other flights showed correlations between nocturnally oxidized isoprene and observed WSOC, non-negligible contributions of NO_3 -isoprene SOA to total organic aerosol, and large nighttime isoprene-SOA compared to photochemical isoprene-SOA, as shown in Table 3. Figure 12 gives one additional example, from the night flight on 7 August, which sampled mainly along the coastline from New Jersey to Massachusetts under conditions of offshore flow. In this case, there was a small organic aerosol loading, $0.5\text{--}1.5\ \mu\text{g C m}^{-3}$, but it was again well correlated and of the same magnitude as the isoprene oxidized after dark. The fraction of secondary organic aerosol attributable to oxidized isoprene was 2–10%. Analysis of the 31 July flight showed a similar result, although the determination of isoprene oxidized after dark was much less certain (see Table 2). By contrast, flights in which the background organic aerosol loading was already substantial (e.g., 11 July, 9 August) showed only small amounts of observed organic aerosol attributable to nocturnal isoprene oxidation (<4%), even though there was a comparable amount of isoprene oxidized on these flights. In these cases, the correlations between oxidized isoprene and WSOC were poor. These observations suggest that the signature of nocturnally-produced isoprene SOA is most apparent when the background aerosol loading is small. Photochemical production of organic aerosol from anthropogenic VOC on the preceding day has the capacity to produce much larger amounts of organic aerosol than nighttime processes,

so that the influence of nighttime chemistry on SOA becomes less apparent if the daytime, background SOA loading is large. Nevertheless, the derived isoprene-SOA from NO_3 , ranging up to 17% of observed organic aerosol and in many cases dominating the calculated SOA derived from isoprene (see Table 3), is a source that may need to be accounted for in models if these results are borne out by further work and if current understanding of SOA yields is correct.

7 Summary and conclusions

This paper has examined nighttime oxidation of isoprene in air masses sampled from an aircraft within the residual daytime boundary layer during the summer of 2004 in the Northeast US. Collocation of NO_x and VOC emissions within this region leads to an increased contribution of NO_3 to total isoprene oxidation. Reaction of isoprene with NO_3 is kinetically equivalent to an enhancement in the rate of ozonolysis since the NO_3 oxidant is itself derived from ozone. The NO_3 reaction differs considerably from ozonolysis, however, in that it has the potential to rapidly consume isoprene in the hours after sunset and thus prevent its overnight transport. During this study, mixing ratios of isoprene in excess of 1 ppbv were observed aloft after dark and were strongly anticorrelated with NO_3 . Further, isoprene was found to be the most important factor determining the lifetime of NO_3 for isoprene levels in excess of 50 pptv. The mass of isoprene oxidized at night was estimated by comparison to its photochemical oxidation products, methyl vinyl ketone and methacrolein, which served as tracers. These estimates showed that up to ~20% of the total isoprene emissions were oxidized after dark within the sampled air masses. The resulting isoprene nitrates were calculated to be 2–9% of total reactive nitrogen and 7–31% of longer-lived organic nitrates, such as PAN. Finally, in air masses with relatively small backgrounds of organic aerosol, nocturnally oxidized isoprene was well correlated with measured water soluble organic carbon. Estimation of nocturnally oxidized isoprene SOA using yields from recent smog chamber measurement showed it to be in the range of 1–17% for low organic aerosol

Nocturnal isoprene oxidation

S. S. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



backgrounds. Isoprene-SOA derived from NO₃ reaction was found to be approximately 50% larger than that from photochemical isoprene oxidation within these air masses.

Results from this study are applicable to isoprene oxidation occurring at night within the daytime residual boundary layer (~0.1–1.5 km) during summertime in a polluted, forested region. Other biogenic VOCs, particularly monoterpenes, were not addressed here because they were not observed in large concentration from the aircraft. However, nocturnal oxidation of these hydrocarbons is rapid via both O₃ and NO₃, such that their oxidation rates and production of organic nitrates and secondary organic aerosol should be strongly influenced by regional NO_x pollution (Fry et al., 2008; Hallquist et al., 1999). Because monoterpene oxidation is temperature but not sunlight dependent, and because these compounds are more reactive with both ozone and NO₃, the signature of their oxidation should be most apparent within the nocturnal boundary layer. As a result, their nocturnal oxidation products may be both more concentrated and more vertically stratified. Study of these oxidation processes will be of considerable future interest but may require vertically resolved measurements above NO_x-impacted forest canopies.

Acknowledgements. This work was supported by the NOAA Atmospheric Chemistry and Climate Program and the New England Air Quality Study.

References

- Aldener, M., Brown, S. S., Stark, H., Williams, E. J., Lerner, B. M., Kuster, W. C., Goldan, P. D., Quinn, P. K., Bates, T. S., Fehsenfeld, F. C., and Ravishankara, A. R.: Reactivity and loss mechanisms of NO₃ and N₂O₅ in a marine environment: results from in-situ measurements during NEAQS 2002, *J. Geophys. Res.*, 111, D23S73, doi:10.1029/2006JD007252, 2006.
- Allan, B. J., McFiggans, G., Plane, J. M. C., Coe, H., and McFadyen, G. G.: The nitrate radical in the remote marine boundary layer, *J. Geophys. Res.*, 105, 24191–24204, 2000.
- Altieri, K. E., Carlton, A. G., Lim, H.-J., Turpin, B. J., and Seitzinger, S. P.: Evidence for oligomer formation in clouds: Reaction of isoprene oxidation products, *Environ. Sci. Technol.*, 40, 4956–4960, 2006.

Nocturnal isoprene oxidation

S. S. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Nocturnal isoprene
oxidation**

S. S. Brown et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Ambrose, J. L., Mao, H., Mayne, H. R., Stutz, J., Talbot, R., and Sive, B. C.: Nighttime nitrate radical chemistry at Appledore Island, Maine during the 2004 International Consortium for Atmospheric Research on Transport and Transformation, *J. Geophys. Res.*, 112, D21302, doi:10.1029/2007JD008756, 2007.
- 5 Apel, E., Riemer, D. D., Hills, A., Baugh, W., Orlando, J., Faloon, I., Tan, D., Brune, W., Lamb, B., Westberg, H., Carroll, M. A., Thornberry, T., and Geron, C.: Measurement and interpretation of isoprene fluxes and isoprene, methacrolein, and methyl vinyl ketone mixing ratios at the PROPHET site during the 1998 Intensive, *J. Geophys. Res.*, 107, 4034, doi:10.1029/2000JD000225, 2002.
- 10 Atkinson, R. and Arey, J.: Gas-phase tropospheric chemistry of biogenic volatile organic compounds: a review, *Atmos. Environ.*, 37, S197–S219, 2003.
- Barnes, I., Bastian, V., Becker, K. H., and Tong, Z.: Kinetics and Products of the Reactions of NO₃ with Monoalkenes, Dialkenes, and Monoterpenes, *J. Phys. Chem.*, 94, 2413–2419, 1990.
- 15 Bench, G., Fallon, S., Schichtel, B., Malm, W., and McDade, C.: Relative contributions of fossil and contemporary carbon sources to PM_{2.5} aerosols at nine Interagency Monitoring for Protection of Visual Environments (IMPROVE) network sites, *J. Geophys. Res.*, 112, D10205, doi:10.1029/2006JD007708, 2007.
- Berndt, T. and Boge, O.: Gas-phase reaction of NO₃ radicals with isoprene: A kinetic and mechanistic study, *Int. J. Chem. Kinet.*, 29, 755–765, 1997.
- 20 Biesenthal, T., Bottenheim, J. W., Shepson, P. B., Li, S. M., and Brickell, P. C.: The chemistry of biogenic hydrocarbons at a rural site in eastern Canada, *J. Geophys. Res.*, 103, 25487–25498, 1998.
- Brown, S. S., Stark, H., and Ravishankara, A. R.: Applicability of the Steady-State Approximation to the Interpretation of Atmospheric Observations of NO₃ and N₂O₅, *J. Geophys. Res.*, 25 108, D174539, doi:10.1029/2003JD003407, 2003.
- Brown, S. S., Dibb, J. E., Stark, H., Aldener, M., Vozella, M., Whitlow, S., Williams, E. J., Lerner, B. M., Jakoubek, R., Middlebrook, A. M., DeGouw, J. A., Warneke, C., Goldan, P. D., Kuster, W. C., Angevine, W. M., Sueper, D. T., Quinn, P. K., Bates, T. S., Meagher, J. F., Fehsenfeld, 30 F. C., and Ravishankara, A. R.: Nighttime removal of NO_x in the summer marine boundary layer, *Geophys. Res. Lett.*, 31, L07108, doi:10.1029/2004GL019412, 2004.
- Brown, S. S., Neuman, J. A., Ryerson, T. B., Trainer, M., Dubé, W. P., Holloway, J. S., Warneke, C., de Gouw, J. A., Donnelly, S. G., Atlas, E., Matthew, B., Middlebrook, A. M., Peltier, R.,

**Nocturnal isoprene
oxidation**

S. S. Brown et al.

Weber, R. J., Stohl, A., Meagher, J. F., Fehsenfeld, F. C., and Ravishankara, A. R.: Nocturnal odd-oxygen budget and its implications for ozone loss in the lower troposphere, *Geophys. Res. Lett.*, 33, L08801, doi:10.1029/2006GL025900, 2006a.

5 Brown, S. S., Ryerson, T. B., Wollny, A. G., Brock, C. A., Peltier, R., Sullivan, A. P., Weber, R. J., Dubé, W. P., Trainer, M., Meagher, J. F., Fehsenfeld, F. C., and Ravishankara, A. R.: Variability in nocturnal nitrogen oxide processing and its role in regional air quality, *Science*, 311, 67–70, 2006b.

10 Brown, S. S., Dubé, W. P., Osthoff, H. D., Stutz, J., Ryerson, T. B., Wollny, A. G., Brock, C. A., Warneke, C., de Gouw, J. A., Atlas, E., Neuman, J. A., Holloway, J. S., Lerner, B. M., Williams, E. J., Kuster, W. C., Goldan, P. D., Angevine, W. M., Trainer, M., Fehsenfeld, F. C., and Ravishankara, A. R.: Vertical profiles in NO_3 and N_2O_5 measured from an aircraft: Results from the NOAA P-3 and surface platforms during NEAQS 2004, *J. Geophys. Res.*, 112, D22304, doi:10.1029/2007JD008883, 2007.

15 Chen, X., Hulbert, D., and Shepson, P. B.: Measurement of the organic nitrate yield from OH reaction with isoprene, *J. Geophys. Res.*, 103, 25563–25568, 1998.

Claeys, M., Graham, B., Vas, G., Wang, W., Vermeylen, R., Pashynska, V., Cafmeyer, J., Guyon, P., Andreae, M. O., Artaxo, P., and Maenhaut, W.: Formation of Secondary Organic Aerosols Through Photooxidation of Isoprene, *Science*, 303, 1173–1176, 2004.

20 de Gouw, J. A., Goldan, P. D., Warneke, C., Kuster, W. C., Roberts, J. M., Marchewka, M., Bertman, S. B., Pszenny, A. A. P., and Keene, W. C.: Validation of proton transfer reaction-mass spectrometry (PTR-MS) measurements of gas-phase organic compounds in the atmosphere during the New England Air Quality Study (NEAQS) in 2002, *J. Geophys. Res.*, 108, D214682, doi:10.1029/2003JD003863, 2003.

25 de Gouw, J. A., Middlebrook, A. M., Warneke, C., Goldan, P. D., Kuster, W. C., Roberts, J. M., Fehsenfeld, F. C., Worsnop, D. R., Canagaratna, M. R., Pszenny, A. A. P., Keene, W. C., Marchewka, M., Bertman, S. B., and Bates, T. S.: Budget of organic carbon in a polluted atmosphere: Results from the New England Air Quality Study in 2002, *J. Geophys. Res.-Atmos.*, 110, D16305, doi:10.1029/2004JD005623, 2005.

30 de Gouw, J. A., Brock, C. A., Atlas, E., Bates, T. S., Fehsenfeld, F. C., Goldan, P. D., Holloway, J. S., Kuster, W. C., Lerner, B. M., Matthew, B. M., Middlebrook, A. M., Onasch, T. B., Peltier, R. E., Quinn, P. K., Senff, C. J., Stohl, A., Sullivan, A. P., Trainer, M., Warneke, C., Weber, R. J., and Williams, E. J.: Sources of particulate matter in the northeastern United States in summer: 1. Direct emission and secondary formation of organic matter in urban plumes, *J.*

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Nocturnal isoprene
oxidation**

S. S. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Geophys. Res., 113, D08301, doi:10.1029/2007JD009243, 2008.
- Dubé, W. P., Brown, S. S., Osthoff, H. D., Nunley, M. R., Ciciora, S. J., Paris, M. W., McLaughlin, R. J., and Ravishankara, A. R.: Aircraft instrument for simultaneous, in-situ measurements of NO_3 and N_2O_5 via cavity ring-down spectroscopy, Rev. Sci. Instrum., 77, 034101, doi:10.1063/1061.2176058, 2006.
- 5 Ervens, B., Carlton, A. G., Turpin, B. J., Altieri, K. E., Kreidenwies, S. M., and Feingold, G.: Secondary organic aerosol yields from cloud-processing of isoprene oxidation products, Geophys. Res. Lett., 35, L02816, doi:10.1029/2007GL031828, 2008.
- Fan, F. and Zhang, R.: Atmospheric Oxidation Mechanism of Isoprene, Environ. Chem., 1, 140–149, 2004.
- 10 Fehsenfeld, F. C., Ancellet, G., Bates, T. S., Goldstein, A. H., Hardesty, R. M., Honrath, R., Law, K., Lewis, A. C., Leatch, R., McKeen, S. A., Meagher, J., Parrish, D. D., Pszenny, A. A. P., Russel, P. B., Schlager, H., Seinfeld, J. H., Talbot, R., and Zbinden, R.: International Consortium for Atmospheric Research on Transport and Transformation (ICARTT): North America to Europe – Overview of the 2004 summer field study, J. Geophys. Res., 111, D23S01, doi:10.1029/2006JD007829, 2006.
- Fischer, R. G., Kaster, J., and Ballschmitter, K.: Levels and pattern of alkyl nitrates, multifunctional alkyl nitrates, and halocarbons in the air over the Atlantic Ocean, J. Geophys. Res., 105, 14473–14494, 2000.
- 20 Fry, J. L., Kiendler-Scharr, A., Rollins, A. W., Wooldridge, P. J., Brown, S. S., Fuchs, H., Dube, W., Mensah, A., dal Maso, M., Tillmann, R., Dorn, H.-P., Brauers, T., and Cohen, R. C.: Organic nitrate and secondary organic aerosol yield from NO_3 oxidation of β -pinene evaluated using a gas-phase kinetics/aerosol partitioning model, Atmos. Chem. Phys. Discuss., 8, 18039–18089, 2008,
- 25 <http://www.atmos-chem-phys-discuss.net/8/18039/2008/>.
- Fuentes, J. D., Lerdau, M., Atkinson, R., Baldocchi, D., Bottenheim, J. W., Ciccioli, P., Lamb, B., Geron, C., Gu, L., Guenther, A., Sharkey, T. D., and Stockwell, W.: Biogenic Hydrocarbons in the Atmospheric Boundary Layer: A Review, Bull. Am. Meteorol. Soc., 81, 1537–1575, 2000.
- 30 Geyer, A., Ackermann, R., Dubois, R., Lohrmann, B., Müller, R., and Platt, U.: Long-term observation of nitrate radicals in the continental boundary layer near Berlin, Atmos. Environ., 35, 3619–3631, 2001.
- Geyer, A. and Platt, U.: Temperature dependence of the NO_3 loss frequency: A new indica-

**Nocturnal isoprene
oxidation**

S. S. Brown et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

tor for the contribution of NO_3 to the oxidation of monoterpenes and NO_x removal in the atmosphere, *J. Geophys. Res.*, 107, D214431, doi: 10.1029/2001JD001215, 2002.

Giacopelli, P., Ford, K., Espada, C., and Shepson, P. B.: Comparison of the measured and simulated isoprene nitrate distributions above a forest canopy, *J. Geophys. Res.*, 110, D01304, doi:10.1029/2004JD005123, 2005.

Grossenbacher, J. W., Couch, T., Shepson, P. B., Thornberry, T., Witmer-Rich, M., Carroll, M. A., Faloon, I., Tan, D., Brune, W., Ostling, K., and Bertman, S.: Measurements of isoprene nitrates above a forest canopy, *J. Geophys. Res.*, 106, 24429–24438, 2001.

Grossenbacher, J. W., Barket, D. J., Shepson, P. B., Carroll, M. A., Olszyna, K., and Apel, E.: A comparison of isoprene nitrate concentrations at two forest-impacted sites, *J. Geophys. Res.*, 109, D11311, doi:10.1029/2003JD003966, 2004.

Guenther, A., Karl, T., Harley, P., Wiedinmyer, C., Palmer, P. I., and Geron, C.: Estimates of global terrestrial isoprene emissions using MEGAN (Model of Emissions of Gases and Aerosols from Nature), *Atmos. Chem. Phys.*, 6, 3181–3210, 2006, <http://www.atmos-chem-phys.net/6/3181/2006/>.

Hallquist, M., Wängberg, I., Ljungstrom, E., Barnes, I., and Becker, K. H.: Aerosol and Product Yields from NO_3 Radical-Initiated Oxidation of Selected Monoterpenes, *Environ. Sci. Technol.*, 33, 553–559, 1999.

Henze, D. K. and Seinfeld, J. H.: Global secondary organic aerosol from isoprene oxidation, *Geophys. Res. Lett.*, 33, L09812, doi:10.1029/2006GL025976, 2006.

Horowitz, L. W., Fiore, A. M., Milly, G. P., Cohen, R. C., Perrin, A., Wooldridge, P. J., Hess, P. G., Emmons, L. K., and Lamarque, J. F.: Observational constraints on the chemistry of isoprene nitrates over the eastern United States, *J. Geophys. Res.*, 112, D12S08, doi:10.1029/2006JD007747, 2007.

Hurst, J. M., Barket, D. J., Herrera-Gomez, O., Couch, T. L., Shepson, P. B., Faloon, I., Tan, D., Brune, W., Westberg, H., Lamb, B., Biesenthal, T., Young, V., Goldstein, A., Munger, J. W., Thornberry, T., and Carroll, M. A.: Investigation of the nighttime decay of isoprene, *J. Geophys. Res.-Atmos.*, 106, 24335–24346, 2001.

Kim, S.-W., Heckel, A., McKeen, S. A., Frost, G. J., Hsie, E. Y., Trainer, M. K., Richter, A., Burrows, J. P., Peckham, S. E., and Grell, G. A.: Satellite-observed US power plant NO_x emission reductions and their impact on air quality, *Geophys. Res. Lett.*, 33, L22812, doi:10.1029/2006GL027749, 2006.

Kleindienst, T. E., Lewandowski, M., Offenberg, J. H., Jaoui, M., and Edney, E. O.: Ozone-

**Nocturnal isoprene
oxidation**

S. S. Brown et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

isoprene reaction: Re-examination of the formation of secondary organic aerosol, *Geophys. Res. Lett.*, 34, L01805, doi:10.1029/2006GL027485, 2007.

Kroll, J. H., Ng, N. L., Murphy, S. M., Flagan, R. C., and Seinfeld, J. H.: Secondary organic aerosol formation from isoprene photooxidation under high-NO_x conditions, *Geophys. Res. Lett.*, 32, L18808, doi:10.1029/2005GL023637, 2005.

Kwok, E. S. C., Aschmann, S. M., Arey, J., and Atkinson, R.: Product formation from the reaction of the NO₃ radical with isoprene and rate constants for the reactions of methacrolein and methyl vinyl ketone with the NO₃ radical, *Int. J. Chem. Kinet.*, 28, 925–934, 1996.

Lelieveld, J., Butler, T. M., Crowley, J. N., Dillon, T. J., Fischer, H., Ganzveld, L., Harder, H., Lawrence, M. G., Martinez, M., Taraborrelli, D., and Williams, J.: Atmospheric oxidation capacity sustained by a tropical forest, *Nature*, 452, 737–740, 2008.

Liggio, J., Li, S.-M., Brook, J. R., and Mihele, C. M.: Direct polymerization of isoprene and α -pinene on acidic aerosols, *Geophys. Res. Lett.*, 34, L05814, doi:10.1029/2006GL028468, 2007.

McLaren, R., Salmon, R. A., Liggio, J., Hayden, K. L., Analuf, K., and Leaitch, W. R.: Nighttime chemistry at a rural site in the Lower Fraser Valley, *Atmos. Environ.*, 38, 5837–5848, 2004.

Neuman, J. A., Huey, L. G., Dissly, R. W., Fehsenfeld, F. C., Flocke, F., Holecek, J. C., Holloway, J. S., Hübler, G., Jakoubek, R., Nicks, D. K., Jr., Parrish, D. D., Ryerson, T. B., Sueper, D. T., and Weinheimer, A. J.: Fast-response airborne in situ measurements of HNO₃ during the Texas 2000 Air Quality Study, *J. Geophys. Res.*, 107, 4436, doi:10.1029/2001JD001437, 2002.

Neuman, J. A.: Reactive nitrogen transport and photochemistry in urban plumes over the North Atlantic Ocean, *J. Geophys. Res.*, 111, D23S54, doi:10.1029/2005JD007010, 2006.

Ng, N. L., Kwan, A. J., Surratt, J. D., Chan, A. W. H., Chhabra, P. S., Sorooshian, A., Pye, H. O. T., Crouse, J. D., Wennberg, P. O., Flagan, R. C., and Seinfeld, J. H.: Secondary organic aerosol (SOA) formation from reaction of isoprene with nitrate radicals (NO₃), *Atmos. Chem. Phys.*, 8, 4117–4140, 2008, <http://www.atmos-chem-phys.net/8/4117/2008/>.

Olivier, J. G. J. and Berdowski, J. J. M.: Global emissions sources and sinks, in: *The Climate System*, edited by: Berdowski, J., Guicherit, R., and Heij, B. J., A. A. Balkema, The Netherlands, 33–78, 2001.

Paulson, S. E. and Seinfeld, J. H.: Development and Evaluation of a Photooxidation Mechanism for Isoprene, *J. Geophys. Res.*, 97, 20703–20715, 1992.

**Nocturnal isoprene
oxidation**

S. S. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Platt, U. F., Winer, A. M., Bierman, H. W., Atkinson, R., and Pitts Jr., J. N.: Measurement of Nitrate Radical Concentrations in Continental Air, *Environ. Sci. Technol.*, 18, 365–369, 1984.
- Poisson, N., Kanakidou, M., and Crutzen, P. J.: Impact of Non-Methane Hydrocarbons on Tropospheric Chemistry and the Oxidizing Power of the Global Troposphere: 3-Dimensional Modelling Results, *J. Atmos. Chem.*, 36, 157–230, 2000.
- Quinn, P. K., Bates, T. S., Coffman, D., Onasch, T. B., Worsnop, D. R., Baynard, T., de Gouw, J. A., Goldan, P. D., Kuster, W. C., Williams, E., Roberts, J. M., Lerner, B. M., Stohl, A., Pettersson, A., and Lovejoy, E. R.: Impacts of sources and aging on submicrometer aerosol properties in the marine boundary layer across the Gulf of Maine, *J. Geophys. Res.*, 111, D23S36, doi:10.1029/2006JD007582, 2006.
- Ryerson, T. B., Huey, L. G., Knapp, K., Neuman, J. A., Parrish, D. D., Sueper, D. T., and Fehsenfeld, F. C.: Design and initial characterization of an inlet for gas-phase NO_y measurements from aircraft, *J. Geophys. Res.*, 104, 5483–5492, 1999.
- Ryerson, T. B., Williams, E. J., and Fehsenfeld, F. C.: An efficient photolysis system for fast response NO_2 measurements, *J. Geophys. Res.*, 105, 26447–26461, 2000.
- Schauffler, S. M., Atlas, E., Blake, D. R., Flocke, F., Lueb, R. A., Lee-Taylor, J. M., Stroud, V., and Travnicek, W.: Distributions of brominated organic compounds in the troposphere and lower stratosphere, *J. Geophys. Res.*, 104, 21513–21536, 1999.
- Schichtel, B., Malm, W. C., Bench, G., Fallon, S., McDade, C. E., Chow, J. C., and Watson, J. G.: Fossil and contemporary fine particulate carbon fractions at 12 rural and urban sites in the United States, *J. Geophys. Res.*, 113, D02311, doi:10.1029/2007JD008605, 2008.
- Sillman, S., Carroll, M. A., Thornberry, T., Lamb, B. K., Westberg, H., Brune, W. H., Faloon, I., Tan, D., Shepson, P. B., Sumner, A. L., Hastie, D. R., Mihele, C. M., Apel, E. C., Riemer, D. D., and Zika, R. G.: Loss of isoprene and sources of nighttime OH radicals at a rural site in the United States: Results from photochemical models, *J. Geophys. Res.*, 107, 4043, doi:10.1029/2001JD000449, 2002.
- Skamarock, W. C., Klemp, J. B., Dudhia, J., Gill, D. O., Barker, D. M., Duda, M. G., Huang, X. Y., Wang, W., and Powers, J. G.: A description of the advanced research WRF version 3, National Center for Atmospheric Research Technical Note, NCAR/TN-475+STR, online: http://www.mmm.ucar.edu/wrf/users/docs/arw_v3.pdf, 2008.
- Skov, H., Hjorth, J., Lohse, C., Jensen, N. R., and Restelli, G.: Products and mechanisms of the reactions of the nitrate radical (NO_3) with isoprene, 1,3-butadiene and 2,3-dimethyl-1,3-butadiene in air, *Atmos. Environ.*, 26A, 2771–2783, 1992.

**Nocturnal isoprene
oxidation**

S. S. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Skov, H., Benter, T., Schindler, R. N., Hjorth, J., and Restelli, G.: Epoxide Formation in the Reactions of the Nitrate Radical with 2,3-Dimethyl-2-Butene, Cis-2-Butene and Trans-2-Butene and Isoprene, *Atmos. Environ.*, 28, 1583–1592, 1994.

Slusher, D. L., Huey, L. G., Tanner, D. J., Flocke, F., and Roberts, J. M.: A thermal dissociation – chemical ionization mass spectrometry (TD-CIMS) technique for the simultaneous measurement of peroxyacetyl radicals and dinitrogen pentoxide, *J. Geophys. Res.*, 109, D19315, doi:10.1029/2004JD004670, 2004.

Sommariva, R., Osthoff, H. D., Brown, S. S., Bates, T. S., Baynard, T., Coffman, D., de Gouw, J. A., Goldan, P. D., Kuster, W. C., Lerner, B. M., Stark, H., Warneke, C., Williams, E. J., Fehsenfeld, F. C., Ravishankara, A. R., and Trainer, M.: Radicals in the marine boundary layer during NEAQS 2004: a model study of day-time and night-time sources and sinks, *Atmos. Chem. Phys. Discuss.*, 8, 16643–16692, 2008, <http://www.atmos-chem-phys-discuss.net/8/16643/2008/>.

Sprengnether, M. M., Demerjian, K. L., Donahue, N. M., and Anderson, J. G.: Product analysis of the OH oxidation of isoprene and 1,3-butadiene in the presence of NO, *J. Geophys. Res.*, 107, 4268, doi:10.1029/2001JD000716, 2002.

Starn, T. K., Shepson, P. B., Bertman, S. B., Reimer, D. D., Zika, R. G., and Olszyna, K.: Night-time isoprene chemistry at an urban-impacted forest site, *J. Geophys. Res.*, 103, 22437–22447, 1998.

Steinbacher, M., Dommen, J., Ordonez, C., Reimann, S., Gruebler, F. C., Staehelin, J., Andreani-Aksoyoglu, S., and Prevot, A. S. H.: Volatile Organic Compounds in the Po Basin. Part B: Biogenic VOCs, *J. Atmos. Chem.*, 51, 293–315, doi:10.1007/s10874-005-3577-0, 2005.

Stohl, A., Forster, C., Frank, A., Seibert, P., and Wotawa, G.: Technical note: The Lagrangian particle dispersion model FLEXPART version 6.2, *Atmos. Chem. Phys.*, 5, 2461–2474, 2005, <http://www.atmos-chem-phys.net/5/2461/2005/>.

Stroud, C. A., Roberts, J. M., Williams, E. J., Hereid, D., Angevine, W. M., Fehsenfeld, F. C., Wisthaler, A., Hansel, A., Martinez-Harder, M., Harder, H., Brune, W. H., Hoenninger, G., Stutz, J., and White, A. B.: Nighttime isoprene trends at an urban forested site during the 1999 Southern Oxidants Study, *J. Geophys. Res.*, 107, 4291, doi:10.1029/2001JD000959, 2002.

Stutz, J., Alicke, B., Ackermann, R., Geyer, A., White, A. B., and Williams, E.: Vertical profiles of NO₃, N₂O₂, O₃, and NO_x in the nocturnal boundary layer: 1. Observations during the Texas

**Nocturnal isoprene
oxidation**

S. S. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Air Quality Study 2000, *J. Geophys. Res.*, 109, D12306, doi:10.1029/2003JD004209, 2004.
- Sullivan, A. P., Peltier, R. E., Brock, C. A., de Gouw, J. A., Holloway, J. S., Warneke, C., Wollny, A. G., and Weber, R. J.: Airborne measurements of carbonaceous aerosol soluble in water over the northeastern United States: Method development and an investigation into water-soluble organic carbon sources, *J. Geophys. Res.*, 111, D23S46, doi:10.1029/2006JD007072, 2006.
- Trainer, M., Williams, E. J., Parrish, D. D., Buhr, M. P., Allwine, E. J., Westberg, H. H., Fehsenfeld, F. C., and Liu, S. C.: Models and observations of the impact of natural hydrocarbons on rural ozone, *Nature*, 329, 705–707, 1987.
- Trainer, M., Buhr, M. P., Curran, C. M., Fehsenfeld, F. C., Hsie, E. Y., Liu, S. C., Norton, R. B., Parrish, D. D., Williams, E. J., Gandrud, B. W., Ridley, B. A., Shetter, J. D., Allwine, E. J., and Westberg, H. H.: Observations and Modeling of the Reactive Nitrogen Photochemistry at a Rural Site, *J. Geophys. Res.*, 96, 3045–3063, 1991.
- Tuazon, E. C., Alvarado, A., Aschmann, S. M., Atkinson, R., and Arey, J.: Products of the gas-phase reactions of 1,3-butadiene with OH and NO₃ radicals, *Environ. Sci. Technol.*, 33, 3586–3595, 1999.
- von Kuhlmann, R., Lawrence, M. G., Pöschl, U., and Crutzen, P. J.: Sensitivities in global scale modeling of isoprene, *Atmos. Chem. Phys.*, 4, 1–17, 2004, <http://www.atmos-chem-phys.net/4/1/2004/>.
- Warneke, C., de Gouw, J. A., Goldan, P. D., Kuster, W. C., Williams, E. J., Lerner, B. M., Brown, S. S., Stark, H., Aldener, M., Ravishankara, A. R., Roberts, J. M., Marchewka, M., Bertman, S., Sueper, D. T., McKeen, S. A., Meagher, J. F., and Fehsenfeld, F. C.: Comparison of day and nighttime oxidation of biogenic and anthropogenic VOCs along the New England coast in summer during New England Air Quality Study 2002, *J. Geophys. Res.*, 109, D10309, doi:10.1029/2003JD004424, 2004.
- Weber, R. J., Orsini, D., Daun, Y., Lee, Y.-N., Klotz, P. J., and Brechtel, F.: A particle-into-liquid collector for rapid measurement of aerosol bulk chemical composition, *Aerosol Sci. Technol.*, 35, 718–727, 2001.
- Weber, R. J., Sullivan, A. P., Peltier, R., Russell, A., Yan, B., Zheng, M., de Gouw, J. A., Warneke, C., Brock, C. A., Holloway, J. S., Atlas, E. L., and Edgerton, E.: A study of secondary organic aerosol formation in the anthropogenic-influenced southeastern United States, *J. Geophys. Res.*, 112, D13302, doi:10.1029/2007JD008408, 2007.
- Werner, G., Kaster, J., Looser, R., and Ballschmitter, K.: Organic Nitrates of Isoprene as Atmo-

spheric Trace Compounds, *Angew. Chem. Int. Edit.*, 38, 1634–1637, 1999.
Winer, A. M., Atkinson, R., and Pitts, J. N. J.: Gaseous Nitrate Radical: Possible Nighttime
Atmospheric Sink for Biogenic Organic Compounds, *Science*, 224, 156–158, 1984.

ACPD

9, 225–269, 2009

Nocturnal isoprene oxidation

S. S. Brown et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Nocturnal isoprene oxidation

S. S. Brown et al.

Table 1. Instruments used for analysis of nocturnal isoprene oxidation.

Measurement/Technique	Accuracy	Frequency	Reference
NO ₃ , N ₂ O ₅ /CRDS ^a	25%	1 Hz	(Dubé et al., 2006)
NO, NO ₂ , O ₃ /Chemiluminescence	3–8%	1 Hz	(Ryerson et al., 1999, 2000)
Rapid VOC/PTRMS ^b	10–20%	0.08 Hz	(de Gouw et al., 2003)
Speciated VOC/Can Samples ^c	5–10%	80/flight	(Schauffler et al., 1999)
HNO ₃ /CIMS ^d	15%	1 Hz	(Neuman et al., 2002)
Speciated PANs/CIMS ^d	20%	0.5 Hz	(Slusher et al., 2004)
Aerosol composition/PiLS ^e	15%	0.4 min ⁻¹	(Weber et al., 2001)
WSOC/PiLS ^e	15%	1 min ⁻¹	(Weber et al., 2001)

^a Cavity ring down spectroscopy,

^b Proton transfer reaction mass spectrometry,

^c Analyzed by GC-MS,

^d Chemical ionization mass spectrometry,

^e Particle into liquid sampler.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Nocturnal isoprene oxidation

S. S. Brown et al.

Table 2. Night flights during NEAQS 2004 and ratio of sunset isoprene, determined from Eq. (7), to observed MVK+MACR.

Date	Location	Sunset Isop/MVK+MACR	Range	Correlation Coef. (<i>r</i>)
11 July	Boston	0.4	0.2–0.7	0.65
31 Jul–1 Aug	CT, RI, MA	0.6	0.2–1.2	0.28
2–3 Aug	NH, MA, RI, NYC	0.6	0.3–0.9	0.72
7–8 Aug	Coastal Areas Offshore flow	0.8	0.2–1.2	0.57
9–10 Aug	Northeast US-Ohio	0.6	0.2–1.2	0.54
10–11 Aug	Maine, Atlantic Ocean	N/A ^a		

^a No determination from this flight.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[I◀](#)
[▶I](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Nocturnal isoprene oxidation

S. S. Brown et al.

Table 3. Reactive nitrogen partitioning and contribution of isoprene oxidation products to SOA.

Date	Isop Nitrate/ NO _y (%)	Isop Nitrate/ Isop Nitrate+PAN (%)	Isop SOA/ WSOC (%)	NO ₃ -Isop SOA/ Total Isop SOA (%) ^b
11 Jul	3±2	7±4	<2	31
31 Jul	6±5	26±22	0.5–17	56
2–3 Aug	4±2	16±8	1–12	60
7–8 Aug	8±4	31±15	1–15	86
9–10 Aug ^a	6±4	26±17	0.5–6	57
Average	6±4%	21±18%	1–11%	58±17%

^a Data for Northeast US portion of this flight only,

^b Total isoprene SOA considers that resulting from gas-phase OH, O₃ and NO₃ oxidation but not SOA formed from, for example, heterogeneous reactions of isoprene.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


**Nocturnal isoprene
oxidation**

S. S. Brown et al.




Fig. 1. Map of the Northeast US showing the MEGAN (Guenther et al., 2006) database for isoprene emissions and the nighttime P-3 flight tracks during the NEAQS 2004 campaign. The dashed box indicates the domain for the model calculation described in Sect. 4.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Nocturnal isoprene
oxidation

S. S. Brown et al.




Fig. 2. Eastern part of the 9–10 August flight after sunset color and size code by NO_3 (a) and PTRMS isoprene (b) mixing ratios. The altitude range for the majority of the flight track shown was 470–950 m. Panel (c) shows the correlation plot between measured isoprene and NO_3 . Red points are PTRMS isoprene (high time resolution), and blue points are canister samples (low time resolution, higher precision). In some cases, there is a small background at the mass used for isoprene detection in the PTRMS ($m/z=69$), evident on the comparison to the canister samples at high NO_3 .

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Nocturnal isoprene
oxidation

S. S. Brown et al.




Fig. 3. Time series of isoprene and NO_3 (top traces) and first-order loss rate coefficient for NO_3 radicals (bottom traces) for the eastern part of the 9–10 August P-3 flight shown in Fig. 2. $\tau(\text{NO}_3)^{-1}$ is the inverse steady state lifetime, or loss frequency, calculated according to Eq. (4), and $k_1(\text{Isop})$ is the first-order loss rate coefficient for NO_3 with measured isoprene from the PTRMS instrument according to Eq. (6).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Nocturnal isoprene
oxidation**

S. S. Brown et al.




Fig. 4. Plot of first-order loss rate coefficient for NO_3 with isoprene against inverse NO_3 lifetime for the data in Fig. 3, averaged to 30 s and for inverse NO_3 lifetimes $>0.05 \text{ min}^{-1}$ (≈ 50 pptv isoprene), above which isoprene loss is the controlling factor.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Nocturnal isoprene
oxidation

S. S. Brown et al.




Fig. 5. Tracks for the 3 August 2004 night flight color and size coded by NO_3 mixing ratio (upper left), isoprene (lower left) and methyl vinyl ketone+methacrolein (MVK+MACR) (lower right). The abrupt end to the flight track over Connecticut was due to a technical difficulty that forced the suspension of data collection at that point. The upper right graph shows the correlation plot between isoprene and NO_3 as in Fig. 2.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Nocturnal isoprene
oxidation

S. S. Brown et al.




Fig. 6. Upper panel: Time series of isoprene, methyl vinyl ketone+methacrolein (MVK+MACR) and isoprene present at sunset calculated according to Eq. (7) for isoprene >100 pptv. Lower panel: Plot of initial isoprene vs. MVK+MACR with a linear fit with a slope of 0.6 ± 0.3 and a correlation coefficient of 0.72. This correlation can be used to estimate sunset isoprene from the MVK+MACR data for the entire flight.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Nocturnal isoprene
oxidation

S. S. Brown et al.




Fig. 7. (A) Sample FLEXPART footprint for one point on the P-3 flight track (asterisk). The plot shows residence time (per unit mass of a uniformly distributed, arbitrary tracer emission) of the air mass in the boundary layer on a logarithmic color scale. (B) Time series of the determined isoprene oxidized in the dark and the integrated isoprene emissions from the preceding day from the FLEXPART model for the 3 August flight. The trace over the top shows the ratio of the determination of isoprene oxidized after dark to integrated FLEXPART isoprene emissions.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Nocturnal isoprene
oxidation

S. S. Brown et al.




Fig. 8. Top: time series of reactive nitrogen compounds for the 2–3 August flight according to the color scale on the legend. The large plume near midnight local time was from New York City, and the smaller plumes that preceded it were from the isoprene rich areas over New Hampshire, Massachusetts and Rhode Island. Center: time series of reactive nitrogen partitioning according to the same color code. Bottom: pie charts for average partitioning for the isoprene rich areas early in the flight (left) and for the entire flight (right). Uncertainties for fractions of reactive nitrogen due to isoprene nitrates are $\pm 50\%$.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Nocturnal isoprene
nitrate oxidation

S. S. Brown et al.




Fig. 9. Plot of calculated isoprene nitrates from NO_3 +isoprene against the calculated NO_2 that had been oxidized after sunset, assuming that reactions of NO_3 were dominant compared to N_2O_5 hydrolysis, for the first half of the 3 August flight.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Nocturnal isoprene
oxidation

S. S. Brown et al.




Fig. 10. Top: time series from the 3 August flight of water soluble organic carbon (WSOC), nocturnally oxidized isoprene (in $\mu\text{g C m}^{-3}$, same axis) and secondary organic aerosol (SOA) derived from that isoprene using yields from Ng et al. (2008). The uncertainty in the shaded region is due to the range of aerosol yields (2.5–12.5% carbon mass, see text) and does not reflect the $\pm 50\%$ uncertainty in the derived nighttime isoprene oxidation. The full uncertainty is given in Table 3. Bottom: plots of nocturnally oxidized isoprene (left) and isoprene-SOA against WSOC from the data in the top graph. Blue squares and red circles on the bottom right graph show the correlation of isoprene SOA against WSOC for 2.5 and 12.5% yields, giving a 2 and 8% contribution to observed WSOC, respectively.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Nocturnal isoprene
oxidation

S. S. Brown et al.




Fig. 11. Calculated isoprene SOA mass for the 3 August flight due to NO_3 , OH and O_3 . Sticks with error bars give average mass and uncertainty for different sampled air masses, and the pie chart gives the relative integrated contribution of each oxidant for the entire flight as percentages. Error bars reflect only the uncertainty in mass yields and not the uncertainties due to the FLEXPART isoprene calculation or the determination of the nighttime oxidized isoprene mass (see text).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Nocturnal isoprene
oxidation

S. S. Brown et al.




Fig. 12. Same as Fig. 10, except for the 7 August flight. The maps in the top panels shows the after-sunset portion of the flight track, color and size coded by NO_3 (left) and isoprene (right).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)