

This discussion paper is/has been under review for the journal *Atmospheric Chemistry and Physics (ACP)*. Please refer to the corresponding final paper in *ACP* if available.

**Halogen chemistry
over the Dead Sea**

L. Smoydzin
and R. von Glasow

Modelling chemistry over the Dead Sea: bromine and ozone chemistry

L. Smoydzin^{1,*} and R. von Glasow¹

¹School of Environmental Sciences, University of East Anglia, Norwich, UK

*now at: Max-Planck-Institute for Chemistry, Department of Atmospheric Chemistry,
P.O. Box 3060, 55020 Mainz, Germany

Received: 22 December 2008 – Accepted: 2 February 2009 – Published: 23 February 2009

Correspondence to: L. Smoydzin (smoydzin@mpch-mainz.mpg.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Measurements of O₃ and BrO concentrations over the Dead Sea indicate that Ozone Depletion Events (ODEs), widely known to happen in polar regions, are also likely to occur over the Dead Sea due to the very high bromine content of the Dead Sea water. However, we show that BrO and O₃ levels as they are detected cannot solely be explained by high Br⁻ levels in the Dead Sea water and the release of gas phase halogen species out of sea borne aerosol particles and their conversion to reactive halogen species. It is likely that other sources for reactive halogen compounds are needed to explain the observed concentrations for BrO and O₃. To explain the chemical mechanism taking place over the Dead Sea leading to BrO levels of several pmol/mol we used the single column model MISTRA which calculates microphysics, meteorology, gas and aerosol phase chemistry. We performed pseudo Lagrangian studies by letting the model column first move over the desert which surrounds the Dead Sea region and then let it move over the Dead Sea itself. To include an additional source for gas phase halogen compounds, gas exchange between the Dead Sea water and the atmosphere is treated explicitly. Model calculations indicate that this process has to be included to explain the measurements.

1 Introduction

Tropospheric ozone depletion events related to highly active bromine chemistry are widely known to take place in polar regions and were first observed by Barrie et al. (1988). Several years ago Hebestreit et al. (1999) detected a similar occurrence at the Dead Sea, Israel. Hebestreit et al. (1999) report maximum BrO mixing ratios of 86 pmol/mol, exceeding even mixing ratios in the Arctic regions. O₃ levels dropped simultaneously below the detection limit of 2 nmol/mol. Those first measurements were taken on a peninsula separating the northern and southern part of the Dead Sea. On all measurement days with southerly winds Hebestreit et al. (1999)

Halogen chemistry over the Dead Sea

L. Smoydzin
and R. von Glasow

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Halogen chemistry
over the Dead Sea**L. Smoydzin
and R. von Glasow

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



could observe a strong increase in BrO mixing ratios until noon. However, no BrO was found when the wind direction was northerly. Matveev et al. (2001) made BrO and O₃ measurements at the same site as Hebestreit et al. (1999) and obtained similar results regarding the high BrO mixing ratios but with a maximum of 176 pmol/mol and correlated O₃ depletion events when surface winds were southerly. However, in contrast to the first mentioned measurement campaign they could also detect high BrO levels when the wind direction was from north to south, even though under those conditions no O₃ depletion could be observed. During two long term measurement campaigns, O₃ was monitored for more than one year at three sites (south, mid-west, north; additionally at a site 400 m above sea level on several days). Tas et al. (2003) and Tas et al. (2005) report high levels of BrO and low levels of O₃ on numerous days at all measurement sites, in summer time as well as in winter time. However, aircraft measurements (Matveev et al., 2001) show how strongly variable the horizontal extent of the ODEs are.

A first very simple boxmodel study by Stutz et al. (1999) indicated that the formation of 0.5 nmol/mol BrO, which is about 3 times the absolute maximum measured BrO mixing ratio, would be required to destroy an average background level of about 60 nmol/mol O₃ within 2–3 h. A recent model study by Tas et al. (2006) claims that the only source for the high bromine levels is the bromine explosion reaction taking place within sea salt aerosols which are emitted from the Dead Sea water and the related release of bromine species out of these aerosol particles. However, as Tas et al. (2006) have chosen a set up constraining their model to a large extent and as several of their assumptions are strongly simplified the conclusions drawn from this study have to be thoroughly reconsidered.

Besides the Dead Sea two other salt lakes have turned out to be regions of highly active halogen chemistry in the midlatitudes. Stutz et al. (2002) could detect BrO mixing ratios of up to 6 pmol/mol at the Great Salt Lake, USA. The authors find a decrease in O₃ mixing ratios simultaneously to the daily BrO peak period, however, the decrease is far lower than for the Dead Sea. Furthermore, Hönninger et al. (2004) found up to

20 pmol/mol BrO correlated with O₃ depletion events at the Salar de Uyuni in Bolivia.

2 Motivation

The main goal of this paper is to find an explanation for the measurements taken along the Dead Sea based on a reasonable chemistry mechanism.

In general the most important source for gas phase halogen species in the marine troposphere are sea salt aerosols. As soon as H⁺ concentrations are large enough, BrCl and Br₂ are formed in the aerosol and degas into the gas phase (Vogt et al., 1996; von Glasow and Crutzen, 2007). Gas phase BrCl and Br₂ are rapidly photolysed during the day and form Br which can react with O₃ and produce BrO. Bromine oxide can react further leading to the production of HOBr. Hypobromous acid is taken up by the aerosol where it takes part in the acid catalysed reactions forming BrCl and Br₂. In addition, BrO can react with NO₂ and produce BrONO₂ which participates in heterogeneous reactions on aerosol surfaces also starting the recycling of gas phase species in the aqueous phase (see Fig. 1). It has been suggested several times that due to the very high bromide content and low pH of the Dead Sea water and thus of emitted sea salt aerosol particles the above explained process also leads to extraordinary high BrO mixing ratios and the related ODEs. Two, very general points speak against this hypothesis (i) The wind speed over the Dead Sea is very low. Consequently the number of sea salt aerosol particles that are produced is low. (ii) As the relative humidity along the Dead Sea valley is very low, aerosol phase chemistry is likely to be of minor importance. A low aerosol liquid water content which consequently has to be the case if ambient relative humidity is extremely low, leads to a slow mass transfer between the aerosol and the gas phase. Thus, only a very limited amount of BrCl or Br₂ can be released out of particles as the total halogen cycling should be weak. Due to the dry conditions it has further to be taken into account that sea salt aerosol particles dry out quickly when they are transported vertically as the ambient relative humidity can be as low as 40% at an altitude of only 10 m above the water surface. As such conditions are

Halogen chemistry over the Dead Sea

L. Smoydzin
and R. von Glasow

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Halogen chemistry
over the Dead Sea**L. Smoydzin
and R. von Glasow

common along the Dead Sea, sea salt aerosols can be considered as sources for gas phase bromine species only in the lowest metres above the water. As the production of BrCl and Br₂ in the aerosol is pH dependant and the exchange process between the aerosol and the gas phase depends on the solubility of the species as well as their gas phase and aqueous phase concentrations, BrCl can only degas directly out of acidified aerosols but not out of open ocean water with an average pH of about 8. However, the pH of Dead Sea water is much lower, leading to larger BrCl and Br₂ production rates in the water. This indicates that beside sea salt aerosols the Dead Sea water itself can be a direct source for gas phase bromine species. In order to investigate this aspect we included the air-sea exchange of bromine species explicitly in a numerical model.

3 Background information about the Dead Sea

The Dead Sea extends 75 km from north to south and 15 km from East to West. On the East as well as the West shore it is flanked by mountains exceeding 1000 m altitude. The southern part of the Dead Sea, the so-called evaporation ponds, are separated from the northern part by a peninsula. These very shallow basins are fed with water from the northern part of the lake by a channel. The water of the Dead Sea has a very high salinity and acidity compared to ocean water with pH values in the order of 5 (southern part) to 6 (northern part, Tas et al., 2005). Bromide as well as chloride concentrations increase significantly from the northern to the southern end of the lake (Cl⁻: 225–405 g/l, Br⁻: 6.1–11 g/l, Niemi et al., 1997). Thus the salinity of the water is about 12 times larger than for ocean water which has an average chloride content of about 19 g/l and a bromide content of 0.065 g/l (Sverdrup et al., 1942).

The topography at the Dead Sea leads to a strongly canalysed wind flow through the valley such that the main wind direction is roughly south to north. The local flow pattern, which is still insufficiently understood is partly influenced by the lake breeze leading to a wind from the centre of the lake to the north/south shore (Clarke, 1970; Tas et al., 2003). Thus the wind direction at the north end of the lake is almost always

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

southerly. The wind direction at the south end seems to be more variable and southerly as well as northerly winds are observed (Tas et al., 2003). The wind speed during day time is very low and is in the range of 2–4 m/s.

As the Dead Sea is surrounded by a desert the temperature is very high with mean values of about 35°C in summer and the relative humidity is very low with about 40–50% only a few metres above the sea surface (Alpert et al., 1997, and references therein, personal communication M. Piot, Inst. of Env. Physics, Univ. Heidelberg, Germany)

4 Model description

For all presented model studies the one dimensional numerical model MISTRA is used. It contains a microphysics scheme and explicitly calculates chemistry in the gas phase, in aerosol particles and cloud droplets (von Glasow et al., 2002a,b). The chemical mechanism contains the most important reactions of O, H, C, N, S, Cl and Br both in the gas phase and the aerosol phase and has been updated according to the IUPAC recommendation (04-2004, see <http://www.iupac-kinetic.ch.cam.ac.uk>). In the model we distinguish between sulphate and sea salt aerosol particles which differ by their size ($r=0.005\text{--}0.5\ \mu\text{m}$ for sulphate, $r=0.5\text{--}15\ \mu\text{m}$ for sea salt) and initial chemical composition. The parameterisation of Monahan et al. (1986) for the emission of sea salt particles by bubble bursting is used. An aerosol size distribution for rural areas based on Jaenicke (1988) is used. All presented model runs are set up such that no clouds are formed. To be able to calculate air-sea exchange between the Dead Sea and the overlying atmosphere, chemistry within the water of the lake is calculated explicitly in addition to the aerosol phase. We assume that the same reaction set which is used for aerosol phase chemistry is also valid for chemistry taking place in the water of the Dead Sea apart from heterogeneous reactions. The only possible interactions that are allowed between the Dead Sea water and the atmosphere is via air-sea exchange of trace gases as is shown exemplarily in Fig. 1. To describe the fluxes of gases across

Halogen chemistry over the Dead Sea

L. Smoydzin
and R. von Glasow

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the air-sea interface a two-layer model as first presented by Liss and Slater (1974) is used. The implementation of the air-sea exchange mechanism into MISTRA follows the work done by Pozzer et al. (2006) (and references therein).

For studying the different formation mechanisms of BrO in detail it is crucial to have a high vertical model resolution. Thus a logarithmically spaced grid for the vertical has been chosen with a grid spacing of 0.5 m for the lowest grid box above the water surface.

4.1 Activation of aerosol-chemistry

Due to the extremely dry conditions along the Dead Sea it is crucial to discuss the way aerosol chemistry is triggered in MISTRA. Aerosol phase chemistry is always calculated if the ambient relative humidity is above the deliquescence humidity of sea salt (75%) or sulphate (70%) aerosols. When humidity decreases to the crystallisation humidity (40% for sulphate, 42% for sea salt) calculation of aerosol phase chemistry is terminated. As it can be assumed that freshly emitted sea salt particles are always present as an aqueous solution, sea salt chemistry is always calculated in the lowest model level (over a water surface). As it is well documented that sea salt aerosol particles remain in a metastable state when the relative humidity drops below the deliquescence humidity, aerosol chemistry is calculated in all model layers where the relative humidity had previously exceeded the deliquescence humidity but is still higher than the crystallisation humidity. Calculation of aqueous phase chemistry is however, terminated if the total water mass in one aerosol size class drops below a threshold value of $10^{-13} \text{ m}_{(aq)}^3 / \text{m}_{(air)}^3$. To re-activate aerosol phase chemistry, the ambient relative humidity has to exceed the deliquescence humidity of the particles.

Halogen chemistry over the Dead Sea

L. Smoydzin
and R. von Glasow

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



5 Model setup

5.1 Lagrangian

To investigate bromine chemistry over the Dead Sea and related ODEs a Lagrangian-kind model setup was chosen which can be understood as follows: The model column first moves over the desert for two days and then crosses the Dead Sea from south to north. The temperature ranges from about 42°C directly above the ground to about 22°C at the top of the boundary layer during day. During the nights temperature decreases below 20°C close to the surface. Relative humidity increases from 40% in the lowest metres above the ground to a value of about 80% at the top of the boundary layer at about 1500 m altitude during day. During the night surface air masses cool down. Due to the decrease in temperature, relative humidity increases up to about 70%. Chemistry is initialised such that average background mixing ratios for O₃ are about 70 nmol/mol and 1 nmol/mol for NO₂ at the morning of day two in the initialisation model run. Figure 2 shows typical vertical profiles of temperature, potential temperature and relative humidity over the Dead Sea for two scenarios which will be introduced below. The temperature inversion indicates the boundary layer height in the model.

To initialise chemistry in the Dead Sea water, a separate model run was calculated for two days prohibiting air-sea exchange in order to have a spin up. It is assumed that the model column crosses the lake from south to north. With this setup the influence of the gradient in chemical composition of the lake's water on gas phase concentrations can be taken into account. By using this pseudo-Lagrangian approach no unrealistic or speculative assumptions about advection into the model domain have to be made. Based on the average horizontal wind speed of 3 m/s it would take six hours to transport an air parcel across the Dead Sea. During those six hours when the model is run over the water surface, the lower boundary conditions are changed representing the measured gradient in the acidity and salinity of the lake water which are factors that are assumed to be responsible for the stronger and more often occurring O₃ de-

Halogen chemistry over the Dead Sea

L. Smoydzin
and R. von Glasow

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



pletion events over the southern part of the Dead Sea compared to the northern part. Based on values reported by Tas et al. (2005) Br^- mixing ratios in the Dead Sea water are decreased from 0.15 mol/l (south) to 0.05 mol/l (north) and the pH of the lake is increased from 4.6 (south) to 6.4 (north). Obviously, this setup only changes the bromide content of the water and not the total salt content. Even the lowest reported Cl^- concentrations for the Dead Sea water (6.34 mol/l Tas et al., 2005) are very large compared to ocean water and sensitivity studies showed that they do not limit the extent of halogen chemistry.

An overview of the different scenarios and the abbreviations of each model run is given in Table 1. In all scenarios the model column moves over the Dead Sea from 10:00 until 16:00. The dry and hot meteorological conditions lead to a boundary layer height that is constantly at ≈ 1300 m if not mentioned otherwise.

We used the following Lagrangian model scenarios to simulate the effect of differing ambient conditions on the trace gas concentrations over the Dead Sea:

- NOEX: Air-sea exchange of gases is not explicitly calculated and only the “standard” gas phase and aerosol phase chemistry of the model MISTRA is taken into account.
- EX: Degassing of the halogen species BrCl , Br_2 and Cl_2 out of the Dead Sea water into the atmosphere is calculated explicitly.
- EXALL: Additionally to BrCl , Br_2 and Cl_2 , the air-sea exchange of HOBr , HOCl , HCl and HBr is considered explicitly. In scenarios EXLIM, ENH and LOWBL the same tracers are exchanged like in EXALL.
- EXLIM: For the first two hours of simulation time vertical transport is strongly limited. Such a situation might arise when hot and dry air masses that had been transported over the desert start to cross the water surface which is slightly cooler than the land surface. An inversion which is not very intense but strong enough to limit vertical transport builds up in the lower most boundary layer. After about two hours it is broken up due to vertical mixing of air masses.

Halogen chemistry over the Dead Sea

L. Smoydzin
and R. von Glasow

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- ENH: Fluxes of bromine species out of the Dead Sea water are increased in the first two hours of simulation time to simulate a potentially stronger bromine source. Salt pans which are mainly present in the southern part of the Dead Sea might be a strong additional bromine source. Furthermore, it is possible that the air-sea parameterisation underestimates the flux of bromine species out of the very shallow and highly concentrated evaporation ponds in the south.
- LOWBL: Initial meteorological conditions were chosen such, that the boundary layer height is constantly at about 500 m. To achieve this the temperature had to be decreased by about 3°C compared to scenario EXALL as can be seen in Fig. 2.

5.2 Stationary run

To investigate the daily variation of trace gas concentrations at one single point above the Dead Sea (corresponding to a measurement site), one stationary model run (STAT) over the south end of the Dead Sea (Dead Sea water with a pH of 4.8 and $[\text{Br}^-]=0.15 \text{ mol/l}$) was performed. The model run was initialised with the same fields for meteorology and chemistry as the Lagrangian studies, but the composition of the water as well as the composition of freshly emitted sea salt particles is not changed during the 38 h of simulation time. The model run starts at the same time as the Lagrangian runs (10:00) and all presented plots show mixing ratios between 10:00 day 1 and 20:00 day 2.

It is rather unrealistic, that the air at one particular site at the Dead Sea stays uninfluenced by the environment. Thus advective fluxes are assumed however, only of O_3 and NO_2 . As it is not possible to calculate advection into the model domain explicitly with a one-dimensional model, the strength of the fluxes is chosen somewhat arbitrarily (10:00 day 1–15:00 day 2: NO_2 : 0.463 pmol/(mol s), O_3 : 0.078 pmol/(mol s); 15:00 day 2–24:00 day 2: NO_2 : 0.347 pmol/(mol s), O_3 : 6.829 pmol/(mol s)). The inflow of both species into the model domain is assumed to be the same in all model levels. The

Halogen chemistry over the Dead Sea

L. Smoydzin
and R. von Glasow

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



assumption of an inflow in general is a strong constraint to the model. This scenario is not valid for representing accurately the conditions and daily variations at one particular site at the Dead Sea, but it should help to understand the general chemical mechanism and its sensitivity to changing background conditions.

6 Results: Lagrangian runs

6.1 Lagrangian runs without air-sea-exchange

First we will discuss the scenario ignoring direct air-sea exchange (NOEX). If not mentioned otherwise all line plots presented herein show mixing ratios at two metres altitude as most measurements were taken at approximately this height. Only the time period when the model column moves over the Dead Sea (from 10:00 to 16:00) is shown in all plots.

The maximum BrO mixing ratio as simulated in scenario NOEX is 0.012 pmol/mol (Fig. 3a). This value which is reached at 16:00, is up to four orders of magnitude lower than values reported from measurements. O₃ mixing ratios stay nearly constant at about 70 nmol/mol (Fig. 3b) during the six hours of simulation time between 10:00 (south) and 16:00 (north).

Without considering degassing out of the Dead Sea water the only source for gas phase bromine species is the transfer of halogen precursor species out of sea salt aerosols. Even though sea salt aerosols over the Dead Sea contain high Br⁻ and Cl⁻ concentrations (≈3 and 40 mmol/l Fig. 3c) and have a high acidity (pH≈1, Fig. 3c), it is not possible to reproduce the measurements with model setup NOEX. As the sea salt production is low and aerosol chemistry is only effective in the lowest 2 m above sea level due to the low relative humidity (see Fig. 2), the Br₂ flux out of aerosol particles ($6 \times 10^{-7} - 2.8 \times 10^{-6}$ pmol/(m³ s)) is too weak to produce significant bromine levels in the gas phase. Beside the vertical limitation of bromine production, strong vertical transport is another limiting factor for the accumulation of BrO in the lowest metres

Halogen chemistry over the Dead Sea

L. Smoydzin
and R. von Glasow

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



above ground as it leads to a quick vertical mixing of the anyway low concentrated tracers all over the boundary layer.

In general mixing ratios of gas phase halogen species increase with time (in terms of the Lagrangian model setup with travel distance), as at the south end of the Dead Sea (10:00) halogen concentrations are close to zero, and the longer halogen chemistry is active (the closer the model column gets to the north end of the lake) the more the concentrations increase. The relative increase (e.g. in BrO mixing ratios) is thus large, however, the absolute mixing ratios are very small.

Generally, the low relative humidity and thus the very low liquid water content of sea salt aerosols ($11 \times 10^{12} \text{ m}_{aq}^3 / \text{m}_{gas}^3$) is the reason for the slow recycling of bromine species on aerosol particles. The lower the liquid water content of an aerosol particle, the lower is the uptake rate of a gas phase species. The production of Br₂ in the aerosol phase by the bromine explosion reaction is essential for the formation of significant gas phase BrO levels. However, in addition to the high acidity and salinity in the aerosol, significant HOBr levels are required. As the halogen cycling between the gas- and aerosol phase is weak, HOBr uptake into aerosols is weak leading to the relatively small HOBr concentrations within sea salt particles ($\approx 5 \times 10^{-4} \text{ pmol/l}$). The recycling pathway via heterogeneous decomposition of BrONO₂ is inefficient as well due to the limited uptake of gas phase species on aerosol particles over the Dead Sea. Clearly the model setup only including the interacting gas – and aerosol phase chemistry is not appropriate to simulate ozone depletion events and BrO levels in the range of 100 pmol/mol over the Dead Sea. Therefore air-sea exchange of halogen species is taken into account.

6.2 Lagrangian runs with air-sea-exchange

6.2.1 Air-sea exchange and water chemistry

First, the fluxes of bromine and chlorine species out of (into) the water of the Dead Sea and chemistry taking place in the Dead Sea water are discussed. Only fluxes

Halogen chemistry over the Dead Sea

L. Smoydzin
and R. von Glasow

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Halogen chemistry
over the Dead Sea**L. Smoydzin
and R. von Glasow

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



for scenario EXALL are shown as the magnitude of fluxes as well as aqueous phase concentrations in the water do not differ significantly between the scenarios however, HOBr fluxes are not included in scenario EX. Figure 4a shows fluxes (in $\text{pmol}/(\text{m}^2 \text{ s})$) for Br_2 (red line), BrCl (black dashed line) and HOBr (blue dotted line) out of the Dead Sea water into the gas phase. Fluxes of all other halogen species are more than three orders of magnitude lower than the BrCl flux (which is the smallest flux shown) and are thus of minor importance and will not be discussed any further.

The strength of degassing of Br_2 ($80\text{--}154 \text{ pmol}/(\text{m}^2 \text{ s})$) stays on the same order of magnitude during the simulation time. The change with time for both other fluxes plotted in the same figure is much larger. BrCl and HOBr fluxes increase from 0.23 to $1.22 \text{ pmol}/(\text{m}^2 \text{ s})$ and 2 to $800 \text{ pmol}/(\text{m}^2 \text{ s})$, respectively. As explained above, Br^- and H^+ concentrations in the Dead Sea water are decreased during the six hours that the model column moves from the south to the north of the Dead Sea. Regarding those background conditions it is surprising that Br_2 fluxes stay nearly constant and HOBr and BrCl fluxes even increase with time.

Concentrations of HOBr in the water rise from $0.29 \text{ nmol}/\text{l}$ to $61 \text{ nmol}/\text{l}$ which is an increase of two orders of magnitude (Fig. 4b). Concentrations of Br_2 in the water slightly increase in the first three hours of simulation time from about $121 \text{ nmol}/\text{l}$ to $142 \text{ nmol}/\text{l}$ and then decrease slightly to about $136 \text{ nmol}/\text{l}$ (Fig. 4b). The most important reaction for activating bromine chemistry is the bromine explosion mechanism (red reaction path in Fig. 5):



Even though bromide concentrations and the acidity in the water are decreased with time in the model, they remain high enough such that the equilibrium reactions produce Br_2 throughout the simulation time (indicated through the larger red arrow in Fig. 5 pointing at Br_2). The forward reaction (of Reaction (1)) is getting only slightly slower in the last two hours between 14:00 and 16:00 as Br^- and H^+ get less abundant with time, leading to the small decrease in Br_2 concentrations followed by a decrease in the

**Halogen chemistry
over the Dead Sea**L. Smoydzin
and R. von Glasow

Br₂ flux. Even though Br⁻ concentrations decrease to about a third of their initial value after 6 h of simulation time (0.15 → 0.05 mol/l), the absolute concentrations at 16:00 are still very large. Furthermore, HOBr concentrations increase strongly with time, which will be explained below. Thus the ratio of the forward to the backward reaction of the equilibrium Reaction (1) even increases slightly with time, but the absolute reaction rate slightly decreases with time. This explains why the bromine explosion reaction does not get significantly slower with time and the decrease in Br₂ concentrations in the Dead Sea water with time is negligibly small. Additionally the increase of the pH to a value of 6.4 (thus the water is still slightly acidic) at 16:00 does not yet limit Br₂ formation significantly compared to a value of 5 (Fickert et al., 1999).

In contrast to Br₂, HOBr concentrations increase nearly exponentially with time. Besides for the conversion of Br⁻ into Br₂, HOBr is needed to convert Cl⁻ into BrCl (green reaction path in Fig. 5):



Due to the very high Cl⁻ and H⁺ concentrations this reaction is very fast. Throughout the 6 h of simulation time the backward reaction of the equilibrium Reaction (2) is faster (indicated by large green arrows in Fig. 5 pointing at HOBr and Cl⁻) than the forward reaction. Thus the backward reaction is a net source for HOBr in the Dead Sea water. In contrast, within aerosol particles, the forward reaction is in general faster than the backward reaction and BrCl is net produced. The explanation for the shift of the equilibria in the Dead Sea water can be found in the very high salt content and the very high and changing bromide content as well as the very low and changing pH of the Dead Sea water. Apart from Reaction (2), BrCl is produced by other pH dependent but non-equilibrium reactions within the water of which the fastest two are shown in Fig. 5 (blue and grey reaction path in Fig. 5):

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Thus the slow production of BrCl via Reaction (2) is overcompensated by BrCl production via Reactions (3) and (4). The increase of HOBr as well as of HOCl concentrations in the Dead Sea water from south to north can be seen as follows:

As both H^+ and Br^- concentrations decrease with time, less HOCl is consumed with time forming BrCl (via Reaction (3), blue reaction path in Fig. 5). This leads to an increase of HOCl concentrations followed by a stronger net production of ClO^- via the equilibrium reaction:



(black reaction path in Fig. 5) followed by a stronger production of BrCl (via Reaction (4), grey reaction path in Fig. 5). This now leads to an increase in BrCl concentrations followed by a stronger conversion to HOBr (via Reaction (2), green reaction path in Fig. 5). The strongly increasing HOBr concentrations in the water determine the strength of the HOBr flux into the atmosphere which exceeds the Br_2 flux (after 13:40) even though hypobromous acid is more soluble than Br_2 .

6.2.2 Gas phase chemistry

For both scenario EXALL and EX, BrO mixing ratios increase rapidly within a few minutes after the model column is assumed to cross the Dead Sea up to 10 pmol/mol and increase further up to 151 pmol/mol (EXALL) and 102 pmol/mol (EX, Fig. 6a) indicating that the Dead Sea water itself has to be considered as an important source for gas phase bromine species as the calculation of air-sea exchange is necessary in the model to simulate BrO mixing ratios on the same order of magnitude as field measurements show. The difference of about 50 pmol/mol BrO between scenario EX and EXALL can be explained with the very high gas phase HOBr mixing ratios in scenario EXALL ($[HOBr]_{max}=0.91$ nmol/mol, EX: $[HOBr]_{max}=0.14$ nmol/mol) caused by the additional HOBr source due to direct degassing out of the Dead Sea water (Fig. 6b). HOBr is under the prevailing circumstances an important precursor species for gas phase bromine as it is rapidly photolysed to form Br atoms (and OH radicals) which

Halogen chemistry over the Dead Sea

L. Smoydzin
and R. von Glasow

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Halogen chemistry
over the Dead Sea**L. Smoydzin
and R. von Glasow

react with O_3 to produce BrO . Under less dry atmospheric conditions, $HOBr$ is taken up into sea salt aerosols quickly due to its high solubility. However, as the liquid water content of aerosol particles is very low, the uptake of gas phase species in aerosol particles is not efficient. As the source of $HOBr$ due to degassing out of the Dead Sea water is large, a significant fraction of $HOBr$ gets photolysed and provides an efficient source for Br atoms.

The steadily and strongly increasing BrO mixing ratios in scenario EXALL and EX are caused by the constantly strong fluxes of bromine species out of the water. The most important source reaction of bromine atoms is photolysis of molecular bromine. Even though gas phase Br_2 gets quickly photolysed to Br (by Reaction (6)):



the reaction rate is in the range of $10\text{--}40 \text{ pmol}/(\text{m}^3 \text{ s})$ at 2 m altitude (EXALL). Regarding all Br_2 source and sink reactions, the large additional source of bromine compounds due to the air-sea exchange ($81\text{--}154 \text{ pmol}/(\text{m}^2 \text{ s})$) leads to an accumulation of bromine in the air. Even though the largest source and sink reactions for Br_2 are on the same order of magnitude, the source of gas phase Br_2 due to air-sea exchange is slightly larger than the Br_2 photolysis throughout the simulation time.

BrO mixing ratios in scenario EXALL tend to increase with time throughout the boundary layer. From 13:00 on mixing ratios exceed $5 \text{ pmol}/\text{mol}$ even at 1000 m altitude (Fig. 6d) and increase further at this height up to $20 \text{ pmol}/\text{mol}$ at 16:00, when the model column reaches the north shore of the Dead Sea. There is a delay of about one hour between the start of the model run at 10:00 when BrO surface values exceed $1 \text{ pmol}/\text{mol}$ after a few minutes until BrO mixing ratios in 1000 m also exceed $1 \text{ pmol}/\text{mol}$ at 11:00.

Even though BrO gets quickly distributed all over the boundary layer, the strong vertical gradient in mixing ratios is striking (note the logarithmic scaling of the contour

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

lines in plots showing BrO mixing ratios in Fig. 6d, e). BrO mixing ratios at e.g. 15:30 are less than half the ground level values at 10 m altitude (240 pmol/mol → 90 pmol/mol) and at 100 m altitude they are almost a factor of four smaller than mixing ratios directly above the Dead Sea water surface.

5 The influence of those high bromine concentrations, as simulated in scenario EXALL (and EX), on O₃ chemistry are still small though. The decrease from 69 to 63 nmol/mol (EXALL) is insignificant compared to the observed depletion events. The difference in O₃ mixing ratios in scenario EXALL means an O₃ reduction of only 9%.

10 In Scenario EXLIM vertical transport is limited in the first two hours of simulation time leading to the sharp increase in BrO mixing ratios between 10:00 and 11:40 from below 1 up to 395 pmol/mol and the likewise quick decrease to about 30 pmol/mol between 11:40 and 13:00 when the inversion is broken up (Fig. 6b, blue dash-dotted line). Simultaneous to the very fast increase of BrO mixing ratios in the first 1.5 h that the model crosses the Dead Sea, O₃ mixing ratios decrease until they reach a minimum of 9.3 nmol/mol at 11:40 (Fig. 6c, blue dash-dotted line). However, as soon as the low-level inversion is compensated, O₃ quickly reaches background mixing ratios of about 69 nmol/mol due to vertical downmixing of O₃.

20 The limited O₃ destruction in scenario EXALL might be surprising at a first glance regarding the extraordinarily high BrO levels. Even though the net O₃ destruction is larger than the net O₃ production close to the Dead Sea water surface, the contribution of the strong O₃ sink due to reaction with halogens to the total O₃ mass balance is still not large enough to lead to a distinctive ODE.

25 As mixing ratios of bromine species show a strong vertical gradient, bromine mixing ratios in the upper boundary layer are not large enough anymore to destroy significant O₃ levels. Thus vertical downmixing of O₃ rich air from the upper boundary layer compensates the net O₃ destruction in the lower most metres of the boundary layer. However, bromine mixing ratios permanently increase with time all over the boundary layer such that O₃ destruction cannot be totally compensated anymore by vertical mixing leading to the slight decrease in O₃ mixing ratios between 13:00 and 16:00.

Halogen chemistry over the Dead Sea

L. Smoydzin
and R. von Glasow

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

As was discussed above O_3 is depleted strongly and quickly in scenario EXLIM between 10:30 and 11:30. As bromine mixing ratios are high only in the lowest metres above sea level due to the limited vertical transport (Fig. 6e), O_3 destruction is also restricted to the lowest 10–20 m of the boundary layer. Above this inversion layer O_3 mixing ratios are again hardly affected by bromine chemistry. As soon as the inversion is broken up, O_3 gets again rapidly well mixed throughout the whole boundary layer. After 13:30, when the boundary layer height quickly rises to 1300 m, BrO is also quickly distributed all over the boundary layer and mixing ratios are between 2 pmol/mol and 50 pmol/mol all over the boundary layer (Fig. 6e).

6.3 Enhanced fluxes

With all of the Lagrangian scenarios presented above neither an ozone depletion event affecting the whole boundary layer could be simulated nor any significant O_3 depletion in case of a deep and well mixed boundary layer. For scenario ENH the fluxes of halogen species out of the Dead Sea water were enhanced for the first two hours of simulation time which is the time the model is roughly assumed to cross the evaporation ponds.

Fluxes were enhanced by a factor of 30 until 10:30, a factor of 20 until 11:00 and a factor of 10 until 12:00. Afterwards, the fluxes as calculated by the air-sea exchange parameterisation are not modified anymore. BrO mixing ratios peak at 10:30, 11:00 and 12:00 as a response to the strong and instantaneous changes in the flux strength of BrO precursor species. Between 10:00 and 11:00 BrO mixing ratios reach values between 600 and 750 pmol/mol (red dotted line, Fig. 7c). Simultaneously O_3 mixing ratios decrease from 65 to 47 nmol/mol (black line, Fig. 7c). As the strength of degassing is again decreased after 11:00, BrO mixing ratios decrease quickly to about 300 pmol/mol and drop further to about 100 pmol/mol after 12:00 and stay more or less constant around this value until 16:00. During the same time span O_3 mixing ratios decrease constantly, but not very fast and reach a minimum value of 13 nmol/mol at 16:00.

Halogen chemistry over the Dead Sea

L. Smoydzin
and R. von Glasow

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Halogen chemistry
over the Dead Sea**L. Smoydzin
and R. von Glasow

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

The slow depletion of O_3 affects the whole boundary layer up to an altitude of 1200 m (Fig. 7a). BrO mixing ratios are approximately 100 pmol/mol all over the boundary layer after 12:00 in scenario ENH whereas mixing ratios never exceeded 50 pmol/mol in altitudes greater than 100 m in scenario EXALL. Due to the enhanced fluxes during the first two hours of simulation time the bromine load all over the boundary layer is significantly higher in scenario ENH than in scenario EXALL during the whole simulation time. As a consequence O_3 is destroyed all over the boundary layer by reaction with bromine in scenario ENH, such that vertical mixing cannot compensate the depletion of O_3 anymore as it happened in scenario EXALL. However, the reaction between bromine and O_3 is not fast enough to lead to an immediate ODE such that a delay between the time of maximum BrO and minimum O_3 mixing ratios can be seen (Fig. 7a, c).

For scenario LOWBL different meteorological background conditions were assumed leading to a constantly much lower boundary layer (500 m) than in all other so far presented Lagrangian studies. Note that therefore the temperature in the atmosphere had to be decreased by more than 3°C (see Fig. 2) compared to the other scenarios and thus the relative humidity in scenario LOWBL is higher than in other scenarios all over the boundary layer (see Fig. 2). This also indicates that aerosol phase chemistry gains in importance as it is not only calculated in the lowest 2–4 m but also constantly in altitudes exceeding 300 m. BrO gas phase mixing ratios increase with time as it was seen before (see Fig. 7b, d). Compared to scenario EXALL, they increase stronger with time ($[BrO]_{max}=207$ pmol/mol (LOWBL) versus $[BrO]_{max}=151$ pmol/mol (EXALL)). Simultaneously to the increase in BrO mixing ratios, O_3 mixing ratios decrease from 65 to 25 nmol/mol at 2 m altitude. As in scenario ENH the depletion of O_3 affects the whole boundary layer (Fig. 7b).

7 Stationary runs

To study the time evolution of trace gases within two days at a single site at the Dead Sea and to study the influence of changing meteorological background conditions on tracer mixing ratios, scenario STAT was calculated. In comparison to scenario EX, BrO mixing ratios on day 1 in scenario STAT are slightly lower: EX: $[\text{BrO}]_{\text{max}}$: 100 pmol/mol, STAT: $[\text{BrO}]_{\text{max}}$: 60 pmol/mol (Fig. 8a). NO_2 mixing ratios are higher during the whole day in scenario STAT compared to scenario EX due to the prescribed advection of NO_2 into the model domain (day 1: $[\text{NO}_2]_{\text{STAT}} \approx 0.8$ nmol/mol, $[\text{NO}_2]_{\text{EX}} \approx 0.3$ nmol/mol). Thus comparatively more BrO is consumed via the reaction



leading mainly to the difference in BrO mixing ratios at day 1 between scenarios STAT and EX. On the second day of simulation time, BrO mixing ratios in scenario STAT increase stronger than on day 1 and reach a maximum value of 140 pmol/mol at 2 m altitude at about 10:00 in the morning. Mixing ratios decrease again after 10:00 (34 h in Fig. 8=10:00 day 2) but are always larger than on day 1. BrO mixing ratios exceed 50 pmol/mol up to an altitude of about 1200 m almost all day long whereas on day 1 they were always lower than 12 pmol/mol in heights larger than 100 m (Fig. 8b, e).

Maximum HOBr mixing ratios at 2 m altitude on day 1 of the simulation time are of similar order of magnitude in scenarios STAT and EX. However, on the second day of simulation time, HOBr mixing ratios also increase to larger values than on day 1 (day 1: $[\text{HOBr}]_{\text{STAT}} \approx 174$ pmol/mol, $[\text{HOBr}]_{\text{EX}} \approx 165$ pmol/mol day 2: $[\text{HOBr}]_{\text{STAT}} \approx 523$ pmol/mol, Fig. 8a, b).

Comparing the results from scenarios STAT and EX, the vertical distribution and the time evolution of O_3 mixing ratios is surprising. On the first day of simulation time, O_3 mixing ratios stay constant at around 70 nmol/mol and are not affected at all by halogen chemistry. On the second day (30 h in Fig. 8=06:00 day 2) O_3 mixing ratios start to decrease immediately after sunrise, when photochemistry and thus bromine chemistry starts getting activated. At 14:00, O_3 mixing ratios at 2 m altitude reach

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

a minimum value of 5 nmol/mol and increase again afterwards mainly because of the prescribed O₃ flux and because mixing ratios of bromine species decrease in the late afternoon hours.

In contrast to the Lagrangian scenarios discussed above, O₃ is strongly depleted up to an altitude of about 1200 m even though the boundary layer is high and well mixed throughout the simulation time. Even though BrO mixing ratios are on the same order of magnitude as in e.g. in scenario EXALL where no influence on O₃ chemistry could be found, the bromine oxide production in scenario STAT is sufficiently large to lead to a significant O₃ depletion event.

The explanation for the strong O₃ depletion is related to slightly changing meteorological background conditions from day 1 to day 2. The temperature is slightly lower and the relative humidity slightly higher on day 2 compared to day 1. Under the very hot and dry conditions prevailing along the Dead Sea, sea salt particles dry out quickly after they have been injected into the atmosphere. As explained in the model description the ambient relative humidity has to exceed the aerosol's deliquescence humidity of 75% (70% for sulphate aerosols) to reactivate the calculation of sea salt (and respectively sulphate) aerosol chemistry in the model. On day 1 of the simulation time the relative humidity is below this threshold value all over the boundary layer but on day 2 it exceeds 75% at altitudes above 800 m (and 70% at altitudes above 650 m) and thus aerosol phase (sulphate and sea salt) chemistry is reactivated (Fig. 8b). As a result, aqueous phase chemistry and the cycling of halogens on aerosol particles gains in importance. The liquid water content of sea salt aerosols (Fig. 8b) is an indicator for levels where aerosol phase chemistry (on sea salt and sulphate aerosols) is calculated. Within all levels having a LWC < 10¹² m³_{aq}/m³_{gas} in Fig. 8b, liquid phase chemistry is not activated (note again the logarithmic scaling of the y-axis in all plots showing vertical distributions!). An efficient recycling process is the heterogeneous decomposition of BrONO₂ on sulphate aerosols and the subsequent release of HOBr out of aerosol particles:



4545

Halogen chemistry over the Dead Sea

L. Smoydzin
and R. von Glasow

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back


Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





Through strong vertical transport, also the lower part of the boundary layer is indirectly influenced by the enhanced halogen chemistry in the upper parts of the boundary layer. This can be seen in the vertical distribution of gas phase HOBr mixing ratios (Fig. 8b).

5 As the total halogen recycling capacity all over the boundary layer is enhanced, O_3 is depleted even though BrO mixing ratios do not exceed 200 pmol/mol.

Even though the cycling of halogen species between the aerosol phase and the gas phase contributes to the increase in the gas phase halogen mixing ratios, the air-sea exchange of Br_2 and BrCl is still the dominant source for BrO especially in the lowest
10 part of the boundary layer. BrO mixing ratios still show a strong vertical gradient. In addition to the changing meteorology, halogen mixing ratios are higher in the morning of the second day of the simulation than on the first day. Br_2 mixing ratios before sunrise on day 2 are 470 pmol/mol whereas they increase from 0 at 10:00 on day 1 when the model simulations start ($[\text{Br}_2]_{\text{STAT}}$: 100 pmol/mol at 10:00 day 2).

15 The inflow of O_3 into the model domain was increased in the afternoon hours of day 2 of simulation time, as otherwise O_3 mixing ratios would not increase anymore. Almost all measurements show a quick increase of O_3 mixing ratios in the late afternoon/early evening hours. However, a flux of 6.8 pmol/(mol s) of O_3 was necessary to reach mixing ratios of at least 40 nmol/mol at 20:00 day 2 in the model.

20 8 Discussion

Regarding the measurements taken at the Dead Sea in more detail, it can be seen that BrO mixing ratios in the north reach very regular maximum mixing ratios at noon and the standard variation of the measurements is very small. In the south however, the time of maximum BrO mixing ratios varies strongly as well as the absolute values of
25 maximum BrO mixing ratios (Fig. 2 in Tas et al., 2003). In addition O_3 seems to be destroyed much stronger and more often over the southern end of the Dead Sea than the

Halogen chemistry over the Dead Sea

L. Smoydzin
and R. von Glasow

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Halogen chemistry
over the Dead Sea**L. Smoydzin
and R. von Glasow

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

northern end (Tas et al., 2005; Zingler and Platt, 2005). With the Lagrangian scenario EXLIM it was possible to reproduce these measurements showing high BrO mixing ratios (>60 pmol/mol) all over the Dead Sea but an ODE only in the south. Results agree well regarding the strength of the ODE, the spatial distribution of O_3 and BrO as well as the temporal evolution of the depletion event meaning a very quick O_3 decrease when air masses are assumed to reach the Dead Sea from the south (see Fig. 6). Nevertheless, the presence of BrO mixing ratios of almost 400 pmol/mol at 2 m altitude are necessary to destroy a significant amount of O_3 in the model. However, it is possible that the effect of a low level inversion is overestimated in the model and the inversion is compensated faster in reality. As a consequence, BrO mixing ratios would not become as high as in scenario EXLIM and O_3 mixing ratios would still decrease rapidly, even though not as strongly as in this model scenario. In order to test this, scenario EXLIM was recalculated only allowing the exchange of Br_2 and BrCl (not shown). In that case BrO mixing ratios reached a maximum mixing ratio of 290 pmol/mol (in contrast to 395 pmol/mol in EXLIM) and O_3 still decreased quickly to 13 nmol/mol (in contrast to 10 nmol/mol in EXLIM). Furthermore, in scenario EXLIM only the lowest 10 to 20 m are affected by the depletion event. Nevertheless, a similar scenario might develop in reality affecting not only the lowest 10, but maybe 100 or 300 m. It is not possible to evaluate those model results, as the above mentioned measurements were only taken at one level at approximately 2 m height and no information about the vertical extent of the real depletion events or BrO mixing ratios is available.

The assumption that the wind direction is southerly all over the lake might be idealised as the wind direction at the south and north end of the Dead Sea might be of opposite direction due to the lake breeze. However, even under such circumstances an air parcel reaching the northern end of the Dead Sea has been transported roughly across half the length of the lake and BrO mixing ratios reached significant values already after three hours of transportation time in scenario EXALL.

Regarding the assumption that air masses reaching the south of the Dead Sea coming from the desert have a negligible halogen load might not be true under all circum-

**Halogen chemistry
over the Dead Sea**L. Smoydzin
and R. von Glasow

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

stances. As was observed by Tas et al. (2005) air masses reaching the Dead Sea from the desert south of the lake can already contain a significant load of halogens (up to 20 pmol/mol Tas et al., 2005) which Tas et al. (2005) attributed to recirculations of air masses. For such cases the Lagrangian setup underestimates initial halogen concentrations. However, the different simulations presented herein indicate that an additional halogen load of this reported order of magnitude would not lead to a much faster and much stronger O₃ decrease than shown in the modelled cases.

Apart from the uncertainties regarding the influence of meteorological background conditions on trace gas concentrations, one additional uncertainty is the actual source strength of bromine species degassing out of the Dead Sea water. As the evaporation ponds are very shallow, the water within the ponds have a highly concentrated bromide content and the depth of each pond is different and is varied regularly, the parameterisation used to calculate air-sea exchange might underestimate the bromine source strength over the evaporation ponds. However, it is difficult to say quantitatively how large this potential error might be. It was already suggested by Matveev et al. (2001) that the bromine source strength might be enhanced as the wet soil at the shore of the lake contains bromine-rich sea salt which can provide an additional reactive surface. This idea is supported by Stutz et al. (2002) in order to explain BrO, ClO and O₃ measurements at the Great Salt Lake in Utah. In addition to the variable wind direction over the southern part of the Dead Sea, the potentially very variable bromine source strength due to a variable bromine release strength of each single pond might explain the unregular pattern of BrO mixing ratios as detected at the evaporation ponds.

In order to test the effect of an additional bromine source over the southern part of the Dead Sea, scenario ENH was calculated. O₃ is depleted in this scenario even though more than 700 pmol/mol BrO are formed. It was observed by Matveev et al. (2001) that ODEs can occur only affecting the northern part of the Dead Sea like it is simulated in scenario ENH. Again it is not possible to evaluate the model results of scenario ENH and scenario LOWBL as well, regarding the vertical extent of the depletion event due to the lack of vertically resolved measurements.



**Halogen chemistry
over the Dead Sea**L. Smoydzin
and R. von Glasow

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

As mentioned in the introduction a few measurements took place on an elevated site 400 m above water level. Only at 3 out of 22 days BrO (≈ 20 pmol/mol) was detected and thus simulated values (EX, EXALL) are on the same order of magnitude at this altitude. Tas et al. (2003) assumed that on those three days where BrO mixing ratios were detected, the boundary layer height was very low. However, in scenario EXALL the boundary layer height is about 1300 m and BrO is transported very efficiently with high concentrations into altitudes around 400 m. This let the question arise, what kind of local wind and transport pattern is responsible for the *lack* of BrO on the other days.

Scenario STAT is the only scenario presented, where it was possible to simulate an ODE affecting quickly the whole BL even though the boundary layer height is constantly high (about 1200 m) and air-sea exchange of tracers was not enhanced. This idealised scenario represents two days where air masses in the Dead Sea valley are not influenced by mixing with air masses from aloft or advection. One of the main conclusions which can be drawn from scenario STAT is that it is important not only to consider chemistry as an isolated system as apparently very small changes in the meteorological background conditions (namely temperature and humidity) give a large feedback to chemistry.

In general results from model simulations presented herein agree very well with a first simple model study by Stutz et al. (1999) who concluded that the production of 500 pmol/mol BrO would be essential to simulate a significant loss in O₃ within at least three hours. However, results from both studies differ significantly from what is presented by Tas et al. (2006) and their follow up paper (Tas et al., 2008).

Gas phase halogen mixing ratios depend and are strongly controlled by the cycling between the aerosol phase and the gas phase. Tas et al. (2006) state correctly that the heterogeneous decomposition of BrONO₂ is an important step in the recycling process of reactive halogen species. However, the chemical mechanism as used by Tas et al. (2006) only treats gas phase chemistry explicitly. Adding one aqueous phase and one heterogeneous reaction as a parameterisation as done by them disrupts the sensitive aerosol phase/gas phase chemical cycling. The approach by Tas et al. (2006) is in-

**Halogen chemistry
over the Dead Sea**L. Smoydzin
and R. von Glasow

consistent as e.g. the heterogeneous decomposition of bromine nitrate is a source for gas phase HOBr but Tas et al. (2006) neglect the uptake of HOBr on aerosol particles as an equivalent sink. Nevertheless, we could show that under the prevailing meteorological conditions at the Dead Sea with temperatures exceeding 40°C regularly in summer and a relative humidity of around 50% or even below (Alpert et al., 1997, and references therein, personal communication M. Piot, Inst. of Env. Physics, Univ. Heidelberg, Germany) heterogeneous or phase transfer reactions are slow. Thus the recycling potential of aerosol particles is overestimated by Tas et al. (2006). However, Tas et al. (2006) do not provide any information regarding temperature or humidity as used in their model study. Furthermore, Tas et al. (2006) chose an unrealistically high reaction rate of 1.35×10^{-6} nmol/s for the parameterised aerosol phase reaction: $\text{HOBr} + \text{Br}^- + \text{H}^+ \rightleftharpoons \text{Br}_2$. Under the prevailing circumstances the first order reaction rate as calculated in scenario NOEX in MISTRA is about two orders of magnitude slower at 2 m altitude ($1\text{--}3 \times 10^{-8}$ nmol/(m³ s)). In general, it has to be questioned that, choosing this reaction rate as the only degree of freedom in a numerical model, as done by Tas et al. (2006), and varying this rate until model simulations agree best with measurements is an appropriate approach to explain observations.

The main reason for the termination of aerosol chemistry in MISTRA is the low water content of aerosol particles due to the low ambient humidity. In order to enhance the recycling potential of halogens, scenarios NOEX and EXALL were calculated including heterogeneous reactions on dry aerosols based on several laboratory measurements (<http://jpldataeval.jpl.nasa.gov>; Sander et al., 2006). This setup leads to higher BrO mixing ratios due to the enhanced cycling potentials of halogen species. However, the enhancement of BrO formation by about 20% compared to scenario EXALL and by about 66% compared to scenario NOEX does (i) not lead to a stronger O₃ destruction and (ii) in comparison to scenario NOEX the formation of BrO (BrO_{max}: 0.02 pmol/mol) is still significantly lower than observations show.

In all of the above presented Lagrangian scenarios (apart from EXLIM), the O₃ depletion process is much slower than observations show it happens. A similar discrepancy

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Halogen chemistry
over the Dead Sea**L. Smoydzin
and R. von Glasow

between box model results and measurements was also stated by Stutz et al. (1999). In the model simulations presented by Tas et al. (2006), O_3 is destroyed quickly however, Tas et al. (2006) do not provide any information about the boundary layer height or the vertical distribution of BrO and O_3 mixing ratios. Tas et al. (2006) chose the strength of fluxes into the model domain such, that their simulations agree best with measurements. A problem in their line of argument is that their results strongly depend on the (“manually”) varied O_3 inflow. Measurements show a quick decrease but also a quick increase in O_3 mixing ratios. Tas et al. (2006) state that *“the decrease in O_3 concentrations beneath a threshold value of 1–2 nmol/mol acts as a motive force for fresh O_3 fluxes to move into the evaporation ponds.”* However, the advection of O_3 is controlled by the wind speed (although it is low over the Dead Sea) and wind direction and not only by the gradient in O_3 concentrations itself.

9 Conclusions

Regarding O_3 depletion events at the Dead Sea, not only the strength of the O_3 destruction but also the short timescale is striking. Additionally, the strong spatial and temporal variability of observed ODEs at the Dead Sea and the comparatively large range of corresponding BrO mixing ratios (≈ 40 – 140 pmol/mol) complicates the finding of an overall explanation of the occurring processes.

In general the model results presented herein clearly indicate that air-sea exchange of halogens between the Dead Sea and the atmosphere have to be considered in order to simulate BrO mixing ratios in the same order of magnitude as observations show.

A fast and strong ODE could only be simulated if an inversion was assumed that limits vertical transport. In addition an ODE develops if either the flux of halogens out of the Dead Sea was temporally enhanced or if the BL height did not exceed 700 m. In both cases however the time evolution of the ODE was much slower than often observed at the Dead Sea. It is difficult to evaluate the model results properly due to missing information regarding meteorological background conditions at the measure-

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ment days. In order to improve our understanding of O₃ and halogen chemistry at the Dead Sea, the knowledge of temperature, relative humidity and the boundary layer height for every single measurement day is absolutely crucial.

It would be of interest to have a data set showing the vertical distribution of BrO and O₃. Model simulations indicate that the vertical gradient in BrO mixing ratios is very strong in the lowest 10 to 20 m above the water surface. All published measurements were done with the DOAS technique at an altitude of approximately 2 m. According to the model results it is even of significance whether measurements are taken at 1.50 m or 2.30 m or 10 m. We have no reliable information regarding the vertical extend of an ODE and the strength of an ODE in the upper boundary layer. Overall it can be concluded that meteorological processes seem to play a key role in understanding the observed fast O₃ depletion events. Chemical processes lead to the formation of BrO and the destruction of O₃ however, the strength and spacial extend of the ODEs as well as the distribution of halogen species within the boundary layer strongly depend on the local meteorology.

Acknowledgements. This project was funded by the Deutsche Forschungsgemeinschaft DFG (Emmy Noether Junior Research Group MarHal GL 353/1-2).



This Open Access Publication is
financed by the Max Planck Society.

MAX-PLANCK-GESELLSCHAFT

References

- Alpert, P., Shafir, H., and Issahary, D.: Recent changes in the climate at the Dead Sea – a preliminary study, *Climatic Change*, 37, 513–537, 1997. 4530, 4550
- Barrie, L. A., Bottenheim, J. W., Schnell, R. C., Crutzen, P. J., and Rasmussen, R. A.: Ozone

ACPD

9, 4525–4565, 2009

Halogen chemistry over the Dead Sea

L. Smoydzin
and R. von Glasow

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- destruction and photochemical reactions at polar sunrise in the lower Arctic atmosphere, *Nature*, 334, 138–141, 1988. 4526
- Clarke, R. H.: Recommended methods for the treatment of the boundary layer in numerical models, *Aust. Meteorol. Mag.*, 18, 51–73, 1970. 4529
- 5 Fickert, S., Adams, J. W., and Crowley, J. N.: Activation of Br₂ and BrCl via uptake of HOBr onto aqueous salt solutions, *J. Geophys. Res.*, 104, 23719–23727, 1999. 4538
- Hebestreit, K., Stutz, J., Rosen, D., Matveev, V., Peleg, M., Luria, M., and Platt, U.: DOAS measurements of tropospheric bromine oxide in mid-latitudes, *Science*, 283, 55–57, 1999. 4526, 4527
- 10 Hönninger, G., Bobrowski, N., Palenque, E. R., Torrez, R., and Platt, U.: Reactive bromine and sulfur emissions at Salar de Uyuni, Bolivia, *Geophys. Res. Lett.*, 31, L04101, doi:10.1029/2003GL018818, 2004. 4527
- Jaenicke, R.: Aerosol Physics and Chemistry, in: Zahlenwerte und Funktionen aus Naturwissenschaften und Technik, Landolt-Börnstein, “Zahlenwerte und Funktionen aus Naturwissenschaften und Technik”, V 4b, 391–457, Springer, 1988. 4530
- 15 Liss, P. S. and Slater, P. G.: Flux of gases across the air-sea interface, *Nature*, 147, 181–184, 1974. 4531
- Matveev, V., Peleg, M., Rosen, D., Tov-Alper, D. S., Hebestreit, K., Stutz, J., Platt, U., Blake, D., and Luria, M.: Bromine oxide – ozone interaction over the Dead Sea, *J. Geophys. Res.*, 106, 10375–10387, 2001. 4527, 4548
- 20 Monahan, E. C., Spiel, D. E., and Davidson, K. L.: A Model of Marine Aerosol Generation via Whitecaps and Wave Disruption, in: *Oceanic Whitecaps*, edited by: Monahan, E. C. and Niocaill, G. M., 167–174, D. Reidel, Norwell, Mass, 1986. 4530
- Niemi, T., Ben-Avraham, Z., and Gat, J.: *The Dead Sea: The Lake and Its Setting*, Oxford Monogr. Geol. Geophys., vol. 36, Oxford Univ. Press, New York, 1997. 4529
- 25 Pozzer, A., Jöckel, P., Sander, R., Williams, J., Ganzeveld, L., and Lelieveld, J.: Technical Note: The MESSy-submodel AIRSEA calculating the air-sea exchange of chemical species, *Atmos. Chem. Phys.*, 6, 5435–5444, 2006, <http://www.atmos-chem-phys.net/6/5435/2006/>. 4531
- 30 Sander, S. P., Finlayson-Pitts, B. J., Friedl, R. R., Golden, D. M., Huie, R. E., Keller-Rudek, H., Kolb, C. E., Kurylo, M. J., Molina, M. J., Moortgat, G. K., Orkin, V. L., Ravishankara, A. R., and Wine, P. H.: *Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies*, JPL Publication 06-2, Jet Propulsion Laboratory, Pasadena, Evaluation Number 15, 2006.

**Halogen chemistry
over the Dead Sea**L. Smoydzin
and R. von Glasow

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Stutz, J., Hebestreit, K., Alicke, B., and Platt, U.: Chemistry of halogen oxides in the troposphere: comparison of model calculations with recent field data, *J. Atmos. Chem.*, 34, 65–85, 1999. 4527, 4549, 4551

5 Stutz, J., Ackermann, R., Fast, J. D., and Barrie, L.: Atmospheric reactive chlorine and bromine at the Great Salt Lake, Utah, *Geophys. Res. Lett.*, 29(10), 1380, doi:10.1029/2002GL014812, 2002. 4527, 4548

Sverdrup, H., Johnson, M., and Fleming, R.: *The Oceans, Their Physics, Chemistry and General Biology*, Prentice-Hall, Englewood Cliffs, N. J., 1942. 4529

10 Tas, E., Matveeva, V., Zingler, J., Luria, M., and Peleg, M.: Frequency and extent of ozone destruction episodes over the Dead Sea, Israel, *Atmos. Environ.*, 37, 4769–4780, 2003. 4527, 4529, 4530, 4546, 4549

Tas, E., Peleg, M., Matveev, V., Zingler, J., and Luria, M.: Frequency and extent of bromine oxide formation over the Dead Sea, *J. Geophys. Res.*, 110, D11304, doi:10.1029/2004JD005665, 2005. 4527, 4529, 4533, 4547, 4548

15 Tas, E., Peleg, M., Pedersen, D. U., Matveev, V., Biazar, A. P., and Luria, M.: Measurement-based modeling of bromine chemistry in the boundary layer: 1. Bromine chemistry at the Dead Sea, *Atmos. Chem. Phys.*, 6, 5589–5604, 2006, <http://www.atmos-chem-phys.net/6/5589/2006/>. 4527, 4549, 4550, 4551

20 Tas, E., Peleg, M., Pedersen, D. U., Matveev, V., Biazar, A. P., and Luria, M.: Measurement-based modeling of bromine chemistry in the Dead Sea boundary layer – Part 2: The influence of NO₂ on bromine chemistry at mid-latitude areas, *Atmos. Chem. Phys.*, 8, 4811–4821, 2008, <http://www.atmos-chem-phys.net/8/4811/2008/>. 4549

25 Vogt, R., Crutzen, P. J., and Sander, R.: A mechanism for halogen release from sea-salt aerosol in the remote marine boundary layer, *Nature*, 383, 327–330, 1996. 4528

von Glasow, R. and Crutzen, P.: *Tropospheric Halogen Chemistry*, in: *Treatise on Geochemistry Update 1*, Vol. 4.02, edited by: Holland, H. D. and Turekian, K. K., 1–67, Elsevier-Pergamon, Oxford, 2007. 4528

30 von Glasow, R., Sander, R., Bott, A., and Crutzen, P.: Modeling halogen chemistry in the marine boundary layer 1. Cloud-free MBL, *J. Geophys. Res.*, 107, 4341, doi:10.1029/2001JD000942, 2002a. 4530

von Glasow, R., Sander, R., Bott, A., and Crutzen, P. J.: Modeling halogen chemistry in the

Halogen chemistry over the Dead Sea

L. Smoydzin
and R. von Glasow

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



marine boundary layer 2. Interactions with sulfur and cloud-covered MBL, J. Geophys. Res., 107, 4323, doi:10.1029/2001JD000943, 2002b. 4530
Zingler, J. and Platt, U.: Iodine oxide in the Dead Sea Valley: Evidence for inorganic sources of boundary layer IO, J. Geophys. Res., 110, D07307, doi:10.1029/2004JD004993, 2005. 4547

ACPD

9, 4525–4565, 2009

Halogen chemistry over the Dead Sea

L. Smoydzin
and R. von Glasow

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Halogen chemistry over the Dead Sea

L. Smoydzin
and R. von Glasow

Table 1. Overview of all scenarios.

NOEX	No air-sea exchange of gases is calculated
EX	Air-sea exchange of only BrCl, Br ₂ , Cl ₂ is calculated
EXALL	The same setup as EX, but air-sea exchange of BrCl, Br ₂ , Cl ₂ , HOCl, HOBr, HCl, HBr is calculated
EXLIM	The same setup as EXALL, but limited vertical transport for first 90 min, strong vertical mixing for rest of simulation time
ENH	The same setup as EXALL, but strong hot spot emissions within the first 2 h
LOWBL	The same setup as EXALL, but boundary layer height only 500 m
STAT	The same initial conditions as EX, but model runs for 38 h over the same site at the south end of the Dead Sea

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Halogen chemistry over the Dead Sea

L. Smoydzin
and R. von Glasow

Table 2. Summary of results of scenarios calculated for the Dead Sea. All values refer to mixing ratios at 2 m altitude.

Scenario	BrO _(max) [pmol/mol]	O _{3(min)} [nmol/mol]	Δ O ₃	
NOEX	0.012	const 69	0%	No air-sea-exchange, BrO < 1 pmol/mol, O ₃ not affected
EX	102	66	4%	High BrO, No ODE
EXALL	151	63	9%	High BrO, No ODE
EXLIM	347	10	84%	ODE affects only lowest 20 m
ENH	375	13	80%	ODE affects whole BL, O ₃ decrease still slow but steady
LOWBL	207	25	63%	ODE affects whole BL, less strong than when enhancing fluxes
STAT	day 1: 61	day 1: 70	0%	No ODE on day 1 but ODE on day 2 affecting the whole BL due to slightly differing meteorological background conditions
	day 2: 139	day 2: 5	92%	

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Halogen chemistry over the Dead Sea

L. Smoydzin
and R. von Glasow




Fig. 1. Simplified overview of the interacting chemistry in the water, the gas phase and aerosol particles.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Halogen chemistry
over the Dead SeaL. Smoydzin
and R. von Glasow

Fig. 2. The plot on the left shows the vertical gradient of the temperature ($^{\circ}\text{C}$, solid lines) and the potential temperature ($^{\circ}\text{C}$, dashed lines) at 11:30 for scenarios EXALL (black) and LOWBL (red). The plot on the right shows the vertical gradient of the relative humidity (in %) at 11:30 for scenarios EXALL (black solid line) as well as scenario LOWBL (red dashed line).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)


Halogen chemistry
over the Dead SeaL. Smoydzin
and R. von Glasow

Fig. 3. Plots (a) and (b) show gas phase mixing ratios at 2 m altitude for scenario NOEX. Aqueous phase concentrations in sea salt aerosols at again 2 m height for scenario NOEX are shown in (c).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)


Halogen chemistry
over the Dead SeaL. Smoydzin
and R. von Glasow

Fig. 4. The fluxes of Br₂ (red solid line), BrCl (black dashed line) and HOBr (blue dotted line) out of the Dead Sea water are shown (a). Plots (b) and (c) show concentrations of several bromine containing species in the Dead Sea Water. All plots refer to scenario EXALL.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Halogen chemistry over the Dead Sea

L. Smoydzin
and R. von Glasow




Fig. 5. Chemical aqueous phase equilibria. Large arrows indicate whether the forward or backward reaction tends to dominate.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Halogen chemistry
over the Dead SeaL. Smoydzin
and R. von Glasow

Fig. 6. Plotted are gas phase mixing ratios at 2 m altitude for scenarios EX (red dotted line), EXALL (black line) and EXLIM (blue dash-dotted line) for the six hours that the model crosses the Dead Sea. The vertical distribution of O₃ (shaded) and BrO (contour lines) is shown in addition for scenarios EXALL (d) and EXLIM (e). Note the logarithmic scaling of the y-axis.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)


Halogen chemistry
over the Dead SeaL. Smoydzin
and R. von Glasow

Fig. 7. Figures in the top row show BrO (contour lines, in pmol/mol) and O₃ (shaded in nmol/mol) mixing ratios over the whole boundary layer for scenarios ENH (a) and LOWBL (b) for the six hours that the model crosses the Dead Sea. Note the logarithmic scaling of the y-axis. Figures in the bottom row show BrO (red dashed lines, in pmol/mol) and O₃ (black solid lines, in nmol/mol) mixing ratios at 2 m altitude for scenarios ENH (c) and LOWBL (d).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)


Halogen chemistry
over the Dead SeaL. Smoydzin
and R. von Glasow

Fig. 8. All plots in this figure refer to scenario STAT. Plot (a) shows the time evolution at 2m altitude of gas phase mixing ratios of HOBr (blue dotted line, pmol/mol), BrO (red dash-dotted line, pmol/mol) and O₃ (black solid line, nmol/mol). Plot (b) shows the vertical distribution of HOBr mixing ratios (shaded, pmol/mol) and the liquid water content of sea salt aerosols (contour lines). Plot (c) shows the vertical distribution of O₃ (shaded, nmol/mol) and BrO mixing ratios (contour, pmol/mol).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)