

Internet Electronic Journal of **Molecular Design**

June 2005, Volume 4, Number 6, Pages 367–380

Editor: Ovidiu Ivanciuc

Proceedings of the Internet Electronic Conference of Molecular Design 2004
IECMD 2004, November 29 – December 12, 2004

Thermochemistry of Organic and Heteroorganic Species. Part XVI. Application of IR Spectra of Unsaturated Aliphatic Molecules to the Thermochemistry of Vinylic and Allylic Free Radicals

Dmitry Ponomarev¹ and Viatcheslav Takhistov²

¹ St. Petersburg Forest Technical Academy, 194021 St. Petersburg, Institutski per.5, Russia

² Center of Ecological Safety RAS, 197042 St. Petersburg, Korpusnaja ul. 18, Russia

Received: November 6, 2004; Revised: February 17, 2005; Accepted: March 15, 2005; Published: June 30, 2005

Citation of the article:

D. Ponomarev and V. Takhistov, Thermochemistry of Organic and Heteroorganic Species. Part XVI. Application of IR Spectra of Unsaturated Aliphatic Molecules to the Thermochemistry of Vinylic and Allylic Free Radicals, *Internet Electron. J. Mol. Des.* **2005**, 4, 367–380, <http://www.biochempress.com>.

Thermochemistry of Organic and Heteroorganic Species. Part XVI. Application of IR Spectra of Unsaturated Aliphatic Molecules to the Thermochemistry of Vinylic and Allylic Free Radicals[#]

Dmitry Ponomarev^{1,*} and Viatcheslav Takhistov²

¹ St. Petersburg Forest Technical Academy, 194021 St. Petersburg, Institutski per.5, Russia

² Center of Ecological Safety RAS, 197042 St. Petersburg, Korpusnaja ul. 18, Russia

Received: November 6, 2004; Revised: February 17, 2005; Accepted: March 15, 2005; Published: June 30, 2005

Internet Electron. J. Mol. Des. 2005, 4 (6), 367–380

Abstract

The application of IR-spectroscopy supplemented by method of isodesmic (formal) reactions which characterize the relative stabilities of free radicals is demonstrated as a useful tool for getting novel data of the enthalpies of formation for certain types of free radicals. In a result, ΔH_f° values for 23 $XC(\bullet)=CH_2$ and 17 $XC(\bullet)=O$ novel radicals were obtained and for 8 more drastically corrected from literature ν_{CH} values in $XCH=CH_2$ and $XCH=O$ molecules, respectively. On the example of $H-CH_2C(X)=O$ molecules it was demonstrated the possibility to estimate the enthalpies of formation of ground state free radicals from ν_{CH} values for excited C–H bonds. Using the results on the latter radicals the ΔH_f° values for 7 novel $\bullet CH_2C(X)=CH_2$ radicals were estimated with the aid of isodesmic reactions. In all these calculations the variable $\Delta \nu_{CH}$ value at $H \rightarrow X$ replacement equivalent to 1 kcal mol⁻¹ in C–H bond dissociation energy was applied: 9 cm⁻¹ for H–CH₂R bond, 14 cm⁻¹ for XC(–H)=O and 11 cm⁻¹ for vinylic C–H bond. The strong destabilization effect in both vinylic type $XC(\bullet)=O$ and $XC(\bullet)=CH_2$ free radicals at $H \rightarrow X$ replacement (X – substituent with lone pair or π -electrons) was found. For $XC(\bullet)=O$ radicals even with such groups as RO or R₂N, traditionally treated as strong electron-donating, destabilization effect at $H \rightarrow X$ replacement was demonstrated. The origin of this effect was suggested in the absence of overlapping of free radical center with lone pair or π -electrons of substituent X. Stabilization (destabilization) of vinylic free radicals is found to be the result of electronegativity/polarizability interplay of a substituent.

Keywords. Thermochemistry; bond dissociation energy; free radicals; IR spectra.

Abbreviations and notations

BDE, bond dissociation energy

PI MS, photoionization mass spectrometry

EN, electronegativity

SE, stabilization energy

PAZ, polarizability

1 INTRODUCTION

Recently the systematic application of known ν_{C-H} frequencies from gas phase IR spectra of R–H molecules to checking, correcting or finding novel values of the enthalpies of formation for

[#] Presented in part at the Internet Electronic Conference of Molecular Design 2004, IECMD 2004.


* Correspondence author; E-mail: juniper@mailbox.alkor.ru.

corresponding R• free radicals has been demonstrated [1–4]. These works followed the fundamental works of McKean with collaborators [5–10] who had shown that $\Delta\nu$ (R–H \rightarrow R'–H) values were correlated with Δ BDE (bond dissociation energies), *i.e.* [BDE(R–H) \rightarrow BDE(R'–H)] values in molecules and consequently with the enthalpies of formation for corresponding R• and R'• free radicals. The problems arising in these correlations are discussed in details in Refs 1–10. For such correlations McKean suggested to take *unique* value in $\Delta\nu$ about 11.5 cm^{-1} for $\text{RH} \rightarrow \text{R}'\text{H}$ replacement equivalent to 1 kcal mol^{-1} for Δ BDE in R–H \rightarrow R'–H molecules [5,6]. Conversely, we showed that the *variable* $\Delta\nu$ equivalents to 1 kcal mol^{-1} in Δ BDE, depending roughly on electronegativity (EN) of C–atom, better fitted such interrelationships [1–4]. Finally, 9 cm^{-1} in alkanes, 11 cm^{-1} at H–C=C double bond, 12 cm^{-1} at H–C \equiv C triple bond and 14 cm^{-1} in RC(–H)=O molecules for $\Delta\nu$ values were adjusted to 1 kcal mol^{-1} in Δ BDE values [1,3,4].

In present work we further explored these correlations to find out novel values of the enthalpies of formation for RC(•)=O, R'C(•)=CH₂ and some •CH₂C(X)=CH₂ free radicals. To find out these values we applied the well known equation suggested by Benson as early as in 1965 for quantitative evaluation of the relative stabilities E_s of free radicals (Eq.1) [11].

$$E_s = \Delta H_f^\circ(\text{R}\bullet) + \Delta H_f^\circ(\text{R}_1\text{H}) - \Delta H_f^\circ(\text{RH}) - \Delta H_f^\circ(\text{R}_1\bullet) \quad (1)$$

For our purpose, estimation of the enthalpies of formation for free radicals, we apply Eq.1 in the form of isodesmic (formal) reaction, Eq. (2), where $Q = E_s$:



The full form of this formal reaction incorporating the ΔH_f° values of all participants of the reaction is given by Eq. (3):

$$\Delta H_f^\circ(\text{R}\bullet) + \Delta H_f^\circ(\text{R}_1\text{H}) \rightarrow \Delta H_f^\circ(\text{RH}) + \Delta H_f^\circ(\text{R}_1\bullet) + Q (= E_s) \quad (3)$$

To obtain the ΔH_f° value of the target $\text{R}_1\bullet$ from that of model free radical $\text{R}\bullet$ we apply Eq. (4) coming out from Eq. (3):

$$\Delta H_f^\circ(\text{R}_1\bullet) = \Delta H_f^\circ(\text{R}\bullet) - \Delta H_f^\circ(\text{RH}) + \Delta H_f^\circ(\text{R}_1\text{H}) - Q \quad (4)$$

The obvious advantage of the use of isodesmic reactions to compare the *relative* stabilities of free radicals is the exclusion of the enthalpies of formation for molecules, *i.e.* free radicals' skeletons and leaving only the enthalpy Q of stabilization (Eq. (3), $Q > 0$) or destabilization (Eq. (3), $Q < 0$) of the target $\text{R}_1\bullet$ compared with the model $\text{R}\bullet$ free radical. Such way of comparison of free radicals relative stabilities is very convenient and easily perceptible. Q value expresses the deviation of the resulted $\Delta H_f^\circ(\text{R}_1)$ value from additive scheme thus making doubtful the elaboration of group contributions for free radicals which still appear in the literature (for example, Refs. [12] and [13]).

2 SUBSTITUTED CARBONYL XC(●)=O FREE RADICALS

We begin presentation of $\Delta v/\text{BDE}$ ($=\Delta\Delta H_f^\circ$) correlations from a representative set of RCH=O molecules (Table 1) (for ΔH_f° values of molecules see Appendix). To extract the enthalpies of formation for free radicals $\text{RC}(\bullet)=\text{O}$ we applied the following calculation scheme based on Eq. (4) and literature $\nu_{\text{CH}}(\text{RCH=O})$ values in gas phase IR spectra.

Table 1. Thermochemical characteristics of $\text{XC}(\bullet)=\text{O}$ free radicals (ν_{CH} values in cm^{-1} – from Refs. [5] and [6], ΔH_f° and Q – in kcal mol^{-1} , derived values – in square brackets).

X-CH=O	ν_{CH}	ΔH_f° (RH)	$\Delta v(\text{H}\rightarrow\text{X})$	Q H \rightarrow X	ΔH_f° XC(●)O calc.		ΔH_f° [XC(●)O] lit.
					from Δv	final	
$\text{Me}_2\text{C=CH}$	2768 [15]	[-35.5]	-45	3.21	-2.7	-2.5	
Me	2770	-39.73	-43	3.07	-6.8	-6.8	-2.4 [12], -4.5 [13], -5.4 [14,19]
Ph	2770	-8.8	-43	3.07	24.1	24	26.1 [13, 14], 27.8 [12]
H	2813	-26.0	0.0	0.0	10.0		9.96 [12], 10.0 [13]
MeS	2838 [16]	[-33.5]	25	-1.8	4.3	4.5	
HS	[2844]	-30	31	-2.2	8.2	8.0	
CF_3	2844	-188.3	31	-2.2	-151.1	-151	-128.4 [12]
Me_2N	[2844]	-45.8	31	-2.2	-6.0	-6.0	
Br_3C	2845	[-1]	32	-2.3	-38.0	-38	
MeNH	[2847]	[-44.5]	34	-2.4	-5.5	-5.5	
CHO	2850	-50.7	37	-2.6	12.1	12	
H_2N	2852	[-43.5]	39	-2.8	-4.7	-4.5	
Cl_3C	2856	[-38.1]	43	-3.1	1.0	1	
$\text{HC}\equiv\text{C}$	2858	24.5	45	-3.2	64.4	64.5	
$\text{BrC}\equiv\text{C}$	2858 [16]	[28]	45	-3.2	67.9	68	
$\text{ClC}\equiv\text{C}$	2860 [16]	[15.5]	47	-3.36	55.4	55.5	
Br	2912.5 [17]	[-32.5]	99.5	-7.1	10.6	10.5	
I	2930 [17]	-17	117	-8.36	27.8	28	
MeO	2930	-85.0	117	-8.36	-40.2	-40	-40.4 [19]
Cl	2934 [17]	[-45]	121	-8.64	-0.5	-0.5	-2.8 [13], -5.2 [13]
HO	2943	-90.5	130	-9.3	-45.2	-45	-46.5 [12], -47.3 [20], -53.3 [13]
HCOO (two rotamers)	2948 [18] 2964 [18]	[-110]	135 151	-9.6 -10.8	-63.9 -62.7	-64	
F	2981.2 [17]	[-93]	168.2	-12.0	-44.4	-44.5	-43.0 [13]

Formaldehyde H-CH=O was taken as a model compound. The ΔH_f° $\text{HC}(\bullet)=\text{O}$ $10.0 \text{ kcal mol}^{-1}$ is well established [12,13]. We give an example of calculation for ΔH_f° $\text{MeC}(\bullet)=\text{O}$ free radical from ν_{CH} MeCH=O molecule. $\Delta H_f^\circ[\text{MeC}(\bullet)=\text{O}] = \Delta H_f^\circ[\text{HC}(\bullet)=\text{O}] - \Delta H_f^\circ(\text{H}_2\text{C=O}) + \Delta H_f^\circ(\text{MeCH=O}) - Q = 10.0 - (-26.0) + (-39.73) - Q$ (Eq. (4), $R = \text{H, Me}$). We gain $Q = \{\nu_{\text{CH}}(\text{H-CH=O}) - \nu_{\text{CH}}[\text{CH}_3\text{C}(\text{-H})=\text{O}]\}$: $14 = (2813 - 2770)$: $14 = 43:14 = 3.07 \text{ kcal mol}^{-1}$. Each 14 cm^{-1} in Δv are equivalent to 1 kcal mol^{-1} in $\Delta \Delta H_f^\circ$ values. The positive sign of Q 3.07 signifies that acetyl $\text{CH}_3\text{C}(\bullet)=\text{O}$ is by $3.07 \text{ kcal mol}^{-1}$ more stable than formyl $\text{HC}(\bullet)=\text{O}$ free radical or, in other words $\text{H} \rightarrow \text{CH}_3$ replacement reveals stabilization of free radical center by $3.07 \text{ kcal mol}^{-1}$. Thus, in general, we use the additive scheme as a primary step and then introduce the correction term Q reflecting quantitatively the electron-donating ($Q > 0$) or electron-withdrawing ($Q < 0$) properties of R -substituent compared with H -atom in formyl $\text{H-C}(\bullet)=\text{O}$ radical. ν_{CH} values for some

molecules were extrapolated using IR data for structurally close compounds, like HSCH=O from data on MeSCH=O or MeNHCH=O and $\text{Me}_2\text{NCH=O}$ from data on $\text{H}_2\text{NCH=O}$ molecule keeping in mind that $\text{H} \rightarrow \text{Me}$ replacement decreases the ν_{CH} value which parallels the decrease of the BDE C–H in molecules.

Analysis of data collected in Table 1 revealed some unexpected features. The first of them is the minor (if any) role of resonance (conjugation) interaction of lone pairs of heteroatom or π -system (Ph, $\text{XC}\equiv\text{C}$) with unshared electron at free radical center. Such substituents as halogens and especially RO- , RS- , R_2N ($\text{R} = \text{H}$, alkyl)-groups, being well known as stabilizing free radical center compared with Me-group, reveal, contrary, the destabilizing effect in the $\text{RC}(\bullet)=\text{O}$ series. Let us compare the Q values (Table 2) for two isodesmic reactions Eqs. (5) and (6). The data for $\text{RCH}_2\cdot$ radicals are taken from Ref. [14] (here ΔH_f° for $\text{CH}_3\cdot$ radical is taken $35.1 \text{ kcal mol}^{-1}$) and for $\text{RC}(\bullet)=\text{O}$ radicals from present work. The isodesmic reactions in Table 2 are given in a brief form like $\text{CH}_3\bullet \rightarrow \text{CH}_2\text{NMe}_2 + Q$, instead of a full form (Eqs. (5) and (6)) or $\text{CH}_3\bullet + \text{CH}_3\text{NMe}_2 \rightarrow \text{CH}_4 + \bullet\text{CH}_2\text{NMe}_2 + Q_1$.





Table 2. Stabilization energies Q for $\text{CH}_3\bullet \rightarrow \text{XCH}_2\bullet$ (Q_1) and $\text{HC}(\bullet)=\text{O} \rightarrow \text{XC}(\bullet)=\text{O}$ (Q_2) (ΔH_f° $\text{R}\bullet$ and Q – in kcal mol^{-1}); ΔH_f° ($\bullet\text{CH}_2\text{X}$) from Ref. [4], ΔH_f° $\text{XC}(\bullet)=\text{O}$ – this work (Table 1).

$\bullet\text{CH}_2\text{X}$		$\Delta H_f^\circ(\bullet\text{CH}_2\text{X})$	Q_1	$\text{XC}(\bullet)=\text{O}$		Q_2
$\text{CH}_3\bullet$	$\rightarrow \bullet\text{CH}_2\text{NMe}_2$	25.8	21.4	$\text{HC}(\bullet)=\text{O}$	$\rightarrow \text{Me}_2\text{NC}(\bullet)=\text{O}$	-2.2
	$\rightarrow \bullet\text{CH}_2\text{NHMe}$	30.9	17.5		$\rightarrow \text{MeNHC}(\bullet)=\text{O}$	-2.4
	$\rightarrow \bullet\text{CH}_2\text{NH}_2$	34.9	12.5		$\rightarrow \text{H}_2\text{NC}(\bullet)=\text{O}$	-2.8
	$\rightarrow \bullet\text{CH}_2\text{OMe}$	-5.0	13.9		$\rightarrow \text{MeOC}(\bullet)=\text{O}$	-8.4
	$\rightarrow \bullet\text{CH}_2\text{OH}$	-3.2	7.9		$\rightarrow \text{HOC}(\bullet)=\text{O}$	-9.3
	$\rightarrow \bullet\text{CH}_2\text{F}$	-7.5	6.1		$\rightarrow \text{FC}(\bullet)=\text{O}$	-12
	$\rightarrow \bullet\text{CH}_2\text{Cl}$	28.5	4.8		$\rightarrow \text{ClC}(\bullet)=\text{O}$	-8.6
	$\rightarrow \bullet\text{CH}_2\text{Br}$	41	2.3		$\rightarrow \text{BrC}(\bullet)=\text{O}$	-7.0
	$\rightarrow \bullet\text{CH}_2\text{I}$	54.5	1.3		$\rightarrow \text{IC}(\bullet)=\text{O}$	-8.4
	$\rightarrow \bullet\text{CH}_2\text{Me}$	27	5.8		$\rightarrow \text{MeC}(\bullet)=\text{O}$	3.1
	$\rightarrow \bullet\text{CH}_2\text{Ph}$	47	17.9		$\rightarrow \text{PhC}(\bullet)=\text{O}$	3.1

All substituents collected in Table 2 stabilize the $\text{CH}_3\cdot$ free radical ($Q_1 > 0$) while in substituted formyl $\text{XC}(\bullet)=\text{O}$ free radicals only Me- and Ph-groups reveal a small stabilizing effect ($Q_2 > 0$). Even such traditionally very strong electron donating groups as $\text{H}_2\text{N-}$, $-\text{OMe}$, MeNH- , $\text{Me}_2\text{N-}$ destabilize formyl radicals ($Q_2 < 0$). All that signifies that in substituted formyl free radicals there is no overlapping between heteroatoms' lone pair and unshared electron of free radical center or, in other words, the conjugation between them is absent. If in the series of halogenated $\text{XCH}_2\bullet$ radicals the resonance stabilization decreases in the row $\text{F} > \text{Cl} > \text{Br} > \text{I}$ in consensus with their R^+ values – 0.52 (F), -0.31 (Cl), -0.30 (Br) and -0.28 (I) [21] the observed effect in halogenated formyl radicals is the opposite one just following inductive σ_{I} [21] or electronegativity values [3]. Thus, it may

appear that substituted formyl radicals may become the models of “pure” inductive effect without involvement of resonance effect.

Another interesting observation is that electron-donating effects of Me- and Ph-groups are identical in their relation to stabilization of formyl free radical while, as expected, the latter demonstrated the much higher substitution effect on $XCH_2\bullet$ radical when compared with $X = Me$. This again can only be treated as the evidence of absence of resonance stabilization of the free radical center by π -system while minor stabilization still takes place owing to polarizability (PAZ) effect of Ph- which is expected to be larger than with Me-group (compare ENs values 4.8 and 5.4 units for Me- and Ph-groups, respectively [3]).

One more specific feature of the $XC(\bullet)=O$ system is contribution of PAZ effect of X-substituents. Its role in stabilization of free radicals on the example of $XC\equiv C\bullet$ species is described in detail in Ref.3. It is expected that the larger is the size of an atom or of a group the larger is the PAZ effect inspired by electron-deficient free radical center. The behavior of $XC(\bullet)=O$ radicals (Tables 1 and 2) parallels that observed with $XC\equiv C\bullet$ free radicals. For the latter only a few “hard” (of small size and low PAZ) substituents F, HO, Cl, and $-N=C$ destabilize $HC\equiv C\bullet$ radical in the framework of isodesmic reaction $HC\equiv C\bullet + XC\equiv CH \rightarrow HC\equiv CH + XC\equiv C\bullet + Q$ ($Q < 0$). All others, including such traditionally treated as strong electron-withdrawing groups as $-C\equiv N$, CF_3 , $COOH$ and even SF_5 stabilize $XC\equiv C\bullet$ radical ($Q > 0$) since ν_{CH} values for $XC\equiv C-H$ molecules with these groups are *lower* than ν_{CH} for unsubstituted $HC\equiv CH$ molecule [1,3].

Since $HC(\bullet)=O$ radical is much more stable [by $32.5 \text{ kcal mol}^{-1}(=Q^\circ)$!] in the framework of isodesmic reaction $HC\equiv C\bullet + CH_2=O \rightarrow HC\equiv CH + HC(\bullet)=O + Q^\circ$ many other X-atoms and groups enter the list of destabilizing substituents. This comes out from our observations and found regularities that some substituents stabilize unstable, like $Me\bullet$ radicals while destabilize the more stable radicals like $Me_2CH\bullet$, that is $Q > 0$ for $Me\bullet + CH_3X \rightarrow MeH + \bullet CH_2X$ and $Q < 0$ for $Me_2CH\bullet + Me_2CHX \rightarrow Me_2CH_2 + Me_2C(\bullet)X$ isodesmic reaction. In the absence of possibility to bring about their resonance effect in $XC(\bullet)=O$ system such substituents as halogens, $RO-$, $RS-$, $RC\equiv C-$ and even R_2N- exhibit electron-withdrawing (destabilizing) effects. Of course, one should not expect to “measure” the PAZ effect directly from IR data since the interplay between EN and PAZ of a group might appear to be very complex [3]. For example, such rather diverse groups as CF_3 , CBr_3 , HS , Me_2N or Cl_3C and $HC\equiv C$ give very close ν_{CH} values. Another complicating factor even in case of structurally close groups like CX_3 (F, Cl, Br) is the possible difference in their geometrics in relation to C-H bond in $RCH=O$ molecule.

3 1-X-SUBSTITUTED VINYL FREE RADICALS

For $XCH=CH_2$ molecules and $XC(\bullet)CH=CH_2$ free radicals the effects similar to those for $XCH=O$ molecules and $XC(\bullet)=O$ radicals, respectively, are expected. The set of compounds

XCH=CH₂ for which the data on IR spectra are available is more narrow owing to chemical instability of many of them like XCH=CH₂ with X=OH, SH, NH₂. Still, the general tendencies in structure/enthalpy of formation are similar in both classes of compounds. The most important and common tendency remains untouched, this is the minor (if any) participation of resonance (conjugation) effect in interaction of X–substituent, possessing either lone pairs or π–electrons, with free radical center and the important role of polarizability effect (Table 3). The best example to demonstrate the role of PAZ effect are the ν_{CH} values for CH₂=CHEH₂ molecules, where E is the element of V (15) group N–Bi (Table 3).

Table 3. Thermochemistry of XC(•)=CH₂ free radicals [ν_{CH} and Δν – in cm⁻¹ refer to vinylic C–H bond), ΔH_f^o and Q values – in kcal mol⁻¹, derived values – in square brackets; for ΔH_f^o of parent molecules see Appendix].

X–CH=CH ₂	ν _{CH}	ΔH _f ^o	Δν(H→X)	Q	ΔH _f ^o XC(•)=CH ₂		literature value
		XCH=CH ₂			H→X	from Δν	
H ₂ Bi	[2940]	[75]	-120	11.0	121.5	121.5	
H ₂ Sb	2949.3[22]	[57.7]	-110.7	10.1	105.14	105	
H ₂ As	2959[22]	[35.5]	-101	9.2	83.8	84	
Me ₂ N	[2975]	[13]	-85	7.7	62.9	63	
H ₂ P	[2980]	[19.8]	-77	7	70.5	70.5	
MeNH	2990[23]	[14.2]	-70	6.4	65.3	65.5	
ClCO (two rotamers)	2990[24] 2998	[-32]	-70	6.4	19.3	19.5	
CH ₃ CO	[2997]	[-28]	-63	5.7	23.6	23.5	27 [20]
Et	2998[16]	0.02	-62	5.6	51.8	52	
CH ₂ =CH	3000[25]	26.3	-60	5.5	78.2	78	
Ph	[3000]	35.35	-60	5.5	87.2	87	74 [20]
Me	3001[5]	4.88	-59	5.4	56.7	56.5	
H ₂ N	[3010]	13.5	-50	4.5	66.3	66.5	57.5 [20]
HC≡C	3012[5]	69.8	-48	4.4	122.7	122.5	108 [29]
ClCH ₂	3021[16]	0.5	-39	3.5	54.2	54	
BrCH ₂	3021[16]	10.8	-39	3.5	64.7	64.5	
FCH ₂	3026[16]	-35.5	-34	3.1	25.1	25	
MeS	3044[25]	[16]	-16	1.5	72.0	72	
Cl ₂ C=C(Cl)	3047[27]	11.5	-13	1.2	67.7	67.5	
H	3060[5]	12.55	0.0	0.0	70.0	70	
MeO	3065[5]	[-24.5]	5.0	-0.6	33.5	33.5	
HS	[3070]	19.5	10	-0.9	78.0	78	
CF ₃	[3070]	-151	10	-0.9	-92.5	-92.5	
Cl	3071[28]	5.2	11	-0.9	63.3	63.5	
N≡C	3078[5]	43.16	18	-1.6	102.2	102	
F	3080[5]	-35	20	-1.8	24.4	24.5	
HO	[3082]	-30.4	22	-2.0	28.9	29	17.5 [20]
Br	3085[5]	18	25	-2.3	77.8	78	
I	[3087]	32	27	-3.5	91.9	92	

Vinyl •CH=CH₂ radical is by 21.55 kcal mol⁻¹ less stable than •CH=O free radical in the framework of isodesmic reaction •CH=O + CH₂=CH₂ → CH₂=O + •CH=CH₂ + Q [Q = ΔH_f^o(•CH=O) + ΔH_f^o(CH₂=CH₂) – ΔH_f^o(CH₂=O) – ΔH_f^o(•CH=CH₂) = 10.0 + 12.55 – (-26.0) – 70 = -21.55 kcal mol⁻¹(destabilization)]. Therefore, many of the substituents which destabilize the XC(•)=O free radicals now become stabilizing ones in XC(•)=CH₂ radicals in the framework of isodesmic reactions. In other words, the less stable vinyl •CH=CH₂ free radical extracts

stabilization from those substituents which in the system of more stable formyl $\bullet\text{C}(\text{H})=\text{O}$ radical were electron-withdrawing, *i.e.* destabilizing the latter ones. This fact stands in line with our previous observations (see above and Refs. [1–4,14]). Among those which now destabilize vinyl radicals we find all halogens, HO, MeO, HS and NC–groups. Using the known experimental data we – by interpolation – ascribed ν_{CH} values for some $\text{XCH}=\text{CH}_2$ molecules for which such data are not available. IR spectra of substituted ethylenes $\text{XCH}=\text{CH}_2$ revealed certain specificity in interaction of halogens with π –double bond. The measured ν_{CH} values in the series of halogenated ethylenes $\text{XCH}=\text{CH}_2$ (F, Cl, Br, I) range only for 7 cm^{-1} (compare with the range of 70 cm^{-1} for $\text{XCH}=\text{O}$ molecules – see above) (Table 3). Thus, it is hardly possible to make any definite conclusions about the role of inductive effect (or EN), polarizability and geometrical fine structure for a particular halogen.

Summarizing the data on thermochemistry of free radicals of vinylic type we now realize that our earlier estimations of the enthalpies of formation of some $\text{XC}(\bullet)=\text{CH}_2$ radicals ($\text{X} = \text{Ph}, \text{H}_2\text{N}, \text{HO}, \text{HC}\equiv\text{C}$) [20,29] (Table 3) were based on erroneous assumptions that either of free radicals saturated and vinylic ones follow similar structure/stability interrelationships. Similarly, the quoted value for ΔH_f° $\text{CF}_3\text{C}(\bullet)=\text{O}$ radical -128.4 [12] compared with the value $-151\text{ kcal mol}^{-1}$ (present work) would give too high destabilization effect of $-23.6\text{ kcal mol}^{-1}$ for $\text{HC}(\bullet)=\text{O} \rightarrow \text{CF}_3\text{C}(\bullet)=\text{O}$ replacement (compare with $-2.2\text{ kcal mol}^{-1}$ – Table 1). This would lead to unprecedented value for ν_{CH} in $\text{CF}_3\text{CH}=\text{O}$ molecule being much more higher than with F–substituent: $\nu_{\text{CH}}(\text{CF}_3\text{CH}=\text{O}) = \nu_{\text{CH}}(\text{CH}_2=\text{O})(2813) + 23.6 \times 14$ (see above) = 3143.4 cm^{-1} (!).

Table 4. Thermochemistry of $\bullet\text{CH}_2\text{C}(\text{X})=\text{O}$ free radicals (ν_{CH} from Ref. 6 – in cm^{-1}); ΔH_f° and Q values – in kcal mol^{-1} , molecules' ΔH_f° – from Ref. 31; derived values – in square brackets).


Molecule RH	ΔH_f° RH	$\nu_{\text{exp.}}$ CH _{as}	$\nu_{\text{calc.}}$ CH _{as}	Q_0 H→X	ΔH_f° $\bullet\text{CH}_2\text{COX}$	$\nu_{\text{exp.}}$ CH _{sym}	Q_1 H→X	ΔH_f° $\bullet\text{CH}_2\text{COX}$
$\text{CH}_3\text{CH}=\text{O}$	-39.73	2945	2900.5	0.0	3.0^a	3002	0.0	9.3
CH_3COCH_3	-51.94	2946	2893.5	-0.1	-9.1^a -10^b	3004	-0.2	-2.7
CH_3COF	-105.66	2980.1	2935.6	-3.9	-59^a -60.5^b	3025.4	-2.6	-54
CH_3COCl	-58.3	2978.5	2934	-3.8	-11.5^a -14^b	3004.0	-0.2	-8.8
CH_3COBr	-45.5	2978.2	2933.7	-3.7	1.0^a -1.2^b	2994.9	0.8	2.7
CH_3COI	-30.2	[2978]	2933.5	-3.7	16^a	2983.3	2.1	16.8
CH_3COOMe	-98.45	2972.7	2928.2	-3.0	-52.5^a -52.4^b	3012	-1.1	-48.3
CH_3COCN	[-3]	2957.5	2913	-1.44	40^a	3010	-0.9	46

^a this work; ^b from Ref. 30

4 ALLYLIC TYPE $\bullet\text{CH}_2\text{C}(\text{X})=\text{O}$ FREE RADICALS

According to Ref. [6] CH_3CHO molecules exhibit in its IR spectrum two bands with ν_{CH} frequencies in CH_3 –group 2945 (CH_{as}) and 3002 (CH_{sym}). The lower ν_{CH} (antisymmetric or *cis*)

corresponds to free radical's conformer $\bullet\text{CH}_2\text{CH}=\text{O}$ in which free radical's center practically does not interact with the double $\text{C}=\text{O}$ bond. The higher ν_{CH} (symmetric or *trans*) belongs to the conformer in which free radical's electron is actively interacting with this double bond performing the destabilizing effect since here CO -group reveals its electron-withdrawing properties. However, even the lower ν_{CH} corresponds to $\bullet\text{CH}_2\text{CH}=\text{O}$ radical being in "excited" rather in its ground state. According to McKean [6] BDEs $\text{H}-\text{CH}_2\text{CH}=\text{O}$ are 99.4 and 92 kcal mol⁻¹, coming, respectively, from ν_{CH} 2945 cm⁻¹ value and from kinetic data (ground state). Thus, all low ν_{CH} values in $\text{H}-\text{CH}_2\text{C}(\text{X})=\text{O}$ molecules (Table 4) correspond to formation of $\bullet\text{CH}_2\text{C}(\text{X})=\text{O}$ radical in "excited" state with *cis*-position of unshared electron to CO -group. We estimated the enthalpies of formation of series of $\bullet\text{CH}_2\text{C}(\text{X})=\text{O}$ radicals applying the known ΔH_f° value for unsubstituted $\bullet\text{CH}_2\text{CH}=\text{O}$ 3 kcal mol⁻¹ (kinetic and mass spectrometric studies [14]) by estimating ν_{CH} $\text{H}-\text{CH}_2\text{C}(\text{X})=\text{O}$ in the ground state with the help of following equations (ΔH_f° and Q – in kcal mol⁻¹)



$$\nu_{\text{CH}}(\text{calc.}) (\text{H}-\text{CH}_2\text{CH}=\text{O}) = \nu_{\text{CH}}(\text{CH}_3-\text{H}) - Q_1(\text{CH}_3\bullet \rightarrow \bullet\text{CH}_2\text{CH}=\text{O}) \times 9 = \quad (8)$$

$$= 2992 - [10.17 \times 9 = 91.5] = 2900.5$$

$$\nu_{\text{CH}}(\text{calc.} \rightarrow \text{exp.}) = 2900.5 - 2945 = -44.5 \text{ cm}^{-1} \quad (9)$$

where 9 cm⁻¹ is the equivalent for 1 kcal mol⁻¹ in BDEs in $\text{H}-\text{CH}_2$ -group. Then we calculated ν_{CH} values for the ground states of all other $\text{H}-\text{CH}_2\text{C}(\text{X})=\text{O}$ molecules using ν_{CH} (exp.) and introducing the common correction term -44.5 cm⁻¹ [Eq. (9)] as for unsubstituted $\text{H}-\text{CH}_2\text{CH}=\text{O}$ molecule [Eqs. (7)–(9)]. For example:

$$\nu_{\text{CH}}(\text{calc.}) [\text{H}-\text{CH}_2\text{C}(\text{F})=\text{O}] = \nu_{\text{CH}}(\text{exp.}) [\text{H}-\text{CH}_2\text{C}(\text{F})=\text{O}] (2980.1) - 44.5 = 2935.6 \text{ cm}^{-1} \quad (10)$$

From these ν_{CH} (calc.) values for ground state $\bullet\text{CH}_2\text{C}(\text{X})=\text{O}$ radicals with *cis*-position of unshared electron and CO -group (Table 4) we then calculated the Q values for isodesmic reactions $\bullet\text{CH}_2\text{CH}=\text{O} \rightarrow \bullet\text{CH}_2\text{C}(\text{X})=\text{O}$ (brief form). For example,

$$\Delta\text{H}_f^\circ [\bullet\text{CH}_2\text{C}(\text{F})=\text{O}] = \Delta\text{H}_f^\circ [\bullet\text{CH}_2\text{CH}=\text{O}] - \Delta\text{H}_f^\circ [\text{CH}_3\text{CH}=\text{O}] + \Delta\text{H}_f^\circ [\text{CH}_3\text{C}(\text{F})=\text{O}] - \quad (11)$$

$$Q\{\nu_{\text{CH}}(\text{calc.}) \text{H}-\text{CH}_2\text{C}(\text{F})=\text{O} : 9\} = 3.0 - (-39.73) - 105.66 [31] - [(2900.5 - 2935.6) : 9] = -59.03 \sim 59 \text{ kcal mol}^{-1}$$

When similar calculation was performed with $\nu_{\text{CH}}(\text{sym.})$ values (Table 4) which were responsible for interaction of CO -group and unshared electron being in *trans*-position to each other such interaction appeared to be rather specific. Halogens showed the yet higher contribution of their PAZ effects to such an extent that only F-atom (very low PAZ) destabilized $\bullet\text{CH}_2\text{C}(\text{F})=\text{O}$ radical compared with unsubstituted $\bullet\text{CH}_2\text{CH}=\text{O}$ free radical ($Q = -2.6$), $\bullet\text{CH}_2\text{C}(\text{Cl})=\text{O}$ demonstrated similar stability with the latter ($Q = -0.2$) while Br- and then I-atoms showed the yet stronger stabilization effects ($Q > 0$). Consequently, for Br- and, especially, for I-substituents both conformers (with either *cis*- or *trans*-position of $\text{C}(\text{X})=\text{O}$ group to unshared electron) become close

in their stabilities: compare ΔH_f° values for *cis*- and *trans*-radicals 1.0 and 2.7 (Br) or 16 and 16.8 kcal mol⁻¹ (X = I). We stress that either $\nu_{\text{CH}}(\text{as})$ or $\nu_{\text{CH}}(\text{sym})$ values correspond to free radicals being in “excited” states.

Owing to essentially different interaction in *cis*- and *trans*-conformers (compare Q_0 and Q_1 – Table 4) it is hardly possible to calculate the ΔH_f° values of the *trans*-substituted radicals in their ground states as it was performed for *cis*-conformers (see above). All these observations coming from data on IR spectra enlighten the possible complexity in interpretation of experimental results on determination of BDEs in H–CH₂C(X)=O [and H–CH₂C(X)=CH₂ – see later] molecules by methods of chemical kinetics which hardly take in account the described effects coming from IR data.


It is important to mention that experimental ΔH_f^0 values for •CH₂COCH₃ (–10) and •CH₂COOCH₃ (–52.4 kcal mol⁻¹) were obtained by photoionization mass spectrometry (PI MS) [30] (Table 4) and are very close to data gained in present work from IR data thus supporting our methodology of calculating ΔH_f° •CH₂C(X)=O free radical from ν_{CH} values related to “excited” C–H bonds. The literature values for •CH₂C(X)=O radicals (F, Cl, Br) were calculated by additive scheme from experimental data on •CH₂C(X)=O radicals (Me, OMe) [30]. We think that ΔH_f° •CH₂COCH₃ –10 kcal mol⁻¹ gained from PI MS [30] is more reasonable than that obtained from IR spectrum of acetone (Table 4) –9.1 [6] or from chemical kinetics –5.7 [19] kcal mol⁻¹. What is strange that both ν_{CH} values for acetone 2946 and 3004 cm⁻¹ are higher than for CH₃CHO molecule 2945 and 3002 cm⁻¹, respectively (Table 4). H → Me replacement might stabilize free radical and thus it is expected that ν_{CH} values should be lower for acetone H–CH₂COCH₃ than for acetaldehyde H–CH₂CHO like it was observed for many other systems at H → Me replacement (Refs. [1–4] and Tables 1–3). The possible source of this discrepancy is involvement of the enol form of acetone CH₂=C(OH)CH₃ molecule. Another possible reason is that namely acetone (for unknown reason) deviates from $\nu_{\text{CH}}(\text{as})$ (exp. → calc.) tendencies which are followed by all other CH₃C(X)=O molecules. Still the data on •CH₂COCH₃ radical coming from IR spectra and from PI MS differ by only 1 kcal mol⁻¹.

The discrepancy between PI MS and kinetic data –10 [30] and –5.7 [19] kcal mol⁻¹ respectively, can now be explained by involvement in kinetic experiment (bimolecular reaction X• + H–CH₂COCH₃ → HX + •CH₂COCH₃) of the stronger C–H bond of *trans* conformer (see above).


5 ALLYLIC TYPE •CH₂C(X)=CH₂ FREE RADICALS

To derive the enthalpies of formation for this type of free radicals we followed the tendencies in structure/ ΔH_f° for •CH₂C(X)=O radicals (see above). In the framework of isodesmic reactions •CH₂CH=CH₂ is by 8.4 kcal mol⁻¹ more stable than •CH₂CH=O free radical [Eq. (12)] (ΔH_f° and Q

– in kcal mol⁻¹).



Since all X–substituents (with exception of Me) destabilize $\bullet\text{CH}_2\text{C}(\text{X})=\text{O}$ free radicals at H \rightarrow X replacement, *i.e.* $Q < 0$ for $\bullet\text{CH}_2\text{CH}=\text{O} \rightarrow \bullet\text{CH}_2\text{C}(\text{X})=\text{O}$ isodesmic reaction (brief form) (Table 4) the same X–substituents will even more destabilize the more stable [Eq. (12)] allyl $\bullet\text{CH}_2\text{C}(\text{X})=\text{CH}_2$ free radicals at H \rightarrow X replacement in β –position to free radical center (see above). We can estimate approximately the Q value for, say, $\bullet\text{CH}_2\text{CH}=\text{CH}_2 + \text{CH}_3\text{C}(\text{F})=\text{CH}_2 \rightarrow \text{CH}_3\text{CH}=\text{CH}_2 + \bullet\text{CH}_2\text{C}(\text{F})=\text{CH}_2 + \text{Q}$ isodesmic reaction [Eq. (13)] applying the known data for less stable $\text{FCH}_2\text{CH}_2\bullet$ [32] and $\bullet\text{CH}_2\text{C}(\text{F})=\text{O}$ (this work) free radicals [Eq. (14)] [the values at arrows signify Q values (in kcal mol⁻¹) for corresponding isodesmic reaction like $\bullet\text{CH}_2\text{CH}=\text{CH}_2 \rightarrow \bullet\text{CH}_2\text{C}(\text{F})=\text{CH}_2$ (brief form) while Eq. (13) is the full form of this reaction].



Here we applied the earlier found observation that introduction of an electron–withdrawing substituent to yet more stable free radical performs yet higher destabilizing effect (see above). The final values of the enthalpies of formation for $\bullet\text{CH}_2\text{C}(\text{X})=\text{CH}_2$ free radicals are collected in Table 5.

Table 5. Thermochemical characteristics of $\bullet\text{CH}_2\text{C}(\text{X})=\text{CH}_2$ free radicals (ΔH_f° and Q – in kcal mol⁻¹; derived values – in square brackets)

X	ΔH_f°	Q(H→X)	$\Delta H_f^\circ(\text{R}\bullet)$
H	4.88	0.0	39.1[14]
F	[-46.5]	-5.7	-7.5
Cl	[-5.0]	-5.55	33.5
Br	[8.5]	-5.35	47.5
I	[23.5]	-5.35	61.5
CN	[32]	-2.45	70
MeO	[-35]	-4.15	2.5
HO	[-41.5]	-5.55	-4.5

6 CONCLUSIONS

In present work we further demonstrated the application of IR spectroscopy as a useful tool for getting novel data of the enthalpies of formation for certain types of free radicals if IR data are supplemented by series of isodesmic (formal) reactions. In a result, ΔH_f° values for 23 $\text{XC}(\bullet)=\text{CH}_2$ and 17 $\text{XC}(\bullet)=\text{O}$ novel radicals were obtained and for 8 more were drastically corrected using literature ν_{CH} values in $\text{XCH}=\text{O}$ and $\text{XCH}=\text{CH}_2$ molecules. For the first time, on the example of H–

CH₂C(X)=O molecules the IR ν_{CH} values for excited C–H bonds were applied to calculate the ΔH_f° values in the ground state. Using the results on the latter radicals the enthalpies of formation for 7 novel $\bullet\text{CH}_2\text{C(X)=CH}_2$ radicals were estimated with the aid of isodesmic reactions.

The strong destabilization effect in both types XC(\bullet)=O and XC(\bullet)=CH₂ free radicals at H → X replacement (X – substituent with lone pair or π -electrons) was demonstrated. The origin of this effect was explained by the absence of overlapping of free radical center with lone pair or π -electrons of X-substituent.

Appendix 1


Most of the known values for the enthalpies of formation for parent molecules are taken from Ref. [31]. The values for ΔH_f° of H₂NCH=CH₂ (13.5), HSCH=CH₂ (19.5), BrCH=CH₂ (18), HOCH=CH₂ (–30.4), ClCH=CH₂ (5.2), CF₃CH=CH₂ (–151), FCH=CH₂ (–35), FCH₂CH=CH₂ (–35.5), ClCH₂CH=CH₂ (0.5) we extracted from Ref. [32]. ΔH_f° CF₃CHO –188 kcal mol^{–1} was earlier obtained in Ref. [33]. ΔH_f° HSCHO –30 kcal mol^{–1} was earlier estimated in Ref. [20] and hence $\Delta H_f^\circ[\text{MeSCH}_2] = \Delta H_f^\circ[\text{HSCHO}] - \Delta H_f^\circ[\text{MeSH}] (-5.5) + \Delta H_f^\circ[\text{MeSMe}] (-9.0) = -33.5$ kcal mol^{–1}. The data for MeNHCHO (or MeNHCH=CH₂) and Me₂NCHO (or Me₂NCH=CH₂) were estimated by additive scheme from ΔH_f° values for H₂NCHO (or H₂NCH=CH₂), H₂NMe and Me₂NH molecules [31]. The enthalpies of formation for HCOX (X = F–I) were calculated by additive scheme from known ΔH_f° values of their homologues CH₃COX molecules ([31] and Table 4). For example, $\Delta H_f^\circ[\text{HCOF}] = \Delta H_f^\circ[\text{HCOOH}] (-90.5) + \Delta H_f^\circ[\text{CH}_3\text{COF}] (-105.7) - \Delta H_f^\circ[\text{CH}_3\text{COOH}] (-103.44) \approx -93$ kcal mol^{–1}. Similarly, $\Delta H_f^\circ[\text{OCH-O-CHO}] = \Delta H_f^\circ[\text{CH}_3\text{CO-O-COCH}_3] (-136.83) - 2\Delta H_f^\circ[\text{CH}_3\text{COOCH}_3] (-98.45) + 2\Delta H_f^\circ[\text{HCOOCH}_3] (-85.0) \approx -110$ kcal mol^{–1}. ΔH_f° values for HC≡CCH=O (24.5) and HC≡CCH=CH₂ (69.8) are taken from Ref. [3]. The heats of formation for ClC≡CCHO and BrC≡CCHO molecules (Table 1) were estimated by additive scheme using ΔH_f° values for ClC≡CH (45.5), BrC≡CH (58) and HC≡CCHO (24.5) [3]. Since what kind of Hal/CHO interaction (repulsive, attractive or zero) might appear in disubstituted acetylene is unknown (for details see Ref. [3]) we just applied the additive scheme without introducing any correction term.

$\Delta H_f^\circ(\text{CH}_3\text{COCN})$ was estimated by macroincremental scheme [20]: $\Delta H_f^\circ[\text{CH}_3\text{COCN}] = \Delta H_f^\circ[\text{CH}_3\text{COCF}_3] (-200.7)[20,33] - \Delta H_f^\circ[\text{CH}_3\text{CF}_3] (-179.5)[4] + \Delta H_f^\circ[\text{CH}_3\text{CN}] (18[4]) = -3.2 \approx -3$ kcal mol^{–1}. The CH₃COCF₃ molecule was selected as a model compound due to close EN values for CF₃ (7.7) and NC-groups (7.9) [3]. Probably, because of expected larger steric CO/CF₃ interaction compared with that for CO/CN pair ΔH_f° CH₃COCN might appear to be even more negative. The literature experimental value is 6.2±6.2 kcal mol^{–1} [34]. The enthalpy of formation for Me₂C=CHCHO molecule was calculated using additive (macroincremental) scheme introducing the small correction term SE: $\Delta H_f^\circ[\text{Me}_2\text{C=CHCHO}] = \Delta H_f^\circ[\text{CH}_2=\text{CHCHO}] (-16.5) + \Delta H_f^\circ[\text{Me}_2\text{C=CH}_2] (-4.0) - \Delta H_f^\circ[\text{CH}_2=\text{CH}_2] (12.55) - \text{SE} (\sim 2) = -35.05 \approx -35.0$ kcal mol^{–1}. The stabilizing energy SE = 2 kcal mol^{–1} appeared due to polar interaction of electron donating Me- and electron withdrawing CHO-groups with EN 4.8 and 7.2 units, respectively [3]. For example, if CH₂=CH₂ → MeCH=CH₂ replacement gives $\Delta\Delta H_f^\circ -7.77$ kcal mol^{–1} [31] such replacement in CH₂=CHCHO → Z-MeCH=CHCHO or CH₂=CHCOOH → E-MeCH=CHCOOH gives SE –1.71 or –1.96 kcal mol^{–1}, respectively. Additive (macroincremental) scheme was applied for estimation of $\Delta H_f^\circ[\text{Cl}_2\text{C=CClCH=CH}_2] = \Delta H_f^\circ[\text{H}_2\text{C=CHCH=CH}_2] (26.29) + \Delta H_f^\circ[\text{Cl}_2\text{C=CHCl}] (-2.3[16]) - \Delta H_f^\circ[\text{CH}_2=\text{CH}_2] (12.55) = 11.44 \approx 11.5$ kcal mol^{–1}.


For calculation of ΔH_f° CCl₃CHO molecule we first estimated ΔH_f° CCl₃COOH. Using ΔH_f° CCl₃COCl (–57.3[31]) and CCl₃COOPr (–109.9) and Benson's contributions [36] we derived ΔH_f° CCl₃COOH –102.71 and –101.79 kcal mol^{–1}, respectively. Taking the mean value –102.25 and using again Benson's group contributions [36] we estimated C(Cl)₃(CO) group contribution –9.05 kcal mol^{–1} and then calculated $\Delta H_f^\circ[\text{CCl}_3\text{CHO}] = [\text{C}(\text{Cl})_3(\text{CO})] + [\text{C}(\text{H})(\text{CO})\text{R}] = -9.05 - 29.0[36] = -38.15 \approx -38.1$ kcal mol^{–1}. Now we have $\Delta\Delta H_f^\circ\{[\text{C}(\text{Cl})_3(\text{C})] (-19.62[36]) \rightarrow [\text{C}(\text{Cl})_3(\text{CO})] (-9.05)\} = 10.57$ kcal mol^{–1}. Since the Br-atom is larger than Cl-atom we ascribe arbitrary larger $\Delta\Delta H_f^\circ\{[\text{C}(\text{Br})_3(\text{C})] (15.84[36]) \rightarrow [\text{C}(\text{Br})_3(\text{CO})]\} \sim 12$ kcal mol^{–1}. Thus, $[\text{C}(\text{Br})_3(\text{CO})] \approx 15.84 + 12 \approx 28$ kcal mol^{–1} and $\Delta H_f^\circ[\text{CBr}_3\text{CHO}] = 28 - 29.1 \approx -1$ kcal mol^{–1}.

The ΔH_f° values for some of XCH=CH₂ molecules were calculated applying enthalpic shift procedure originally suggested by Benson [37] then in Ref. [20] and in many other works (see Ref. [3]) and introduced in practice as a completed methodology in Ref. 3. As a model system for XCH=CH₂ molecules the substituted benzenes PhX were selected with now well known enthalpic shift $\Delta\Delta H_f^\circ(\text{CH}_2=\text{CHX} \rightarrow \text{PhX}) \approx 7.2$ kcal mol^{–1} [3,20,38–40]. Thus, taking $\Delta H_f^\circ -24.67$ [31] for PhCOCl, –20.72 [31] for PhCOCH₃, –17.27 [32, 41] for PhOCH₃ and adding ~ -7.2 kcal mol^{–1} we

obtained ΔH_f° values -32 , -28.5 and -24.5 kcal mol $^{-1}$ for $\text{CH}_2=\text{CHCOCl}$, $\text{CH}_2=\text{CHCOCH}_3$ and $\text{CH}_2=\text{CHOCH}_3$ molecules, respectively. A larger procedure was performed for calculation of the enthalpies of formation for $\text{CH}_2=\text{CHEH}_2$ (E = As, Sb and Bi) molecules. The ΔH_f° values for model PhEH_2 molecules are derived by interpolation of the data for EH_3 and EPh_3 molecules [interpolated $\Delta\Delta H_f^\circ$ and ΔH_f° values in kcal mol $^{-1}$ – in square brackets]



Now, taking ΔH_f° (calc.) for PhAsH_2 , PhSbH_2 and PhBiH_2 we add $\sim - (7.2-7.5)$ kcal mol $^{-1}$ [3,20] and obtain ΔH_f° values for $\text{CH}_2=\text{CHAsH}_2$, $\text{CH}_2=\text{CHSbH}_2$ and $\text{CH}_2=\text{CHBiH}_2$ molecules 35.5, 57 and 75 kcal mol $^{-1}$. Earlier ΔH_f° $\text{CH}_2=\text{CHPH}_2$ 19 kcal mol $^{-1}$ was estimated by other methods [32]. We leave this value for calculation of $\text{CH}_2=\text{C}(\bullet)\text{PH}_2$ free radical although estimation of $\text{CH}_2=\text{CHPH}_2$ from ΔH_f° Ph_3P gives close results (ΔH_f° and $\Delta\Delta H_f^\circ$ – in kcal mol $^{-1}$):



The ΔH_f° of Ph_3P is rather uncertain. Several values are quoted: 71.9 ± 3.8 [44], 76.5 [45], 78 ± 5 [46] and 78.4 ± 5 [43] kcal mol $^{-1}$. We suggest the value 75 kcal mol $^{-1}$ as the compromise value. Now, we calculate $\Delta H_f^\circ\{\text{CH}_2=\text{CHPH}_2\} = 26.5 - 7.2 = 19.3$, compare with 19 kcal mol $^{-1}$ [32]. From Q values for $\text{CH}_2=\text{CH}\bullet + \text{CH}_2=\text{CHEH}_2 + \text{CH}_2=\text{C}(\bullet)\text{EH}_2 + \text{Q}$ isodesmic reactions (E = N, As, and Sb) (Table 3) we estimate by inter- and extrapolation those Q values for E = P and Bi:

EH_2 :	H_2N	PH_2	AsH_2	SbH_2	BiH_2
$\text{Q (kcal mol}^{-1}\text{):}$	4.76	[7]	9.2	10	[11]

then calculate the enthalpies of formation for $\text{CH}_2=\text{C}(\bullet)\text{PH}_2$ and $\text{CH}_2=\text{C}(\bullet)\text{BiH}_2$ free radicals, e.g. $\Delta H_f^\circ [\text{CH}_2=\text{C}(\bullet)\text{PH}_2] = \Delta H_f^\circ [\text{CH}_2=\text{CH}\bullet] - \Delta H_f^\circ [\text{CH}_2=\text{CH}_2] + \Delta H_f^\circ (\text{CH}_2=\text{CHPH}_2) - \text{Q} = 70 - 12.55 + 19.3 - 7 = 70$ kcal mol $^{-1}$ and $\nu_{\text{CH}} [\text{CH}_2=\text{CHPH}_2] = \nu_{\text{CH}} (\text{CH}_2=\text{CH}_2)(3060) - \text{Q} (7) \times 11$ (11 cm $^{-1}$ – equivalent for 1 kcal mol $^{-1}$ for vinylic =C–H bond) = ~ 2980 cm $^{-1}$.

For estimation of ΔH_f° $\text{CH}_2=\text{C}(\text{X})\text{Me}$ molecules for two of them (X = Me and Cl) we possess experimental values, i.e. -4.0 (Me) and -5.0 ± 2.2 (Cl) kcal mol $^{-1}$ [31] the latter having rather large uncertainty. From these data we gain $\Delta\Delta H_f^\circ [\text{CH}_2=\text{CHMe}(4.88) \rightarrow \text{CH}_2=\text{CMe}_2(-4.0)] = -8.88$ and $\Delta\Delta H_f^\circ \{\text{CH}_2=\text{CHCl}(-5.2) \rightarrow \text{CH}_2=\text{C}(\text{Cl})\text{Me}(-5)\} = -10.2$ kcal mol $^{-1}$. The latter value seems quite reasonable since the observed difference $-10.2 - (-8.8) = -1.4$ kcal mol $^{-1}$ (revealing stabilization of system due to CH_3/Cl polar effect) might be smaller compared with 1,2-isomers (see above). Taking the known EN values for X-substituents 10 (F), 9.0 (OH), 7.9 (CN), 7.4 (MeO), 7.1 (Cl), 6.2 (Br), 5.3 (I), 4.8 (Me) [3] and $\Delta\Delta H_f^\circ$ values -8.88 [$\text{CH}_2=\text{CHMe} \rightarrow \text{CH}_2=\text{C}(\text{Me})_2$] and -10.2 [$\text{CH}_2=\text{CHCl} \rightarrow \text{CH}_2=\text{C}(\text{Me})\text{Cl}$] kcal mol $^{-1}$ we roughly assign the following $\Delta\Delta H_f^\circ$ [$\text{CH}_2=\text{CHX} \rightarrow \text{CH}_2=\text{C}(\text{Me})\text{X}$] values -11.5 (F), -11 (OH), -11 (CN), -10.5 (MeO), -10.2 (Cl), -9.7 (Br), -9.2 (I) and using the ΔH_f° values for $\text{CH}_2=\text{CHX}$ obtain the following ΔH_f° (calc.) for $\text{CH}_2=\text{C}(\text{X})\text{Me}$ molecules: -46.5 (X = F), -41.4 (OH), 32 (CN), -35 (MeO), -5.0 (Cl) [31], 8.3 (Br), 22.8 (I). According to Ref.20 such substituents as OH, OMe, CN, Cl and Me have similar steric requirements while Br and I are larger in their “size” compared with Me-group in $\text{CH}_2=\text{CMe}_2$ molecule. Therefore, we slightly increase their ΔH_f° values to 8.5 and 23.5 kcal mol $^{-1}$, respectively.

Recently, the $\Delta H_f^\circ -65.26$ kcal mol $^{-1}$ for $\bullet\text{CH}_2\text{C}(\text{F})=\text{O}$ free radical was estimated [47] (compare with our value -59 or -60.5 [30] kcal mol $^{-1}$). When this one is inserted in isodesmic reaction $\bullet\text{CH}_2\text{CHO}$ (3) + $\text{CH}_3\text{C}(\text{F})=\text{O}$ (-105.66) \rightarrow $\bullet\text{CH}_2\text{C}(\text{F})=\text{O}$ (-65.26 [47]) + CH_3CHO (-39.73 kcal mol $^{-1}$) + Q the positive Q value $+2.33$ kcal mol $^{-1}$ is obtained

revealing the stabilization (?) of the free radical center at H-replacement by electron-withdrawing F-atom in non-conjugative position. This is doubtful result.

7 REFERENCES

- [1] D. A. Ponomarev, V. V. Takhistov, Structural Aspects and Thermochemical Approach to Isomerization and Fragmentation of Negative Ions: Skeletal Rearrangements, Molecular Ion and Artifacts. *J. Mol. Struct.*, in press.
- [2] D. A. Ponomarev, V. V. Takhistov, Isodesmic Reactions and Thermochemistry of Free Radicals. *Abstracts of Fifth Int. Conf. Chem. Kinet.*, July 16–20, Gaithersberg, USA, , **2001**, 61–62.
- [3] A. V. Golovin, V. V. Takhistov, Thermochemistry of organic and heteroorganic species. Part XII. Mono- and disubstituted acetylenes and ethynyl free radicals. New electronegative scale. *J. Mol. Struct.* **2004**, *701*, 57–91.
- [4] A. V. Golovin, V. V. Takhistov, Interrelationship for ν_{CH} values in RH molecules and enthalpy of formation of R• free radicals in aliphatic compounds, *J. Mol. Struct.* **2005**, *734*, 219–233.
- [5] D. C. McKean, Individual CH bond strength in simple organic compounds, *Chem. Soc. Rev.* **1978**, *7*, 399–422 and refs therein.
- [6] D. C. McKean, CH bond dissociation energies, isolated stretching frequencies and radical stabilisation energy, *Int. J. Chem. Kinet.* **1989**, *21*, 445–464.
- [7] D. C. McKean, I. L. Duncan, L. Blatt, CH-stretching frequencies, bond lengths and dissociation energies, *Spectrochim. Acta* **1973**, *23A*, 1037–1049.
- [8] D. C. McKean, Quantum-chemical studies of fluoroethanes: vibrational assignments, isolated CH stretching frequencies, valence force constants, and bond lengths relationships, *J. Phys. Chem.* **2000**, *A104*, 8995–9008.
- [9] D. C. McKean, A. L. McPhail, H. G. M. Edwards, I. R. Lewis, V. S. Mastryukov, J. E. Roggs, Vibrational spectra in the $\nu(\text{SiH})$ and $\nu(\text{SiD})$ regions of chloro- and bromodisilanes and ab initio geometry studies of $\text{C}_2\text{H}_5\text{Cl}$, $\text{Si}_2\text{H}_5\text{Cl}$ and 1,1- $\text{Si}_2\text{H}_4\text{Cl}_2$, *Spectrochim. Acta*, **1993**, *49A*, 1079–1094.
- [10] D. C. McKean, CH-stretching frequencies, bond lengths and strengths in halogenated ethylenes, *Spectrochim. Acta*, **1975**, *31A*, 1167–1186.
- [11] S. W. Benson, Bond Energies, *J. Chem. Educ.* **1965**, *42*, 502–518.
- [12] Y.-R. Luo, *Handbook of Bond Dissociation Energies*, CRC Press, Boca Raton, 2003.
- [13] Yu. D. Orlov, Yu. A. Lebedev, I. Sh. Saifullin, *Thermochemistry of Organic Free Radicals (Russ.)*, Nauka (Science), Moscow, 2001.
- [14] D. A. Ponomarev, V. V. Takhistov, Some regularities in thermodynamic stability of free radicals, *J. Mol. Struct.* **1997**, *435*, 259–274.
- [15] V. I. Tulin, L. A. Koroleva, L. N. Bazhenova, S. V. Krasnoshchekov, L. S. Surmina, V. V. Ivannikova, Yu. A. Pentin, Vibrational spectra and rotational isomerism of the β,β -dimethylacrolein molecule, *Russ. J. Phys. Chem.* **2004**, *78*, 1088–1096.
- [16] M. Frenkel, K. N. Marsh, R. C. Wilhoit, G. J. Kabo, G. N. Roganov. *Thermodynamics of Organic Compounds in the Gas Phase*, Thermodynamics Research Center, College St. TX, USA. 1994.
- [17] I. Barnes, K. H. Becker, J. Starcke, The gas-phase IR-spectra of formyl iodide and carbonyl iodide, *Chem. Phys. Lett.* **1995**, *246*, 594–600.
- [18] G. Wu, S. Shlykov, C. Van Arsenoy, H. J. Geise, E. Sluyts, B. J. Van der Veken, Formic anhydride in the gas phase, studied by electron diffraction and microwave and IR spectroscopy, supplemented with ab initio calculations of geometries and force fields, *J. Phys. Chem.* **1995**, *99*, 8589–8598.
- [19] D. F. McMillen, D. M. Golden, Hydrocarbon bond dissociation energies, *Ann. Rev. Phys. Chem.* **1982**, *33*, 493–532.
- [20] V. V. Takhistov, *Organic Mass Spectrometry (Russ.)*, Nauka (Science), Leningrad, 1990.
- [21] C. Hansch, A. Leo, R. W. Taft, A survey of Hammett substituent constants and resonance and field parameters, *Chem. Rev.* **1991**, *91*, 165–195.
- [22] A. Benidar, R. LeDoucen, J.-C. Guillemin, O. Mo, M. Yanez, Vibrational spectra of vinylarsine and vinylstibine. An experimental and theoretical study, *J. Phys. Chem.* **2002**, *106A*, 6262–6270.
- [23] N. L. Allinger, L. Yan, K. Chen, Molecular mechanics calculations (MM2 and MM3) on enamine and aniline derivatives, *J. Comput. Chem.* **1994**, *15*, 1321–1330 and references therein.
- [24] J. E. Katon, W. R. Fairheller, Jr., Vibrational spectra and rotational isomerism in acrylyl chloride and related α,β -unsaturated acyl chlorides, *J. Chem. Phys.* **1967**, *47*, 1248–1255.
- [25] C. M. Richards, J. R. Nielsen, Raman spectrum of 1,3-butadiene in the gaseous and liquid state, *J. Opt. Soc. Amer.* **1950**, *40*, 438–441.
- [26] J. R. Durig, D. T. Durig, T. J. Dickson, M. Jalilian, Y. Jin, J. F. Sullivan, IR and Raman spectra, conformational

- stability, barriers to internal rotation, vibrational assignment, and ab initio calculations for methyl vinyl sulfide, *J. Mol. Struct.* **1998**, *442*, 71–92.
- [27] Yu. N. Panchenko, V. I. Mochalov, Yu. A. Pentin, Some trends in vibrational spectra and the structure of chloro- and methyl-substituted butadienes-1,3, *Vestnik Moscow Univ. (Russ.)*, ser.2, **1984**, 131–144.
- [28] M. Z. El-Sabban, B. J. Zwolinski, Normal Coordinate Treatment and Assignment of Fundamental Vibrations of Vinyl Chloride and the seven Deuterated Chlorides, *J. Mol. Spectrosc.* **1968**, *27*, 1–16.
- [29] I. N. Domnin, V. V. Takhistov, D. A. Ponomarev, Thermochemistry of organic and heteroorganic species. Part V. Photoionization studies of ionization and fragmentation of polysubstituted cyclopropenes: 3-methyl-3-acetyl and related compounds. *Eur. Mass. Spectrom.* **1999**, *5*, 169–182.
- [30] V. M. Orlov, A. D. Misharev, V. V. Takhistov, I. I. Ryabinkin, Enthalpy of formation of acetyl radical and its analogs in the gas phase. *Russ. Chem. Bull. (Engl. Transl.)* **1989**, 1337–1338.
- [31] J. B. Pedley, R. D. Naylor, S. P. Kirby. *Thermochemical Data of Organic Compounds*, 2nd ed., Chapman and Hall, New York, 1986.
- [32] Yu. D. Orlov, G. Bouchoux, V. V. Takhistov, D. A. Ponomarev, A new estimation scheme of the enthalpies of formation for RCH₂• free radicals, *J. Mol. Struct.* **2001**, *608*, 109–122.
- [33] V. V. Takhistov, T. A. Pashina, N. G. Ismagilov, V. M. Orlov, A. A. Rodin, V. G. Barabanov, Experimental and Theoretical Determination of the Enthalpies of Formation of Molecules, Cations, and Radicals Containing Fluoroalkyl and Other fluorinated Groups, *Russ. J. Org. Chem. (Engl. Transl.)*. **1995**, *31*, 1582–1588.
- [34] M. Born, S. Ingemann, N. M. M. Nibbering, Experimental determination of the enthalpies of formation of formyl cyanide in the gas phase, *J. Phys. Chem.* **1996**, *100*, 17662–17769.
- [35] A. V. Golovin, D. A. Ponomarev, V. V. Takhistov, Thermochemistry of Organic and Elementoorganic species. Part VIII. Ketene and structurally related species, *J. Mol. Struct.* **2000**, *524*, 259–280.
- [36] S. W. Benson, *Thermochemical Kinetics*, J. Wiley, New York. 1976.
- [37] S. W. Benson, Thermochemistry and Kinetics of Sulfur-containing Molecules and Radicals, *Chem. Rev.* **1978**, *78*, 23–36.
- [38] V. V. Takhistov, *Thesese Dr.Sci.Dissertation*, Leningrad (Russia), 1989.
- [39] J. F. Liebman, A. Greenberg, Survey of the heats of formation of three-membered-ring species *Chem. Rev.* **1989**, *89*, 1225–1246.
- [40] Y.-R. Luo, J. L. Holmes, σ - π and n - π Stabilization energies in vinyl and phenyl compounds, *J. Phys. Org. Chem.* **1994**, *7*, 403–411 and refs therein.
- [41] J. D. Cox, G. Pilcher, *Thermochemistry of Organic and Organometallic Compounds*, Academic Press, London, 1970.
- [42] D. A. Ponomarev, V. V. Takhistov, The enthalpies of formation of the hydrides of main group elements, *J. Mol. Struct.* **1999**, *477*, 91–103.
- [43] G. Pilcher, in: *Int.Rev.Sci., Phys.Chem., Ser.2, Vol.10, Chapter 2*. Skinner H.A., ed. Butterworth, London, 1975 (according to quotation in Ref. 44).
- [44] W. V. Steele, The standard enthalpies of formation of the triphenyl compounds of the group V elements. 2. Triphenylbismuth and the phenyl-bismuth mean bond dissociation energy, *J. Chem. Thermodyn.* **1979**, *11*, 187–192.
- [45] D. R. Kirklin, E. S. Domalski, Enthalpies of combustion of triphenylphosphine and triphenylphosphine oxide, *J. Chem. Thermodyn.* **1988**, *20*, 743–754.
- [46] S. G. Lias, G. E. Bartmess, J. F. Liebman, J. L. Holmes, R. D. Levin, W. G. Mallard. Gas-phase ion and neutral thermochemistry, *J. Phys. Chem. Ref. Data.* **1988**, *17*, Suppl 1.
- [47] A. A. Hoops, J. R. Gascooke, K. E. Kautzman, A. E. Faulhaber, D. M. Neumark, Photodissociation spectroscopy and dynamics of the CH₂CFO radical, *J. Chem. Phys.* **2004**, *120*, 8494–8504.