

Internet Electronic Journal of Molecular Design

July 2004, Volume 3, Number 7, Pages 379–386

Editor: Ovidiu Ivanciuc

Proceedings of the Internet Electronic Conference of Molecular Design, IECMD 2003
November 23 – December 6, 2003
Part 1

Computational Chemical Analysis of Enantiomer Separations of Derivatized Amino Acids in Reversed–phase Liquid Chromatography

Toshihiko Hanai¹

¹ Health Research Foundation, Institute Pasteur 5F, Sakyo–ku, Kyoto 606–8225 Japan

Received: October 22, 2003; Accepted: January 31, 2003; Published: July 31, 2004

Citation of the article:

T. Hanai, Computational Chemical Analysis of Enantiomer Separations of Derivatized Amino Acids in Reversed–phase Liquid Chromatography, *Internet Electron. J. Mol. Des.* **2004**, *3*, 379–386, <http://www.biochempress.com>.

Computational Chemical Analysis of Enantiomer Separations of Derivatized Amino Acids in Reversed-phase Liquid Chromatography[#]

Toshihiko Hanai^{1,*}

¹ Health Research Foundation, Institute Pasteur 5F, Sakyo-ku, Kyoto 606-8225 Japan

Received: October 22, 2003; Accepted: January 31, 2003; Published: July 31, 2004

Internet Electron. J. Mol. Des. 2004, 3 (7), 379–386

Abstract

A fast quantitative structure–retention relationship method is required in chromatography for rapid optimization of chromatographic separation conditions. Chromatographic data of amino acid enantiomers were analyzed using a computational chemical method to simulate chromatographic separation. Using computational chemical calculations, the direct interaction between a model–phase and an enantiomer was calculated as an energy value using the MM2 calculation. Computational chemistry using a model adsorbent is a new method for quantitative analysis of retention in reversed-phase liquid chromatography. The correlation coefficient was 0.938 ($n = 14$) between the retention factors of derivatized amino acids and interaction energy values of the final structure (ΔFS) between an analyte and a model pentyl–bonded phase.

Keywords. QSRR; liquid chromatography; computational chemistry; derivatized amino acid enantiomers.

Abbreviations and notations

| | |
|-----------------------------|---|
| FS, final structure energy | VW, van der Waals energy |
| HB, hydrogen bonding energy | MM, molecular mechanics |
| ES, electrostatic energy | FEF, (+)–1–(9–fluorenyl)ethyl chloroformate |

1 INTRODUCTION

A variety of chiral recognition phases have been synthesized and their chiral selectivity has been analyzed using a computational chemical method, especially for Pircle type phases [1–9]. The calculated energy values predicted the elution order of enantiomers but did not reflect the separation factor (α) of the enantiomers. One reason for this is that the density of bonded chiral phases is not consistent compared to alkylsilane–bonded phases, whose carbon contents and reproducibility in chromatography are consistent [10]. Thus, the reproducibility of retention time is not as reliable for chiral phases, and it is difficult to develop a model phase for computational chemical analysis. Untreated silanols affect the retention of analytes because hydrogen bonding is the predominant

[#] Presented in part at the Internet Electronic Conference of Molecular Design, IECMD 2003.

* Correspondence author; phone: 81–45–547–4871; fax: 81–45–547–4874; E–mail: thanai@attglobal.net.

retention force for enantiomer separations in normal-phase liquid chromatography.

On the other hand, analytes derivatized with a chiral derivatization reagent have been separated in reversed-phase liquid chromatography [11–18]. A computational chemical simulation of the retention of phenolic compounds in reversed-phase liquid chromatography was achieved using a molecular mechanics calculation with a model phase where the retention factors of phenolic compounds, even partially ionized phenolic compounds, were related to calculated energy values with high precision [19].

In this present paper, reversed-phase liquid chromatography of derivatized amino acids was analyzed using a molecular mechanics calculation, and the retention factors and separation factors were measured in isocratic elution related to energy value changes between an analyte and a model-phase.

2 COMPUTATIONAL CHEMICAL ANALYSIS

A Power Macintosh G3 computer equipped with a 450 MHz processor and 512 MB memory, and a Dell model Latitude C840 equipped with a 2 GHz processor and 1024 MB memory were used. The molecular properties of analytes and model phases and molecular interactions were calculated by molecular mechanics (MM2) using version 5 of the CAChe™ program from Fujitsu, Tokyo, Japan.

The standard parameters used were bond stretch, bond angle, dihedral angle, improper torsion, van der Waals force, hydrogen bond, and electrostatic energy (MM2 bond dipoles). The van der Waals force cut-off distance was 9Å. The energy unit was kcal/mol (1 kJ/mol = 4.18 kcal/mol). The Cricket-Graph™ program from Computer Associates (San Diego, CA, USA) and Project Reader of CAChe™ program were used for data analysis.




Figure 1. Model dimethylpentylsilane bonded-silica gel. Small white ball, hydrogen; large white ball, carbon; small black ball, oxygen; large black ball, silicone.

3 RESULTS AND DISCUSSION

A model phase was constructed based on dimethylpentylsilane, consisting of 991 atoms, 1051 bonds, and 15193 connectors, containing 171 silicones, 328 oxygens, 143 carbons, and 349 hydrogens. Dimethylpentylsilane (20) and trimethylsilane (1) were bonded within 900 \AA^2 in the polysilicone dioxide phase. The trimethylsilane was considered to be an end-capped molecule. The optimized energy value change was less than 0.00001 kcal/mol . The molecular size and alkyl-chain length were determined by the calculation capacity of the computer used and the alkyl-chain length effect for the hydrophobicity [10]. An optimized structure of the model phase and a complex between this model phase and an enantiomer is shown in Figures 1 and 2, respectively. Dimethylpentyl groups are in close proximity due to their steric hindrance. Some of them are disposed in free space after the optimization of molecular interactions.




Figure 2. Adsorption of FEF-S-arginine in the dimethylpentylsilane bonded silica gel.

Table 1. Chromatographic data and calculated energy values

| Amino acid | log k ^a | α^b | α^c |
|-----------------|--------------------|------------|------------|
| Serine R | 0.303 | | |
| Serine S | 0.320 | 1.04 | 1.09 |
| Aspartic acid R | 0.375 | | |
| Aspartic acid S | 0.405 | 1.07 | 1.03 |
| Threonine R | 0.468 | | |
| Threonine S | 0.520 | 1.13 | 1.00 |
| Glutamic acid R | 0.513 | | |
| Glutamic acid S | 0.567 | 1.14 | 1.07 |
| Arginine R | 0.549 | | |
| Arginine S | 0.605 | 1.10 | 1.09 |
| Alanine R | 0.751 | | |
| Alanine S | 0.792 | 1.00 | 0.95 |
| Proline R | 0.741 | | |
| Proline S | 0.741 | 1.09 | 1.04 |
| Asparagine R | 0.127 | | |
| Asparagine S | 0.164 | 1.09 | 1.04 |
| Glutamine R | 0.111 | | |
| Glutamine S | 0.164 | 1.13 | 1.04 |

^a converted from retention factors in [14]

^b separation factor (R/S) calculated from reference retention factors [14]

^c separation factor (R/S) calculated from molecular interaction energy ΔF_S

| Amino acid | FE | HB | ES | VW |
|-----------------|----------|---------|----------|----------|
| Serine R | -35.288 | -20.927 | -7.579 | 6.052 |
| Serine S | -33.205 | -19.384 | -7.165 | 5.786 |
| Aspartic acid R | -60.000 | -21.349 | -26.546 | 7.460 |
| Aspartic acid S | -59.709 | -24.120 | -26.975 | 6.120 |
| Threonine R | -31.208 | -11.589 | -7.939 | 5.806 |
| Threonine S | -38.323 | -24.938 | -8.073 | 6.657 |
| Glutamic acid R | -49.242 | -19.914 | -19.216 | 4.161 |
| Glutamic acid S | -48.433 | -17.487 | -18.687 | 6.944 |
| Arginine R | -50.046 | -14.696 | -33.766 | 6.954 |
| Arginine S | -52.529 | -8.815 | -34.164 | 5.952 |
| Alanine R | -30.544 | -10.952 | -12.045 | 5.970 |
| Alanine S | -35.001 | -13.680 | -11.973 | 6.923 |
| Proline R | -15.360 | -3.649 | -10.543 | 7.054 |
| Proline S | -15.683 | -3.735 | -10.490 | 6.595 |
| Asparagine R | -65.290 | -20.718 | -41.294 | 7.529 |
| Asparagine S | -66.426 | -23.673 | -40.734 | 7.351 |
| Glutamine R | -55.929 | -15.865 | -30.937 | 6.201 |
| Glutamine S | -54.911 | -17.048 | -30.485 | 6.926 |
| Model phase | -648.713 | 0.000 | -403.448 | -400.646 |

In this bonded-phase, dimethylpentyl groups surrounded one trimethyl group. Silanol groups around the trimethylsilanol group were completely covered by alkyl groups. The silanol group effect might not be a factor. The first circle of the dimethylpentyl groups might not be pushed down by the presence of an analyte. The second circle of dimethylpentyl groups should support the first circle of dimethylpentyl groups. The calculated energy values of the amino acid enantiomers

derivatized with (+)-1-(9-fluorenyl)ethyl chloroformate by MM2 are listed in Table 1 with chromatographic data measured in reversed-phase liquid chromatography from reference 14.

Table 1. (Continued)

| Amino acid | FS1 | HB1 | ES1 | VW1 |
|-----------------|----------|---------|----------|----------|
| Serine R | -710.131 | -18.668 | -410.908 | -421.484 |
| Serine S | -710.505 | -24.279 | -409.569 | -418.956 |
| Aspartic acid R | -736.477 | -18.916 | -429.722 | -419.922 |
| Aspartic acid S | -737.038 | -17.503 | -430.095 | -425.358 |
| Threonine R | -712.766 | -14.450 | -411.557 | -422.818 |
| Threonine S | -719.963 | -24.770 | -411.591 | -423.652 |
| Glutamic acid R | -729.851 | -11.721 | -422.989 | -426.524 |
| Glutamic acid S | -731.292 | -22.132 | -422.767 | -424.641 |
| Arginine R | -731.092 | -12.143 | -438.115 | -423.195 |
| Arginine S | -736.621 | -16.415 | -437.406 | -426.794 |
| Alanine R | -716.373 | -15.449 | -414.971 | -424.942 |
| Alanine S | -718.915 | -16.922 | -415.852 | -424.892 |
| Proline R | -701.278 | -3.615 | -413.991 | -429.290 |
| Proline S | -703.224 | -3.660 | -413.679 | -431.488 |
| Asparagine R | -745.840 | -25.170 | -444.725 | -421.302 |
| Asparagine S | -748.306 | -23.438 | -444.369 | -422.947 |
| Glutamine R | -738.278 | -21.341 | -436.870 | -424.949 |
| Glutamine S | -738.620 | -14.028 | -434.048 | -432.012 |




Figure 3. Relationship between Δ energy and $\log k$, No.1, R-Ser; 2, S-Ser; 3, R-Asp; 4, S-Asp; 5, R-Thr; 6, R-Glu; 7, S-Thr; 8, R-Arg; 9, S-Glu; 10, S-Arg; 11, R-Pro; 12, S-Pro; 13, S-Ala; 14, R-Ala; 15, R-Gln; 16, R-Asn; 17, S-Asn; 18, R-Gln; Δ energy = 22.127 ($\log k$) + 20.692, $r = 0.938$.

The calculated energy values are final (FS), hydrogen bonding (HB), electrostatic (ES), and van der Waals (VW) energy. The energy values of individual complexes between the model pentyl phase and an enantiomer are listed in Table 2 as FS1, HB1, ES1, and VW1. For docking between an analyte and the model phase, the most hydrophobic site of the analyte was faced towards the hydrophobic model phase whose silanol activity was eliminated.

The molecular interaction energy (Δ energy) was calculated from the following equation: Δ energy = energy value of an analyte + energy value of the model phase – energy value of a complex. The Δ energy values were related to retention factors of the amino acid derivatives. The correlation coefficient between Δ FS and $\log k$ is shown in Figure 3.

$$\Delta FS = 22.127(\log k) + 20.692, r=0.938, n=14 \quad (1)$$

A one-to-one molecular interaction model might be ideal for chromatography using a Pircle-type bonded phase, however this model can be applied to reversed-phase liquid chromatography of enantiomer separations, such as for phenolic compounds. Amino acids R- and S-asparagine and R- and S-glutamine do not fit in the above correlation.

Under these liquid chromatographic conditions, the elution order of amino acids was Asn < Gln < Ser < Asp < Thr < Glu < Arg < Ala < Pro. The interaction energy values of Asn and Gln, however, were high. The elution order of amino acids derivatized with orthophthaldialdehyde was Asp < Glu < Cys < Asn < Ser < Gln < Thr < Arg < Ala [20].

The difference in elution order depends on the organic modifiers used in the eluent. The former system [14] used tetrahydrofuran and the latter system used acetonitrile and methanol. Tetrahydrofuran is a selective organic modifier in reversed-phase liquid chromatography. The elution order of analytes did not have a linear relation with the octanol-water partition coefficient, $\log P$ relative hydrophobicity, in reversed-phase liquid chromatography using tetrahydrofuran as an organic modifier [21].

Acetonitrile is the most suitable organic modifier to obtain a high correlation coefficient between $\log k$ and $\log P$. Therefore, if the purpose is to predict the elution order, the retention factors should be measured in reversed-phase liquid chromatography using acetonitrile as the organic modifier. In the chromatographic results, all R-amino acids eluted before the S-amino acids, however the interaction energy value of S-alanine was smaller than that of R-alanine (Table 1).

More precise calculations obtained using a better model phase and solvent effect might resolve this discrepancy. Even though the solvent effect is not predicted by the above calculation, computational chemical simulation of reversed-phase liquid chromatography is useful for determining enantiomer separation.

4 CONCLUSIONS

The retention factors and the separation factors of amino acid enantiomers derivatized with (+)-1-(9-fluorenyl)ethyl chloroformate measured in isocratic elution on reversed-phase liquid chromatography were related to energy value changes between an analyte and a model-phase. The energy values of their final structures was correlated well to log *k* values of the amino acid derivatives.

5 REFERENCES

- [1] K. B. Lipkowitz, Atomistic modeling of enantioselection: applications in chiral chromatography, *Theor. Comput. Chem.* **1998**, *5*, 329–379.
- [2] T. Hanai, H. Hatano, N. Nimura, T. Kinoshita, Computational chemical analysis of chiral recognition in liquid chromatography, selectivity of N-(R)-1-(α -naphthyl)ethylaminocarbonyl-(R or S)-valine and N-(S)-1-(α -naphthyl)ethylaminocarbonyl-(R or S)-valine bonded aminopropyl silica gel, *Anal. Chim. Acta* **1996**, *332*, 213–224.
- [3] T. Hanai, H. Hatano, N. Nimura, T. Kinoshita, Computational chemical analysis of the separation of derivatized R- and S-amino acid enantiomers on N-(tert-butylaminocarbonyl)-(S)-valylaminopropylsilica gel and (R)-1-(α -naphthyl)ethylaminocarbonyl-glycylaminopropyl silica gel by liquid chromatography, *J. Liq. Chromatogr. & Rel. Technol.* **1996**, *19*, 1189–1204.
- [4] T. Hanai, H. Hatano, N. Nimura, T. Kinoshita, Molecular recognition in chromatography aided by computational chemistry, *Supramol. Chem.* **1994**, *3*, 243–247.
- [5] K. B. Lipkowitz, A. G. Anderson, Computational aspects in supramolecular chemistry: Chiral discrimination in chromatography, *NATO ASI Series, Series C: Mathematical and Physical Sciences* **1994**, *426*, 183–198.
- [6] R. Kalisz, T. A. G. Noctor, I. W. Wainer, Stereochemical aspects of benzodiazepine binding to human serum albumin. II. Quantitative relationships between structure and enantioselective retention in high performance liquid affinity chromatography, *Mol. Pharmacol.* **1992**, *42*, 512–517.
- [7] R. Daepfen, H. R. Karfunkel, F. J. J. Leusen, Computational chemistry applied to the design of chiral stationary phases for enantiomeric separation, *J. Comp. Chem.* **1990**, *11*, 181–193.
- [8] S. Topiol, M. Sabio, Computational chemical studies of chiral stationary phase models. Complexes of methyl N-(2-naphthyl)alaninate with N-(3,5-dinitrobenzoyl)leucine n-propylamide, *J. Chromatogr.* **1989**, *461*, 129–137.
- [9] K. Akira, H. Hasegawa, Y. Shinohara, M. Imachi, T. Hashimoto, Stereoselective internal acyl migration of 1 β -O-acyl glucuronides of enantiomeric 2-phenylpropionic acids, *Biol. Pharm. Bull.* **2000**, *23*, 506–510.
- [10] T. Hanai, New developments in liquid chromatographic stationary phases, *Adv. Chromatogr.* **2000**, *40*, 315–357.
- [11] N. Nimura, A. Toyama, Y. Kasahara, T. Kinoshita, Reversed-phase liquid chromatographic resolution of underivatized D,L-amino acids using chiral eluents, *J. Chromatogr.* **1982**, *239*, 671–675.
- [12] D. W. Aswad, Determination of D- and L-aspartate in amino acid mixtures by high-performance liquid chromatography after derivatization with a chiral adduct of o-phthalaldehyde, *Anal. Biochem.* **1984**, *137*, 405–409.
- [13] N. Nimura, T. Kinoshita, o-Phthalaldehyde-N-acetyl-L-cysteine as a chiral derivatization reagent for liquid chromatographic optical resolution of amino acid enantiomers and its application to conventional amino acid analysis, *J. Chromatogr.* **1986**, *352*, 169–177.
- [14] S. Einarsson, B. Josefsson, P. Moller, D. Sanchez, Separation of amino acid enantiomers and chiral amines using precolumn derivatization with (+)-1-(9-fluorenyl)ethyl chloroformate and reversed-phase liquid chromatography, *Anal. Chem.* **1989**, *59*, 1191–1195.
- [15] R. Schuster, Determination of amino acids in biological, pharmaceutical, plant and food samples by automated precolumn derivatization and high-performance liquid chromatography, *J. Chromatogr.* **1988**, *432*, 271–284.
- [16] C. Carducci, M. Birarelli, V. Leuzzi, G. Santaga, P. Serafini, I. Antonozzi, Automated method for the measurement of amino acids in urine by high-performance liquid chromatography, *J. Chromatogr. A* **1996**, *729*,

173–180.

- [17] U. Spengler, J. Kirschbaum, I.-G. Baek, H. Bruckner, *Proc. Germ. Nutr. Soi.* **1999**, *1*, 57.
- [18] N. Todoroki, K. Shibata, T. Yamada, Y. Kera, Ryo-hei Yamada, Determination of N-methyl-D-aspartate in tissues of bivalves by high-performance liquid chromatography, *J. Chromatogr. B* **1999**, *728*, 41–47.
- [19] T. Hanai, C. Mizutani, H. Homma, Computational chemical simulation of chromatographic retention of phenolic compounds, *J. Liq. Chromatogr. Rel. Technol.* **2003**, *26*, 2031–2039.
- [20] Y. Ishida, Amino acid; in: *Liquid Chromatography in Biomedical Analysis*, Ed. T. Hanai, Elsevier, Amsterdam, 1991, pp. 47–80.
- [21] T. Hanai, (Ed) *HPLC A Practical Guide*, Royal Chemical Society, 1999, pp. 57–65.

Biographies

Toshihiko Hanai is research scientist at Health Research Foundation. After obtaining Ph.D. degree in analytical chemistry from Kyoto University, Dr. Hanai undertook postdoctoral research with Professor B. Karger at Northeastern University, and with Professors H. Walton and R. Sieverse at Colorado University at Boulder. Dr. Hanai was postdoctoral research fellow then research assistant at Universite de Montreal. He was research manager at GasukuroKogyo. Dr. Hanai collaborated on pharmaceutical analysis with Professors T. Kinoshita and H. Homma at Kitasato University. Dr. Hanai published 13 scientific books including 7 books in English about HPLC. His research interesting is Chromatography and Computational Chemistry for Drug Discovery to accelerate drug discovery as an analytical chemist.