Highly diastereoselective synthesis of 2,3-dihydro-9b*H*- thiazolo[2,3*a*]isoindolin-5-ones

Steven M. Allin,^{*a} Darshan G. Vaidya,^a Michael I. Page^b, and Alexandra M. Z. Slawin^a

*a Department of Chemistry, Loughborough University, Loughborough, Leicestershire, LE11 3TU, England ^b Department of Chemical & Biological Sciences, University of Huddersfield, Queensgate, Huddersfield HD1 3DH, England E-mail: <u>S.M.Allin@lboro.ac.uk</u> (received 18 Nov 99; accepted 25 Apr 00; published on the web 03 May 00)

Abstract

Condensation of 2-acetyl benzoic acid with aminothiols proceeds with extremely high diastereoselectivity to produce the desired 2,3-dihydro-9b*H*-thiazolo[2,3-*a*]isoindolin-5-one products in good yield. The relative stereochemistry of the major diastereoisomer, formed exclusively when using non-racemic aminothiol substrates, has been determined for the first time by single crystal X-ray analysis.

Keywords: Thiazolo[2,3-*a*]isoindolin-5-ones, diastereoselective synthesis

Introduction

The chemistry of the isoindolinone ring system is currently an area of interest for many research groups due to the actual and potential biological activities of many derivatives.¹ We have recently initiated a programme of study aimed at developing the chemistry of isoindolin-1-ones such as (1), and we have developed new and stereoselective routes to 3-substituted isoindolinones (2) and isoindoloisoquinolines (3) from such tricyclic lactam precursors.²



Due to our general interest in this area of heterocyclic chemistry, we turned our attention to the 2,3-dihydro-9bH-thiazolo[2,3-a]isoindolin-5-one ring system (4). Chiral compounds of this type are of interest as non-nucleosidic reverse transcriptase inhibitors.³ In this communication we

present a highly diastereoselective synthesis of the 2,3-dihydro-9bH-thiazolo[2,3-*a*]isoindolin-5-one ring system.

Results and Discussion

In general, equimolar amounts of 2-acylbenzoic acids and the corresponding $\tilde{}$ -aminothiols were heated at reflux under Dean-Stark conditions in xylene solvent for 24 h (Scheme 1). Analysis of the crude product mixture by 250 MHz ¹H-NMR spectro-scopy showed clean and efficient conversion to the desired products (5) [Table]. With the non-racemic, chiral β -aminothiol substrates, th e product was observed as a single diastereoisomer. Purification was achieved by flash column chromatography. The relative stereochemistry of the non-racemic products derived from L-cysteine ethyl and methyl esters was confirmed by single crystal X-ray analysis; the X-ray of the ethyl ester derivative is presented in the Figure.



Scheme 1

 Table 1. Synthesis of 2,3-Dihydro-9bH-thiazolo[2,3-a]isoindolin-5-ones, (5)

Product	R	R^1	Yield (%)	diastereoselectivity
5a	Н	Н	95	N/A
5b	Н	Me	88	N/A
5c	Н	Et	40	N/A
5d	Н	Bn	20	N/A
5e	CO ₂ Et	Н	80	exclusive ^b
5f	CO ₂ Et	Me	70	exclusive
5g	CO ₂ Me	Н	80	exclusive
5h	CO ₂ Me	Me	65	exclusive

(a) determined by 250 MHz ¹H-NMR spectroscopy (b) minor isomer not visible by 250 MHz ¹H-NMR spectroscopy



Figure 1

As can be seen from the Figure,the major product diastereoeisomer adopts a "bowl-like" shape, with the ester substituent lying on the outer (convex) face. Based on our earlier work in related areas,² we propose a mechanism to explain the stereochemical outcome of the reaction outlined in Scheme 1 which involves initial formation of an iminothiol/thiazolidine intermediate (Scheme 2).



Scheme 2

Reversible cyclization of the iminothiol (6) could produce both the *trans* thiazolidine (7) and the *cis* thiazolidine (8). Ring closure of (6) with concomitant loss of water yields the observed *trans* tricyclic product (5). In the case of (8) however, cyclization to (9) appears from simple molecular models to be disfavored due to more remote orientation of the reactive functional groups. Tricyclic compound (5) appears to be the thermodynamically more favorable product

isomer. No interconversion from (5) to (9) is observed by NMR spectroscopy on stirring a solution of (5) at room temperature for 7 days.

In summary, we report a highly efficient and direct method for the synthesis of 2,3-dihydro-9b*H*-thiazolo[2,3-*a*]isoindolin-5-ones products in good yield. The relative stereochemistry of the major diastereoisomer, formed exclusively when using non-racemic aminothiol substrates, has been determined for the first time by X-ray crystal analysis.

Experimental Section

General Procedures. All solvents, where necessary, were dried, distilled and stored over 4Å molecular sieves prior to use. Reagent chemicals were purchased from Lancaster Synthesis Ltd. and the Aldrich Chemical Co. Ltd, and were purified before use as required. Flash-column chromatography was carried out using Merck silica gel (70 - 230 Mesh ASTM). Pressure was applied at the column head using hand bellows. Analytical thin layer chromatography (TLC) was carried out using aluminium-backed plates coated with 0.2 mm silica. Plates were visualised under UV light (at 254 nm) or by staining with either potassium permanganate solution or iodine. Infra-red spectra were recorded in the range 4000600 cm¹, using Perkin Elmer Paragon 100 FT-IR Spectrometer, with internal calibration. Solid samples were run as Nujol mulls, and liquids as thin films. ¹H Nuclear magnetic resonance spectra were recorded using either a Bruker Avance 400 MHz Spectrometer or a Bruker AC 250 MHz Spectrometer. Multiplicities were recorded as broad peaks (br), singlets (s), doublets (d), triplets (t), quartets (q), double doublets (dd) and multiplets (m) etc. ¹³C Nuclear magnetic resonance spectra were recorded using either a Bruker Avance DPX-400 MHz Spectrometer (run at 100 MHz) or a Bruker AC 250 MHz Spectrometer (run at 67.5 MHz). Mass spectra were recorded using a Fisons VG Quattro II SQ instrument and Accurate-Mass mass spectra were recorded using a Kratos MS80 instrument. X-Ray crystallography was carried out using a Rigaku AFC7S diffractometer with graphite monochromated Cu-Ka radiation. Crystallographic data has been deposited at the Cambridge Crystallographic Data Centre (ref: CCDC140138). Elemental analyses were carried out on a Perkin Elmer 2400 CHN Elemental Analyser. Melting points were determined on a 9100 Thermal Apparatus melting point block and are uncorrected Yields are for isolated pure products. Optical rotations were performed using an Optical Activity AA - 10 Automatic Polarimeter.

General procedure for synthesis of 2,3-dihydro-9b*H*-thiazolo[2,3-*a*]isoindolin-5-ones

To a stirred solution of the keto acid in xylene (mixture of isomers, 150 mL) was added an equimolar amount of the aminothiol along with a three-fold molar excess of sodium acetate portion-wise over 5 h under reflux. The mixture was heated for a further 24 h under Dean-Stark conditions. The resultant solution was filtered and the xylene removed under reduced pressure. Purification was achieved by flash column chromatography with ether/petrol mixtures as eluent.

250 MHz ¹H NMR analysis of crude non-racemic products confirmed the formation of a single diastereoisomer. Recrystallisation was achieved from ether/hexanes mixtures.

2,3-Dihydro-9bH-thiazolo[**2,3-a**]**isoindolin-5-one** (**5a**). ⁴Off white crystals; 3.019 g, 95%; mp 99 101 °C (ether/hexanes) lit.⁴ 98100.5 °C; v_{max} (Nujol)/cm¹ 1701 (N-C==O), 1468 (Ar); $\delta_{\rm H}$ (CDCl₃, 400MHz) 3.33 - 3.42 (3H, m, -NCH₂CH₂S-), 4.38 - 4.57 (1H, m, -NCH₂CH₂MS-), 5.85 (1H, s, N-CH-S), 7.46 - 7.59 (3H, m, Ar-H), 7.78 - 7.80 (1H, m, Ar-H); $\delta_{\rm C}$ (CDCl₃, 400MHz) 36.9, 44.8, 66.4, 123.6, 125.2, 129.6, 131.5, 133.0, 145.5, 171.2; *m*/*z* (EI) M⁺ 191 (100%), 158 (42), 145 (90), 132 (28), 117 (23), 104 (11), 90 (15), 77 (10); (Found 191.0404. C₁₀H₉NOS requires 191.0404).

9b-Methyl-2,3-dihydro-9bH-thiazolo[2,3-a]isoindolin-5-one (5b). Orange oil; 1.40 g, 88%; v_{max} (thin film)/cm¹ 1701 (N-C==O), 1466 (Ar); δ_{H} (CDCl₃, 250MHz) 1.91 (3H, s, CH₃), 3.33 - 3.47 (3H, m, -NCH₂CH₂S-), 4.52 - 4.60 (1H, m, -NCH₂CH₂S-), 7.43 - 7.62 (3H, m, Ar-H), 7.77 (1H, d, *J* 7.3, Ar-H); δ_{C} (CDCl₃, 400MHz) 28.9, 38.0, 43.4, 76.0, 122.1, 124.5, 129.5, 129.9, 133.3, 150.9, 170.8; *m*/*z* (EI) 205 (100%), 190 (15), 158 (60), 146 (55), 130 (19), 103 (10); (Found 205.0564. C₁₁H₁₁NOS requires 205.0561).

9b-Ethyl-2,3-dihydro-9bH-thiazolo[**2,3***-a*]**isoindolin-5-one** (**5c**). Yellow oil; 0.44 g, 40%; v_{max} (thin film)/cm¹ 1705 (N-C==O), 1466 (Ar): δ_{H} (CDCl₃, 250MHz) 0.72 (3H, t, *J* 7.3, -CH₂CH₃), 2.27 (2H, q, *J* 7.8, -CH₂CH₃), 3.31 - 3.43 (3H, m, -NCH₂CH₂S-), 4.49 - 4.61 (1H, m, -NCH₂CH₂S-), 7.41 - 7.46 (2H, m, Ar-H), 7.55 - 7.61 (1H, m, Ar-H), 7.75 - 7.78 (1H, m, Ar-H); δ_{C} (CDCl₃, 400MHz) 9.3, 34.1, 37.2, 43.2, 79.4, 121.9, 123.9, 127.7, 130.2, 132.8, 148.8, 171.0; *m/z* (EI) M⁺ 219 (82%), 204 (14), 190 (100), 172 (67), 160 (23), 130 (14), 119 (13), 91 (8), 77 (9); (Found 219.0718. C₁₂H₁₃NOS requires 219.0717).

9b-(PhenyImethyl)-2,3-dihydro-9bH-thiazolo[2,3-*a*]isoindolin-5-one (5d). Colourless crystals; 0.094 g, 20% (Found: C, 72.09; H, 5.30; N, 4.97%. $C_{17}H_{15}NOS$ requires C, 72.60; H, 5.34; N, 4.98%). mp 108 - 110 °C (ether/hexanes); v_{max} (Nujol)/cm¹ 1701 (N-C==O), 1465 (Ph-H); δ_{H} (CDCl₃, 250MHz); 3.16 - 3.33 (3H, m, -NCH₂CH₂S-), 3.46 (2H, q, *J* 8.8, CH₂Ph), 4.45 - 4.51 (1H, m, -NCH₂CH₂S-), 7.06 - 7.10 (2H, m, Ar-H), 7.12 - 7.17 (3H, m, Ar-H), 7.37 - 7.46 (2H, m, Ar-H), 7.52 - 7.59 (1H, m, *J* 1.2 7.6, Ar-H), 7.65 (1H, dt, *J* 0.9 6.45, Ar-H); δ_{C} (CDCl₃, 400MHz) 37.9, 44.3, 47.4, 79.6, 122.6, 122.7, 125.5, 127.1, 127.4, 128.3, 129.2, 130.2, 130.7, 132.9, 137.8, 149.5, 171.3; *m/z* (EI) M⁺ 281 (4%), 234 (38), 220 (18), 204 (14), 191 (100), 178 (30), 163 (20), 130 (61), 102 (32), 91 (51), 77 (17); Found 281.0870. $C_{17}H_{15}NOS$ requires 281.0874).

(3*S*, 9b*R*)-Ethyl-2,3-dihydro-9b*H*-thiazolo[2,3-*a*]isoindolin-5-on-3-carboxylate(5e). White crystals; 2.69 g, 80% (Found C, 59.21; H, 4.84; N, 5.39%. C₁₃H₁₃NO3S requires C, 59.32; H, 4.94; N, 5.32%); mp 77 - 80 °C (ether/hexanes); $[\alpha]_D = 225.2^\circ$ [c = 1.84, CHCl₃]; v max (Nujol)/cm¹1731 (COOEt), 1710 (N-C==O), 1468 (Ar); $\delta_{\rm H}$ (CDCl₃, 250MHz) 1.31 (3H, t, *J* 7.2, COOCH₂CH₃) 3.55 - 3.68 (2H, m, -S-CH₂-), 4.29 (2H, q, *J* 7.2, COOCH₂CH₃), 5.24 (1H, dd, *J* 5.0 7.2, -NCHCOOEt), 6.08 (1H, s, N-CH-S-), 7.45-7.61 (3H, m, Ar-H), 7.80 (1H, d, *J* 7, Ar-H); $\delta_{\rm C}$ (CDCl₃, 400MHz) 15.4, 40.1, 58.1, 62.4, 66.6, 123.7, 124.9, 129.7, 130.2, 133.4, 145.3,

170.3, 170.6; m/z (EI) M⁺ 263 (78%), 190 (80), 132 (33), 91 (39); (Found 263.0618. C₁₃H₁₃O₃NS requires 263.0616).

(3S,9bR)-3-Ethyl-9b-methyl-2,3-dihydro-9bH-thiazolo[2,3-a]isoindolin-5-on-3-

carboxylate(5f). White crystals; 2.07 g, 70% (Found: C, 60.56; H, 5.49; N, 5.38%. $C_{14}H_{15}NO_3S$ requires C, 60.65; H, 5.42; N, 5.05%); mp 110 - 113 °C (ether/hexanes); $[\alpha]_D = 273.10^\circ$ [c = 1.1, CHCl₃]; v_{max} (Nujol)/cm¹ 1731 (COOEt), 1710 (N-C==O), 1468 (Ar); δ_H (CDCl₃, 250MHz) 1.33 (3H, t, *J* 7.1, COOCH₂CH₃), 1.96 (3H, s, CH₃), 3.77 -3.98 (2H, m, -S-CH₂-), 4.24 - 4.34 (2H, m, COOCH₂CH₃), 5.14 (1H, dd, *J* 6.3 8.7, -NCHCOOEt), 7.47 - 7.64 (3H, m, Ar-H), 7.78 - 7.82 (1H, m, Ar-H); δ_C (CDCl₃, 250MHz) 14.5, 28.5, 40.6, 58.4, 62.3, 122.3, 124.9, 126.3, 129.4, 129.5, 135.0, 149.4, 170.7, 170.9; *m*/*z* (EI) M⁺ 277 (100%), 262 (23), 204 (80), 146 (88); (Found 277.0774. $C_{14}H_{15}NO_3S$ requires 277.0772).

(35,9bR)-3-Methyl-2,3-dihydro-9b*H*-thiazolo[2,3-a]isoindolin-5-on-3-carboxylate (5g). White needles; 2.69 g, 80% (Found C, 57.62; H, 4.49; N, 5.67%. $C_{12}H_{11}NO_3S$ requires C, 57.83; H, 4.42; N, 5.62%); mp 83 - 86 °C (ether/hexanes); $[\alpha]_D = 233.6^\circ$ [c = 2.31, CHCl₃]; v_{max} (Nujol)/cm¹ 1731 (COOEt), 1710 (N-C==O), 1468 (Ar); δ_H (CDCl₃, 400MHz) 3.60 - 3.69 (2H, m, -SCH₂) 3.82 (3H, s, COOCH₃), 5.25 (1H, dd, *J* 4.8 7.6, -NCHCOOMe), 6.08 (1H, s, -N-CH-S), 7.47 - 7.52 (2H, m, Ar-H), 7.57 - 7.59 (1H, m, Ar-H), 7.80 - 7.82 (1H, m, Ar-H); δ_C (CDCl₃, 400MHz) 40.0, 53.3, 57.9, 66.7, 123.8, 125.0, 129.8, 130.8, 133.4, 145.1, 170.8, 170.9; *m/z* (EI) M⁺ 249 (15%), 227 (4), 209 (36), 190 (14), 135 (100), 119 (7), 107 (59), 93 (50); (Found 249.0455. $C_{12}H_{11}O_3NS$ requires 249.0459).

(3*S*,9*bR*)-3-Methyl-9b-methyl-2,3-dihydro-9*bH*-thiazolo[2,3-a]isoindolin-5-on-3carboxylate (5h). White needles; 2.07 g, 70%; mp 113 - 116 °C (ether/hexanes); [α]_D = 289.6° [c = 1.75, CHCl₃]; v_{max} (Nujol)/cm¹ 1750 (COOEt), 1693 (N-C==O), 1466 (Ar); δ_{H} (CDCl₃, 400MHz) 1.94 (3H, s, COOCH₃), 3.79 - 3.94 (2H, m, -SCH₂), 3.81 (3H, s, CH₃), 5.14 (1H, dd, *J* 6.4 8.8, -NHCHCOOMe), 7.47 - 7.50(2H, m, Ar-H), 7.57 - 7.59 (1H, m, Ar-H), 7.77 - 7.79 (1H, m, Ar-H); δ_{C} (CDCl₃, 400MHz) 27.9, 40.1, 52.8, 57.7, 77.3, 121.8, 125.3, 128.9, 129.3, 137.2, 148.9, 170.2, 170.9; *m/z* (EI) M⁺ 263 (5%), 163 (26), 160 (100), 119 (20), 100 (33), 87 (11); (Found 263.0619. C₁₃H₁₃NO₃S requires 263.0616).

Acknowledgements

We thank the Universities of Loughborough and Huddersfield for facilities and financial support of this research.

References

1. Csende, F.; Szabo, Z.; Stajer, G. *Heterocycles* **1993**, *36*, 1809. (b) De Clercq, E. *J. Med. Chem.* **1995**, *38*, 2491. (c) Takahashi, I.; Hirano, E.; Kawakami, T.; Kitajima, H.

Heterocycles **1996**, *43*, 2343. (d) Epsztajn, J.; Grzelak, R.; Jozwiak, A. *Synthesis* **1996**, 1212. (e) Marchalin, S.; Decroix, B. *Heterocycles***1995**, *41*, 689. (f) Zhuang, Z.P.; Kung, M.P.; Mu, M.; King, H.F. J. Med. Chem. **1998**, *41*, 157.

- Allin, S.M.; Northfield, C.J.; Page, M.I.; Slawin, A.M.Z. Tetrahedron Lett., 1997, 38, 3627;.(b) Allin, S.M.; Northfield, C.J.; Page, M.I.; Slawin, A.M.Z. Tetrahedron Lett., 1998, 39, 4905. (c) Allin, S.M.; Northfield, C.J.; Page, M.I.; Slawin, A.M.Z. Tetrahedron Lett. 1999, 40, 141. (d) Allin, S.M.; Northfield, C.J.; Page, M.I.; Slawin, A.M.Z. Tetrahedron Lett. 1999, 40, 141. (d) Allin, S.M.; Northfield, C.J.; Page, M.I.; Slawin, A.M.Z. Tetrahedron Lett.
- Schafer, W.; Friebe, W.-G.; Leinert, H.; Mertens, A.; Poll, T.; von der Saal, W.; Zilch, H.; Nuber, B.; Ziegler, M.L. *J. Med. Chem.* 1993, *36*, 726. (b) Mertens, A.; Zilch, H.; Konig, B.; Schafer, W.; Poll, T.; Kampe, W.; Seidel, H.; Leser, U.; Leinert, H. *J. Med. Chem.* 1993, *36*, 2526. (c) De Clercq, E. *J. Med. Chem.* 1995, *38*, 2491.
- 4. Hiskey, R.G.; Dominianni, S.J. J. Org. Chem., **1965**, 30, 1506.