

New Fungal Anthraquinones¹

Melvyn Gill* and Peter M. Morgan

School of Chemistry, The University of Melbourne, Victoria, 3010

E-mail: melvyn@unimelb.edu.au

Dedicated to Prof. D. W. Cameron to mark his retirement and contributions²

(received 01 May 01; accepted 19 Dec 01; published on the web 27 Dec 01)

Abstract

The 1-*O*-methyl ethers **2** and **11** of physcion and emodin, respectively, and emodin anthrone **13** are isolated for the first time from Basidiomycotina. They are found, along with the parent quinones **1** and **10**, the tetrahydroanthraquinones (*S*)-atrochryson **8** and (*S*)-torosachryson **9**, a mixture of atropisomeric flavomannin 6,6'-di-*O*-methyl ethers **4**, and the rare anthraquinone fallacinol **7**, from among a group of Australian and New Zealand toadstools belonging to the genus *Dermocybe*.

Keywords: Natural products, fungal pigments, anthraquinones, asidiomycotina, Agaricales

Introduction


We have reported extensively on the chemical constituents of toadstools indigenous to Australia and New Zealand.^{3,4} Many of these secondary metabolites are derived from the polyketide pathway and are either dihydroanthracenones (monomeric⁵ and dimeric⁶) or anthraquinones proper.⁷ We describe here the isolation of three compounds: two anthraquinones and an anthrone, that are recorded for the first time from Basidiomycotina. Several new records of other compounds of this class are also recorded here for the first time.

Results and Discussion

The diminutive fruiting bodies of *Dermocybe* sp. WAT 22963⁸ were collected in the Kinglake and Otway Ranges National Parks, Victoria from under mixed *Eucalyptus* sp. and *Nothofagus cunninghamii*, respectively. The sporophore is characterised by red-brown cap skin and bright orange gills. We have reported previously on the major red pigments⁹ and their glycosidic derivatives¹⁰ from *D.* sp. WAT 22963 and a description of the fungus has been published.⁹ Drawn


by the complex TLC profile obtained from the organic soluble extractives obtained from this fungus, the minor constituents were examined.

Fresh fungal material was chopped and macerated in ethanol overnight and the extract was evaporated to dryness. The residue was partitioned between water and ethyl acetate and the organic phase was concentrated and purified sequentially by gel permeation and prep. TLC. The most mobile yellow zone (R_f 0.80) obtained by prep. TLC was isolated as orange needles, mp 204–205 °C in a yield of $1.7 \times 10^{-3}\%$ of the fresh weight of the fungus. The molecular formula $C_{16}H_{12}O_5$ followed from HR EIMS while UV–visible absorption at 430 nm and IR absorption at 1681 and 1628 cm^{-1} suggested an anthraquinone structure.¹¹ The 1H NMR spectrum (Table) revealed the presence of *C*-methyl and *O*-methyl groups, four aromatic protons and the protons of two hydrogen bonded phenolic hydroxy groups. These data identified this pigment as physcion **1**, a conclusion that was confirmed by direct comparison with an authentic sample.¹² Physcion **1** is widespread but only as a trace constituent in *Cortinarius* and *Dermocybe* species.⁷




- 1** $R^1 = R^2 = H$
2 $R^1 = H, R^2 = Me$
3 $R^1 = Me, R^2 = H$

A second yellow zone (R_f 0.73) gave orange needles, mp 191–192 °C in a yield of $8 \times 10^{-3}\%$ of the fresh weight of the fungus. The EIMS shows a molecular ion at m/z 298 ($C_{17}H_{14}O_5$) that corresponds to the elements CH_2 more than physcion **1**. The H NMR spectrum confirms the presence of two methoxy groups (δ 3.91 and 4.06), one *C*-methyl group (δ 2.51), four aromatic protons (δ 6.70, 7.15, 7.31 and 7.77) and one phenolic hydroxy group (δ 13.30). The aromatic protons at δ 6.70 and 7.31 are *meta*-coupled ($J = 2.45$ Hz), while the signals at δ 7.15 and 7.77 are broadened by both *meta*- and allylic coupling. In the absence of further evidence these data do not differentiate between the 1- and the 8-*O*-methyl ethers **2** and **3**, respectively, of physcion **1**. The ether **2** has been isolated from plant tissue cultures,¹³ while **3** is a metabolite of plants and microorganisms.¹⁴ Comparison of the physical and spectroscopic data for the quinone from *Dermocybe* sp. WAT 22963 with those published for synthetic physcion 1-*O*-methyl ether **2**¹⁵ and for physcion 8-*O*-methyl ether **3**¹⁴ established its identity with the former (Table). This is the first time that physcion 1-*O*-methyl ether **2** has been isolated from Basidiomycotina.




The least mobile zone (R_f 0.18) from prep. TLC gave flavomannin 6,6'-di-*O*-methyl ether **4** as a yellow-green powder, mp 201–204 °C, in a yield of 6.7×10^{-2} % of the fresh weight of the fungus. A molecular ion at m/z 574 in the EIMS was consistent with the molecular formula $C_{32}H_{30}O_{10}$ while the 1H NMR spectrum, which shows signals from only one half of the number of protons in the molecule, is characteristic of a 7,7'-linked dihydroanthracenone dimer.⁷ Flavomannins and their ethers, such as **4**, are known from *Cortinarius* and *Dermocybe* in various stereochemical forms.⁷ The presence in each half of the molecule of (at least) one stereogenic centre (C-3) coupled with atropisomerism about the hindered biaryl axis introduces considerable stereochemical complexity into these molecules that has only recently been addressed.¹⁶ The absolute configuration at the biaryl axis in compounds of the flavomannin series is conveniently determined from the CD spectrum. Thus, the signs of the strong bisignate Cotton effects close to 275 nm can be related directly to the stereochemistry at the axis.¹⁷ The CD spectrum of flavomannin 6,6'-di-*O*-methyl ether **4** from *D. sp.* WAT 22963 shows a negative Cotton effect ($\Delta\epsilon$ -9.0) at 293 nm and a positive Cotton effect ($\Delta\epsilon$ +17.1) at 271 nm consistent with (*M*)-chirality at the axis.^{16,17} The stereochemistry of the C-3 and C-3' chiral centres in **4** was determined by adding, in turn, authentic samples of the (3*R*,3'*R*,*M*)- and (3*S*,3'*S*,*M*)-stereoisomers of **4**, from *Cortinarius citrinus* and *C. pseudosulphureus*, respectively,¹⁸ to a sample of **4** in deuteriochloroform and recording the 1H NMR spectrum. The 1H NMR spectrum of the solution containing **4** together with the (3*R*,3'*R*,*M*)-isomer showed a single set of resonances with no discernible broadening or doubling of the C-6/6' methoxy or C-8/8' hydroxy resonances. In contrast, the 1H NMR spectrum of the solution containing **4** together with the (3*S*,3'*S*,*M*)-isomer showed distinct broadening of the signal at δ 3.83 (C-6/6'-OMe) and clear doubling of the signal at δ 9.98 (C-8/8'-OH). These results establish unequivocally that the flavomannin 6,6'-di-*O*-methyl ether isolated from *D. sp.* WAT 22963 is predominantly the (3*R*,3'*R*', 3*M*)-stereoisomer shown in formula **4**.


Having established that the flavomannin derivative **4** was not present in *D. sp.* WAT 22963 in admixture with any of its diastereoisomers, we considered next its enantiomeric purity. (3*R*,3'*R*,*M*)-Flavomannin 6,6'-di-*O*-methyl ether **4** was earlier isolated by us from the Australian *Dermocybe sp.* WAT 20880.¹⁶ That sample exhibited extrema in the CD spectrum at 287 nm ($\Delta\epsilon$ -68.3) and 265 nm ($\Delta\epsilon$ +72.4), which suggests that the sample described here is an anisochiral mixture¹⁹ of **4** with its (3*S*,3'*S*,*P*)-enantiomer *ent*-**4** in a ratio of no more than 62:38 (24% ee) in favour of the former.



Dermocybe sp. WAT 26644 was collected from under *Nothofagus menziesii* in the Caitlins River Park, South Otago, New Zealand in May 1995. The fruiting bodies very closely resemble those of *Dermocybe* sp. WAT 22963 described previously,⁹ and a taxonomic relationship between the two species was supported by preliminary TLC analysis of the ethanolic extracts of the New Zealand taxon. The TLC reveals the presence in *D.* sp. WAT 26644 of five pigments with the same chromatographic properties as compounds previously isolated from *D.* sp. WAT 22963.^{9,10} The extracts of *D.* sp. WAT 26644 were subsequently partitioned between ethyl acetate and water and the organic phase was separated and the contents purified by gel permeation and prep. TLC. Physcion **1**, its methyl ether **2**, and the xanthorin derivatives **5** and **6**, previously reported from *D.* sp. WAT 22963, were identified by direct comparison with authentic materials.⁹ A yellow zone (R_f 0.25) that is not present in the TLC of *D.* sp. WAT 22963, was isolated as orange needles, mp 242–244 °C, in a yield of $1.0 \times 10^{-2}\%$. The molecular formula $C_{16}H_{12}O_6$ was determined by HR EIMS and the pigment was identified as fallacinol **7** from the IR (ν 1630 and 1671 cm^{-1}) and UV–visible data (λ 412 nm) and from the 1H NMR spectrum. Finally, comparison of all of the spectroscopic data for the yellow pigment from *D.* sp. WAT 26644 with the literature data for fallacinol **7** confirmed the identity of the pigment.²⁰ Fallacinol **7** has been isolated previously only from *Dermocybe cinnabarina*²⁰ but has been detected chromatographically in extracts of several other *Cortinarius* species.²¹ Prior to our recent discovery of *w*-hydroxyxanthorin 1-*O*-methyl ether **6**,⁹ fallacinol **7** was the only *w*-substituted anthraquinone known from Basidiomycetes.



8 R = H
9 R = Me



10 R¹ = R² = H
11 R¹ = H, R² = Me
12 R¹ = Me, R² = H


The green-yellow zone (R_f 0.18), $C_{32}H_{30}O_{10}$, from the New Zealand *Dermocybe* sp. WAT 26644 was identified as a mixture of (3*R*,3'*R'*,*M*)-flavomannin 6,6'-di-*O*-methyl ether **4** and its (3*R*,3'*R'*,*P*)-atropisomer (not shown) from the 1H NMR and CD spectra. Thus, the 1H NMR spectrum showed distinct doubling of the signals due to the C-6/6' methoxy groups (δ 3.82 and 3.84) and the C-8/8' hydroxy groups (δ 9.98 and 10.14) consistent with the presence of two diastereoisomers. That these are atropisomers rather than epimers is apparent from the CD spectrum, which shows no significant net absorption in the vicinity of 275 nm. Weak absorption at shorter wavelength (Experimental) accords with the CD spectrum of (*R*)-torosachryson **ent-9** and supports the (3*R*,3'*R*) central chirality in both of the flavomannin isomers present in *D. sp.* WAT 26644. An atropisomeric mixture of flavomannin 6,6'-di-*O*-methyl ethers of undefined absolute configuration has been isolated before from *Tricholoma auratum*.⁷

It is clear from the chemical analysis of the New Zealand taxon *D. sp.* WAT 26644 that it must be very close taxonomically to the Australian taxon *D. sp.* WAT 22963. The only differences in the pigment composition of the two species being (i) the presence in the former of fallacinal **7** and (ii) the stereochemical constitution of the flavomannin 6,6'-di-*O*-methyl ether that is present in each species.

Fruit bodies of *Dermocybe* sp. WAT 26813 were collected in leaf litter under *Nothofagus cunninghamii* at Mait's Rest, Otway Ranges National Park, Victoria. They have a pale orange-brown cap and a yellow-green stipe. TLC analysis of the ethanolic extract revealed the presence of two green pigments (R_f 0.24 and 0.35) and three yellow pigments (R_f 0.42, 0.48 and 0.65). The residue remaining after evaporation of the solvent was partitioned between ethyl acetate and water and the organic phase was purified by prep. TLC. Green zones (R_f 0.24 and 0.35) were identified as (*S*)-atrochryson **8** and (*S*)-torosachryson **9**, respectively, from their physical, chiroptical and spectroscopic properties and by direct comparison with authentic materials.²² (*S*)-Atrochryson **8** has been reported previously from several European *Dermocybe* species⁷ but only once before from an Australian member of the genus.²² (*S*)-Torosachryson **9**, on the other hand, is common among Australian dermocybes⁴ but has yet to be recorded from members of the genus indigenous to other parts of the World.

The most mobile yellow pigment (R_f 0.65) from *D. sp.* WAT 26813 was identified as emodin

10 ($8.3 \times 10^{-3}\%$), a common natural product.^{11,13,14} from the spectroscopic data and by comparison with a commercial sample. The more polar yellow zones (R_f 0.48 and 0.42) contained metabolites that have not previously been recorded from *Dermocybe* or *Cortinarius*. The least polar of the two was isolated as orange needles, mp 265–267 °C, in a yield of $3.3 \times 10^{-3}\%$ of the fresh weight of the fungus. The formula $C_{16}H_{12}O_5$ followed from HR EIMS of the molecular ion at m/z 284, while the presence of an emodin chromophore was indicated by a long wavelength absorption at 429 nm [λ (emodin): 436 nm] in the UV–visible spectrum. The IR spectrum implied the presence of free and hydrogen bonded quinonoid carbonyl groups with absorptions at 1663 and 1625 cm^{-1} , respectively. The 1H NMR spectrum (Table) contains signals characteristic of aromatic *C*- and *O*-methyl groups, two pairs of *meta*-coupled aromatic protons and a hydrogen bonded phenolic hydroxy group. These data are accommodated by both the 1- and the 8-*O*-methyl ether structures **11** and **12**, respectively, of emodin **10**. Both ethers are known: the former was isolated recently from the 'aspen fungus' *Phialophora alba*, which grows on *Populus tremuloides*,²³ and from cultures of *Fusarium aquaeductuum*.²⁴ Emodin 8-*O*-methyl ether **12** is a widespread natural product.¹⁴ The methyl ethers **11** and **12** have been synthesised by Cameron²⁵ and Brassard,²⁶ respectively. Comparison of the 1H NMR data of the pigment from *D.* sp. WAT 26813 with those reported for the synthetic ethers (Table) reveals that it corresponded more closely to emodin 1-*O*-methyl ether **11** than it did to the isomer **12**. The identity of the quinone from *Dermocybe* sp. WAT 26813 with the 1-*O*-methyl ether **11** of emodin was confirmed by direct comparison with a synthetic sample.²⁵ Both the natural and synthetic specimens of **11** had the same mp and a mixed mp showed no depression. Furthermore, a single set of proton resonances was observed in the 1H NMR spectrum when the natural and synthetic materials were mixed together. Emodin 1-*O*-methyl ether **11** is an inhibitor of the growth of the parasitic fungus *Phellinus tremulae* at a concentration of 1 mg mL⁻¹.²³ This is the first report of the isolation of emodin 1-*O*-methyl ether **11** from the Basidiomycotina.



The least mobile yellow zone (R_f 0.42) gave yellow-green plates, mp 250–254 °C (decomp.) in a yield of $4.2 \times 10^{-3}\%$. The EIMS contains a molecular ion at m/z 256 while the FAB MS exhibits $[M+Na]^+$ and $[M+1]^+$ ions at 279 and 257, respectively. The HR EIMS established the molecular formula $C_{15}H_{12}O_4$, which corresponds to two hydrogen atoms more and one oxygen atom less than the molecular formula for emodin **10**. In the 1H NMR spectrum (see Experimental) this pigment gives rise to singlets due to two hydrogen bonded phenolic hydroxy groups, a three proton singlet corresponding to an aromatic methyl group and a two proton singlet at δ 4.31. In the aromatic proton region two pairs of *meta*-disposed protons reveal a

typical emodin substitution pattern. Consideration of the molecular formula and the two proton singlet at δ 4.31 in the ^1H NMR spectrum led to the emodin anthrone structure **13** for this constituent of *D. sp.* WAT 26813. Comparison of the physical and spectroscopic data recorded here with those published for emodin anthrone²⁷ confirmed the identity of the pigment. This is the first report of emodin anthrone **13** from Basidiomycotina.

Table 1. H NMR data of the anthraquinones **1**, **2**, **3**, **11** and **12** (CDCl_3 , 400 MHz)

H -2	7.09 br s	7.15 br s	7.14 br s	7.01 br s	7.16 br s	7.16 br s	7.10 br s
3-Me	2.45 s	2.51 s	2.50 s	2.38 s	2.51 s	2.51 s	2.38
H -4	7.64 br s	7.77 br s	7.74 br s	7.49 br s	7.77 br s	7.76 br s	7.40 br s
H -5a	7.38 d	7.31 d	7.29 d	7.36 d	7.21d	7.22 d	7.21 d
H -7a	6.70 d	6.70 d	6.69 d	6.69 d	6.67 d	6.67	6.85 d
1-OH	12.13 s	-	-	11.99 s	-	-	-
6-OH	-	-	-	-	-	-	-
8-OH	12.33 s	13.30 s	13.33 s	-	13.27 s	13.28 s	-
1-OMe	-	4.06 s	4.06 s	-	4.06 s	4.06 s	-
6-OMe	3.94 s	3.91 s	3.93 s	3.96 s	-	-	-
8-OMe	-	-	-	3.92 s	-	-	3.92 s

^{a)}From *Dermocybe* sp WAT 22963.

^{b)} Authentic sample, see reference 25.

^{c)} Recorded in *d*₆-DMSO at 100 MHz.

^{d)} JH5-H7 = 2.4–2.6 Hz.

In summary, from two indigenous Australian and one indigenous New Zealand toadstool the 1-*O*-methyl ethers **2** and **11** of emodin and physcion, respectively, and emodin anthrone **13** are reported for the first time from Basidiomycotina. In addition, flavomannin 6,6'-di-*O*-methyl ether **4** is reported in different degrees of stereochemical purity from some closely related members of this group of fungi and the occurrence of the rare (*S*)-atrochryson **8** and (*S*)-torosachryson **9** is extended to several new indigenous taxa.

Experimental Section

General Procedures. Mps: Kofler hotstage apparatus; uncorr; Optical rotation: Perkin–Elmer 241 polarimeter; CD: AVIV 62DS spectrometer; Prep. TLC: Merck Kieselgel GF₂₅₄; TLC: Macherey–Nagel SIL G-25; toluene/HCO₂Et/HCO₂H (50:49:1); CC: Sephadex LH-20

(Pharmacia); NMR: JEOL JNM GX-400 spectrometer (^1H at 399.65 MHz and ^{13}C at 100.4 MHz); IR: Perkin–Elmer 983G grating spectrophotometer; UV–visible: Varian SuperScan 3 spectrophotometer using 10 mm quartz cells and EtOH as the solvent; EIMS: VG Micromass 7070F and JEOL JMS-AX505HF spectrometers (70 eV).

Plant material

Fungi were collected from the following locations (lyophilised specimens of each species are lodged in the herbarium of the Royal Botanic Garden, Edinburgh, under the accession numbers quoted): *Dermocybe* sp. WAT 22963: Kinglake (1993) and Otway Ranges (1995) National Parks, Victoria, Australia from under *Nothofagus cunninghamii*. *D.* sp. WAT 26644: Catlin's River Park, South Otago, New Zealand, from under *Nothofagus menziesii* (1995). *D.* sp. WAT 26813: Mait's Rest, Otway Ranges National Park, Victoria, from under *Nothofagus cunninghamii*.

Extraction and isolation

From *Dermocybe* sp. WAT 22963

Fresh fruiting bodies (6 g) were finely chopped and soaked overnight in EtOH (300 mL) at room temperature. The solid material was filtered off and the solvent was evaporated. The green-brown residue was partitioned between EtOAc (150 mL) and H₂O (150 mL). A second collection (24 g), from Kinglake National Park, was extracted in the same way.

The organic phase was separated, dried (Na₂SO₄) and evaporated to dryness to afford a green-brown residue (40 mg) which was purified by PLC using toluene/HCO₂Et/HCO₂H (50:49:1) as eluant to give the following compounds: physcion **1** (0.5 mg, 1.7×10^{-3} % fr. wt.) as orange needles (EtOAc–PE); mp 204–205 °C (lit.¹¹ 207–209 °C); TLC: *R_f* 0.80; IR ν (KBr) cm⁻¹: 3505, 1681, 1628; UV–Vis λ (EtOH) nm (log ϵ): 258 sh (4.28), 268 (4.34), 286 (4.33), 430 (4.14); NMR: Table; EIMS *m/z* (rel. int.): 284 (M⁺, 100); physcion 1-*O*-methyl ether **2** (2.4 mg, 8.0×10^{-3} % fr. wt.) as orange needles (CHCl₃–PE); mp 191–192 °C (lit.¹³ 188–191 °C); TLC: *R_f* 0.73; HR EIMS (rel. int.): found 298.0821 [M]⁺ (100%), requires C₁₇H₁₄O₅ 298.0845; IR ν (KBr) cm⁻¹: 3420, 1690, 1625; UV–Vis λ (EtOH) nm (log ϵ): 222 sh (3.98), 269 (4.60), 320 (3.03), 429 (3.86); NMR: Table 1; EIMS *m/z* (rel. int.): 298 (M⁺, 100), 280 (39), 269 (20), 252 (44), 149 (75); xanthorin 1-*O*-methyl ether **5** (2.7 mg, 9.0×10^{-3} % fr. wt.) as red needles (CHCl₃); mp 204–205 °C (lit.⁹ 205–209 °C); TLC: *R_f* 0.55; IR ν (KBr) cm⁻¹: 3434, 1660, 1624; UV–Vis λ (EtOH) nm (log ϵ): 221 (2.91), 230 (2.84), 236 (2.83), 256 (2.89), 294 (2.32), 469 (2.18), 495 (2.26), 530 (2.06); ^1H NMR δ 2.53 (3H, s, 3-Me), 3.99 (3H, s, 6-OMe), 4.07 (3H, s, 1-OMe), 6.71 (1H, s, H-7), 7.17 (1H, br s, H-2), 7.86 (1H, br s, H-4), 13.43 (1H, s, 5-OH), 13.98 (1H, s, 8-OH); EIMS *m/z* (rel. int.): 314 (M⁺, 100), 296 (53), 253 (22); ω -hydroxyxanthorin 1-*O*-methyl ether **6** (1.5 mg, 5.0×10^{-3} % fr. wt.) as red needles (CHCl₃); mp 220–222 °C (lit.⁹ 220–223 °C); TLC: *R_f* 0.20; IR ν (KBr) cm⁻¹: 3432, 1650, 1618, 1583; UV–Vis λ (EtOH) nm (log ϵ): 220 (2.95), 230 (2.86), 236 (2.87), 255 (3.00), 294 (2.45), 465 (2.32), 495 (2.34), 530 (2.15); ^1H NMR (CDCl₃) δ 3.99 (3H, s, 6-OMe), 4.10 (3H, s, 1-OMe), 4.89 (2H, s, 3-CH₂OH), 6.72 (1H, s, H-7), 7.51 (1H, br s, H-2), 7.98 (1H, br s, H-4), 13.41 (1H, s, 5-OH), 13.96 (1H, s, 8-OH); EIMS

m/z (rel. int.): 330 (M^+ , 74), 312 (46), 284 (30), 56 (100); and flavomannin 6,6'-di-*O*-methyl ether **4** (20.1 mg, $6.7 \times 10^{-2}\%$ fr. wt.) as a bright yellow-green powder (EtOAc-PE); mp 201–204 °C (lit.¹⁶ 207–209 °C); TLC: R_f 0.18; $[a]_{546}^{16}$ –240 22 (CHCl₃; c 0.1) (lit.¹⁶ $[\alpha]_{546}^{16}$ –860 [CHCl₃; c 0.02]); IR ν (KBr) cm^{-1} : 3384, 1627, 1550; UV-Vis λ (EtOH) nm (log ϵ): 235 (4.51), 261 (4.55), 270 (4.50), 285 (4.51), 320 (3.98), 415 (3.20); CD λ nm ($\Delta\epsilon$) 440 (+3.3), 400 (–3.0), 323 (–0.4), 293 (–9.0), 271 (+17.1), 242 (–2.8), 229 (–1.0), 216 (–2.9) nm; ¹H NMR (CDCl₃) δ 1.44 (6H, s, 3/3'-Me), 2.83 (4H, br s, H-2/2'), 3.06 (4H, br s, H-4/4'), 3.84 (6H, s, 6/6'-OMe), 6.67 (2H, s, H-5/5'), 6.93 (2H, s, H-10/10'), 9.98 (2H, s, 8/8'-OH), 16.15 (2H, s, 9/9'-OH); EIMS m/z (rel. int.): 574 (M^+ , 32), 538 (29), 507 (100), 254 (17).

From *Dermocybe* sp. WAT 26644

Fresh toadstools (7 g) were chopped and soaked overnight in EtOH (100 mL) at room temperature. Evaporation of the solvent gave a green-brown residue that was partitioned between EtOAc (50 mL) and H₂O (50 mL). The organic phase was dried (Na₂SO₄) and evaporated to afford a yellow-brown residue (20 mg) that was separated by PLC to give six pure compounds: physcion **1** (0.5 mg, $7.1 \times 10^{-3}\%$ fr. wt.); physcion 1-*O*-methyl ether **2** (0.3 mg, $4.3 \times 10^{-3}\%$ fr. wt.); xanthorin 1-*O*-methyl ether **5** (2.1 mg, $3.0 \times 10^{-2}\%$ fr. wt.); ω -hydroxyxanthorin 1-*O*-methyl ether **6** (1.3 mg, $1.9 \times 10^{-2}\%$ fr. wt.), all identical with materials described previously; fallacinal **7** (0.7 mg, $1.0 \times 10^{-2}\%$ fr. wt.) as orange needles (EtOAc-PE); mp 242–244 °C (lit.²⁰ 245–247 °C); TLC: R_f 0.25; IR ν (KBr) cm^{-1} : 3452, 1671, 1630, 1624; UV-Vis λ (EtOH) nm (log ϵ): 221 (4.22), 261 (4.21), 289 sh (3.98), 421 (3.60); UV-Vis λ (EtOH + 1% NaOH) nm (log ϵ): 512 (3.45); ¹H NMR (CDCl₃) δ 3.95 (3H, s, 6-OMe), 4.82 (2H, s, 3-CH₂OH), 6.71 (1H, d, $J = 2.2$, H-7), 7.30 (1H, br s, H-2), 7.40 (1H, d, $J = 2.2$, H-5), 7.79 (1H, br s, H-4), 12.19 and 12.29 (2 x 1H, each s, 1/8-OH); EIMS m/z (rel. int.): 300 (M^+ , 100), 271 (43), 149 (14), 98 (21), 57 (66); and flavomannin 6,6'-di-*O*-methyl ether **4** + **atrop-4** (5.6 mg, $8.0 \times 10^{-2}\%$ fr. wt.) as an orange powder (EtOAc); mp 287–289 °C; TLC: R_f 0.18; $[a]_{546}^{13}$ –412 (CHCl₃; c 0.1); IR ν (KBr) cm^{-1} : 3384, 1627, 1550; UV-Vis λ (EtOH) nm (log ϵ): 235 (4.51), 261 (4.55), 270 (4.50), 285 (4.51), 320 (3.98), 415 (3.20); CD λ (EtOH) nm ($\Delta\epsilon$) 273 (–0.9), 226 (+2.4), 211 (–4.1); ¹H NMR δ 1.44 (12H, s, 3/3'-Me), 2.83 (8H, br s, H-2/2'), 3.06 (8H, br s, H-4/4'), 3.82 (6H, s, 6/6'-OMe), 3.84 (6H, s, 6/6'-OMe), 6.67 (4H, s, H-5/5'), 6.93 (4H, s, H-10/10'), 9.98 (2H, s, 8/8'-OH), 10.14 (2H, s, 8/8'-OH), 16.15 (4H, br s, 9/9'-OH); EIMS m/z (rel. int.): 574 (M^+ , 32), 538 (29), 507 (100).

From *Dermocybe* sp. WAT 26813

Fresh fruiting bodies (12 g) were finely chopped and soaked overnight in EtOH (300 mL) at room temperature. Solid material was filtered off and the solvent was evaporated to dryness and the residue was partitioned between EtOAc (150 mL) and H₂O (150 mL). The organic phase was dried (Na₂SO₄) and evaporated to give a dark green-brown residue (40 mg). The residue was purified by PLC using toluene/HCO₂Et/HCO₂H (50:49:1) and CHCl₃/EtOH/HCO₂H (100:20:1) as the eluant. Five compounds were isolated in pure form: (*S*)-torosachryson **9** (4.6 mg, $3.8 \times 10^{-2}\%$ fr. wt.) as citrine needles (EtOAc-PE); mp 191–192 °C (lit.²² 191–194 °C); TLC: R_f 0.35; $[a]_D^{12.5}$ +7.8 (MeOH; c 0.5) (lit.²² $[\alpha]_D^{30.5}$ +7.0 [MeOH; c 1.0]); IR ν (KBr) cm^{-1} : 3505, 1640,

1590; UV-Vis λ (EtOH) nm (log ϵ): 228 (4.52), 270 (4.77), 320 (4.11), 400 (4.00); ^1H NMR (CDCl_3) δ 1.46 (3H, s, 3-Me), 2.84 (2H, m, H-2), 3.05 (2H, m, H-4), 3.89 (3H, s, 6-OMe), 6.49 (1H, d, $J = 2.2$, H-7), 6.55 (1H, d, $J = 2.2$, H-5), 6.87 (1H, s, H-10), 9.80 (1H, s, 8-OH), 16.11 (1H, s, 9-OH); EIMS m/z (rel. int.): 288 (M^+ , 100), 270 (30), 246 (10), 230 (25), 43 (20); (*S*)-atrochrysonone **8** (2.6 mg, $2.2 \times 10^{-2}\%$ fr. wt.) as green-yellow needles (CHCl_3 -PE); mp 232–236 °C (decomp.) (lit.²² 234–238 °C [decomp.]); TLC: R_f 0.24; $[\alpha]^{30.5} +6.5$ (MeOH; c 0.8) (lit.²² $[\alpha]^{25}_{546} +8.2$ [MeOH; c 0.2]); IR ν (KBr) cm^{-1} : 3460, 1642, 1600; UV-Vis λ (EtOH) nm (log ϵ): 227 (3.25), 272 (3.60), 320 (2.75), 333 sh (2.63), 398 (3.03); ^1H NMR (d_6 -acetone) δ 1.40 (3H, s, 3-Me), 2.77 (1H, dd, $J = 1.95$ and 17.3, H-2), 2.90 (1H, d, $J = 17.3$, $\text{H}_{\text{ax}}-2$), 3.00 (1H, br d, $J = 16.1$, $\text{H}_{\text{eq}}-4$), 3.08 (1H, d, $J = 16.1$, $\text{H}_{\text{ax}}-4$), 4.03 (1H, s, 3-OH), 6.38 (1H, d, $J = 2.2$, H-7), 6.58 (1H, d, $J = 2.2$, H-5), 6.84 (1H, s, H-10), 9.20 (1H, s, 6-OH), 9.84 (1H, s, 8-OH), 16.52 (1H, s, 9-OH); EIMS m/z (rel. int.): 274 (M^+ , 100), 256 (80), 232 (12), 216 (32), 43 (22); emodin 1-*O*-methyl ether **11** (0.4 mg, $3.3 \times 10^{-3}\%$ fr. wt.) as orange needles (EtOAc-PE); mp 265–267 °C (lit.²⁵ 262–265 °C); TLC: R_f 0.48; HR EIMS m/z (rel. int.) found 284.0590 [M^+] (100%), $\text{C}_{16}\text{H}_{12}\text{O}_5$ requires 284.0686; IR ν (KBr) cm^{-1} : 3400, 1665, 1625, 1590; UV-Vis λ (EtOH) nm (log ϵ): 216 (4.68), 249 (4.34), 267 sh (4.30), 282 (4.41), 429 (3.84); ^1H NMR (CDCl_3) δ 2.51 (3H, s, 3-Me), 4.06 (3H, s, 1-OMe), 6.67 (1H, d, $J = 2.45$, H-5), 7.16 (1H, br s, H-2), 7.21 (1H, d, $J = 2.45$, H-7), 7.77 (1H, br s, H-4), 13.27 (1H, s, 8OH); EIMS m/z (rel. int.): 284 (M^+ , 100), 266 (39), 255 (25), 238 (69); emodin **10** (1.0 mg, $8.3 \times 10^{-3}\%$ fr. wt.) as orange needles (EtOAc); mp 257–258 °C; TLC: R_f 0.65; IR ν (KBr) cm^{-1} : 3448, 1660, 1628; UV-Vis λ (EtOH) nm (log ϵ): 218 (4.64), 250 (4.24), 262 (4.20), 287 (4.36), 436 (3.98); ^1H NMR (CDCl_3) δ 2.47 (3H, s, 3-Me), 6.67 (1H, d, $J = 2.45$, H-5), 7.15 (1H, br s, H-2), 7.26 (1H, d, $J = 2.45$, H-7), 7.58 (1H, br s, H-4), 12.08 and 12.20 (2H, s, 1/8-OH); EIMS m/z (rel. int.): 270 (M^+ , 100); and emodin anthrone **13** (0.5 mg, $4.2 \times 10^{-3}\%$ fr. wt.) as yellow-green plates (CHCl_3); mp 250–254 °C (decomp.) (lit.²⁷ 250–258 °C [decomp.]); TLC: R_f 0.42; HR EIMS m/z (rel. int.) found 256.0701 [M^+] (100%), $\text{C}_{16}\text{H}_{12}\text{O}_5$ requires 256.0740; IR ν (KBr) cm^{-1} : 3400, 1622, 1599, 1559; UV-Vis λ (EtOH) nm (log ϵ): 258 (4.00), 268 (4.02), 302 (3.93), 360 (4.14); ^1H NMR (d_6 -acetone) δ 2.59 (3H, s, 3-Me), 4.31 (2H, s, H-10), 5.92 (1H, d, $J = 2.00$, H-5), 6.15 (1H, br s, H-2), 6.20 (1H, d, $J = 2.00$, H-7), 6.42 (1H, br s, H-4), 12.21 and 12.40 (2H, s, 1/8-OH); FAB MS m/z (rel. int.): 279 ([M^+Na^+], 45), 257 ([$\text{M}+1$] $^+$, 12), 237 (100); EIMS m/z (rel. int.): 256 (M^+ , 100), 241 (17), 228 (13), 213 (18).

Acknowledgements

We are grateful to the Australian Research Council for financial support and to Professors D. W. Cameron, for provision of a synthetic sample of emodin 1-*O*-methyl ether, and W. Steglich, for samples of natural flavommanins. P. M. M. is the recipient of a Melbourne University Postgraduate Research Scholarship. We are thankful to the Departments of Chemistry and Plant and Microbial Sciences at the University of Canterbury, Christchurch, New Zealand for

hospitality and facilities to collect and extract New Zealand species. Dr Roy Watling, Edinburgh, is acknowledged for help and encouragement (over many years) and, in the present case, for lodging herbarium specimens and crucial help with taxonomy.

References and Notes

1. Part 67 In the series Pigments of Fungi. For Part 66, see Gill, M.; Saubern, S.; Yu, J. *Aust. J. Chem.* **2000**, *53*, 213.
2. This paper is dedicated to Professor D. W. Cameron to mark his retirement as Professor of Organic Chemistry from the University of Melbourne and in recognition of his outstanding contributions to the structural elucidation and synthesis of naturally occurring quinones.
3. Gill, M. *Aust. J. Chem.* **1995**, *48*, 1.
4. Gill, M. *Nat. Prod. Rep.* **1994**, *11*, 67; *Nat. Prod. Rep.* **1996**, *13*, 513; *Nat. Prod. Rep.* **1999**, *16*, 301.
5. Gill, M.; Gimenez, A.; Watling, R. *J. Nat. Prod.* **1992**, *55*, 372.
6. See for example: Gill, M.; Gimenez, A.; Jhingran, A. G.; Milanovic, N. M; Palfreyman, A. *R. J. Chem. Soc., Perkin Trans. 1* **1998**, 3431; and references cited therein.
7. Gill, M.; Steglich, W. In *Progress in the Chemistry of Organic Natural Products*; Springer, Vienna 1987, Vol. 51, pp 1-317.
8. The code refers to the accession number under which voucher specimens are held in the herbarium of the Royal Botanic Garden, Edinburgh.
9. Gill, M.; Qureshi, A; Watling, R. *J. Nat. Prod.* **1992**, *55*, 517.
10. Gill, M.; Morgan, P. M. *J. Nat. Prod.* **1999**, *62*, 1298.
11. Thomson, R. H. In *Naturally Occurring Quinones*, 2nd Ed.; Academic Press: London, 1971.
12. Gill, M.; Gimenez, A. *J. Chem. Soc., Perkin Trans. 1* **1995**, 645.
13. Thomson, R. H. In *Naturally Occurring Quinones IV, Recent Advances*; Blackie: London, 1997.
14. Thomson, R. H. In *Naturally Occurring Quinones III, Recent Advances*; Chapman and Hall: London, 1987.
15. Savard, J.; Brassard, P. *Tetrahedron* **1984**, *40*, 3455.
16. Gill, M.; Gimenez, A.; Jhingran, A. G.; Palfreyman, A. R. *Tetrahedron: Asymmetry* **1990**, *1*, 621.
17. Billen, G.; Karl, U.; Scholl, K. D.; Stroeck, K. D; Steglich, W. In *Natural Products Chemistry III*; Springer: Berlin, Atta-ut-Rahman, Le Quesne, P. W. Eds.; 1988, pp 305-315.
18. Oertel, B., Dissertation, University of Bonn, 1984.
19. Cornforth, J. W. *Aust. J. Chem.* **1993**, *46*, 157.
20. Steglich, W.; Reininger, W. *Chem. Ber.* **1972**, *105*, 2922.
21. Keller, G.; Ammirati, J. F. *Mycotaxon* **1983**, *18*, 357.

22. Gill, M.; Gimenez, A.; Jhingran, A. G.; Smrdel, A. F. *Phytochemistry* **1989**, *28*, 2647.
23. Ayer, W. A.; Trifonov, L. S. *J. Nat. Prod.* **1994**, *57*, 317.
24. Shu, Y. -Z.; Arcuri, M.; Koslowski, M. R.; Wang, R. R.; Lam, K. S.; Chang, L. -P.; Pernik, D. *J. Antibiotics* **1994**, *47*, 1328.
25. Cameron, D. W.; Crossley, M. J. *Aust. J. Chem.* **1977**, *30*, 1161.
26. Banville, J.; Brassard, P. *J. Chem. Soc., Perkin Trans. 1* **1976**, 1852.
27. (a) Lobadie, R. P. *Pharm. Weekblad.* **1970**, *105*, 189 (b) Lobadie, R. P.; Otten, J.; Svendsen, A. B. *Pharm. Weekblad.* **1972**, *107*, 541.