

## Synthesis of the C3-C12 Fragment of Laulimalide

Hyo Won Lee\* and Joong-Yeoun Hong

Department of Chemistry, Chungbuk National University, 48 Gaeshin-dong, Cheongju, Chungbuk 361-763, Korea

Received September 26, 2003


**Key Words :** Antimitotic agent, Laulimalide, RCM

The antimiototic 20-membered macrolide laulimalide (**1**) also known as fijanolid B, isolated from sea sponges such as *Cacospongia mycofijiensis*,<sup>1</sup> *Hyatella* sp. *Spongia mycofijiensis*,<sup>2</sup> and *Fasciospongia rimosa*,<sup>3</sup> shows a potent biological activity against cell lines resistant to paclitaxel or epothilones since laulimalide binds to a site of microtubules in a mode distinct from taxoids.<sup>4</sup> The stabilization of polymerization by laulimalide leads to the inhibition of mitotic spindle and apoptosis of cells.

The structure of laulimalide determined by X-ray crystallographic study was reported in the literature.<sup>3a</sup> The remarkable biological activity as well as its structural features of laulimalide has inspired interests of organic chemists in its synthesis.<sup>5-10</sup> Previously we reported the synthesis of dihydropyran rings of laulimalide as a part of our synthetic effort toward laulimalide.<sup>11</sup> Herein we would like to report our synthetic efforts on the fragment C3-C12 of laulimalide.



Our retrosynthetic analysis of laulimalide (**1**) leads to two fragments of **2** and **3**. The subunit **3** can be prepared from compound **4** by exploiting the Wittig reaction on aldehyde. The dihydropyran ring of compound **4** can be provided by employing olefinic ring-closing metathesis (RCM). The subsequent transformation utilizing manipulations such as alkylation on chiral oxazolidinone and Horner-Wadsworth-Emmons reaction on side chains of **4** will provide subunit **3**.

At first, we thought the side chain at C-9 of the dihydropyran ring of **3** could be introduced by simple elongation of the side chain of compound **4**. (Eq. 1) However the nucleo-



philic substitution reaction of an enolate from (4*R*,5*S*)-4-methyl-5-phenyl-3-propionyl-oxazolidin-2-one with iodide **7** did not yield any desired product. Likewise the reaction with a carbanion of malonate ester showed a lack of reactivity of **7** toward nucleophiles. Even when we change the iodide to a mesylate, the result was same. This behavior may be attributed to the presence of dihydropyran ring, which might render the approach of nucleophiles unfeasible. Thus we decided not to convert a dihydropyran ring prior to the elongation of backbone. This became our synthetic strategy toward subunit **3**.

Using this reaction route, compound **5**<sup>11a</sup> was converted to iodide **7** by the reduction with LiAlH<sub>4</sub> followed by treatment with iodine and triphenylphosphine (Scheme 1). The conversion of **7** to carboxylic acid **8** of extended backbone was performed by decarboxylation of the intermediate of



**Scheme 1.** Reagents: a. LiAlH<sub>4</sub>, THF, 98%; b. PPh<sub>3</sub>, I<sub>2</sub>, imidazole, 98%; c. Diethyl malonate, NaH, DMF, 98%; d. LiI, collidine, 97%; e. PvCl, Et<sub>3</sub>N, THF; *n*-BuLi, (4*R*,5*S*)-4-Methyl-5-phenyl-2-oxazolidinone, 64%; f. NaHMDS, MeI, THF, 61%; g. Cl<sub>2</sub>(Cy<sub>3</sub>P)<sub>2</sub>Ru=CHPh, CH<sub>2</sub>Cl<sub>2</sub>, 90%; h. LiAlH<sub>4</sub>, THF, -78 °C, 85%; i. Ms<sub>2</sub>O, Et<sub>3</sub>N, CH<sub>2</sub>Cl<sub>2</sub>, 99%; LiBr, DMF, 40 °C, 94%; j. NaH, (EtO)<sub>2</sub>P(O)CH<sub>2</sub>CO<sub>2</sub>Et, DMSO, 50 °C, 6 h; CH<sub>2</sub>O, K<sub>2</sub>CO<sub>3</sub>, 60%; k. DIBAL, -78 °C, 98%; l. MOMCl, *i*Pr<sub>2</sub>NEt, 71%.

malonate ester using LiI. The resulting **8** was transformed to the derivative of oxazolidinone by the reaction of mixed anhydride with an enolate of chiral oxazolidinone. The methylation upon this intermediate gave compound **9**. Now with this compound **9** in hand, it is a time to introduce a dihydropyran ring by carrying out the RCM reaction using Grubbs catalyst. The resulting oxazolidinone **10** from RCM reaction of **9** was reduced to alcohol **11**. The next step along the sequence was the introduction of an  $\alpha,\beta$ -unsaturated ester group using the Horner-Wadsworth-Emmons reaction.<sup>12</sup> Thus, compound **11** was converted to a bromide intermediate *via* reaction of a mesylate with LiBr. Then, this bromide was transformed to a phosphate and subsequent *in-situ* reaction of this intermediate with formaldehyde provided the desired unsaturated ester **12**. The reduction of an ester group followed by protection with MOM group in due course finally provided the desired compound **13**.

In summary, we have reported the successful synthesis of the C3-C12 fragment of laulimalide using the introduction of a dihydropyran ring after the elongation of a side chain. Our on-going synthetic effort toward the synthesis of laulimalide will be reported in the near future.

**Acknowledgement.** This work was supported by Korea Science and Engineering Foundation (KOSEF-2000-1-123-002-5) and in part by BK21.

## References

1. Quinoa, E.; Kakao, Y.; Crews, P. *J. Org. Chem.* **1988**, *53*, 3642-3644.
2. Corley, D. G.; Herb, R.; Moore, R. E.; Scheuer, P. J.; Paul, V. J. *J. Org. Chem.* **1988**, *53*, 3644-3646.
3. (a) Jefford, C. W.; Bernardinelli, G.; Tanaka, J.; Higa, T. *Tetrahedron Lett.* **1996**, *37*, 159-162. (b) Tanaka, J.; Higa, T.; Bernardinelli, G.; Jefford, C. W. *Chem. Lett.* **1996**, *25*, 255-256.
4. Pryor, D. E.; O'Brate, A.; Bilcer, G.; Díaz, J. F.; Wang, Y.; Wang, Y.; Kabaki, M.; Jung, M. K.; Andreu, J. M.; Ghosh, A. K.; Giannakou, P.; Hamel, E. *Biochemistry* **2002**, *41*, 9109-9115.
5. Williams, D. R.; Mi, L.; Mullins, R. J.; Stites, R. E. *Tetrahedron Lett.* **2002**, *43*, 4841-4844.
6. Crimmins, M. T.; Stanton, M. G.; Allwein, S. P. *J. Am. Chem. Soc.* **2002**, *124*, 5958-5959.
7. (a) Mulzer, J.; Hanbauer, M. *Tetrahedron Lett.* **2002**, *43*, 3381-3383. (b) Mulzer, J.; Ohler, E. *Angew. Chem., Int. Ed.* **2001**, *40*, 3842-3846. (c) Enev, V. S.; Kaehlig, H.; Mulzer, J. *J. Am. Chem. Soc.* **2001**, *123*, 10764-10765.
8. Wender, P. A.; Hegde, S. G.; Hubbard, R. D.; Zhang, L. *J. Am. Chem. Soc.* **2002**, *124*, 4956-4957.
9. Ghosh, A. K.; Wang, Y.; Kim, J. T. *J. Org. Chem.* **2001**, *66*, 8973-8982.
10. Paterson, I.; De Savi, C.; Tudge, M. *Org. Lett.* **2001**, *3*, 3149-3152.
11. (a) Lee, H. W.; Jeong, C.-S.; Yoon, S. H.; Lee, I.-Y. C. *Bull. Korean Chem. Soc.* **2001**, *22*, 791-792. (b) Lee, H. W.; Yoon, S. H.; Lee, I.-Y. C.; Chung, B. Y. *Bull. Korean Chem. Soc.* **2001**, *22*, 1179-1180.
12. Vasil'ev, A. A.; Engman, L.; Serebryakov, E. P. *J. Chem. Soc., Perkin Trans. 1* **2000**, 2211-2216.