

- 1053 (1982).
16. D. M. Golden, R. K. Solly, and S. W. Benson, *J. Phys. Chem.* **75**, 1333 (1972).
17. D. C. McKean, J. L. Duncan, and R. K. M. Hay, *Spectrochim. Acta* **23A**, 605 (1966).
18. K. I. Lazaar and S. H. Bauer, *J. Phys. Chem.* **88**, 3052 (1984).
19. D. H. Borchardt and S. H. Bauer, *J. Chem. Phys.* **85**, 4980 (1986).
20. D. W. Octoby and S. A. Rice, *J. Chem. Phys.* **65**, 1676 (1976).
21. E. W. Schlag and R. D. Levine, *Chem. Phys. Lett.* **163**, 523 (1989).

## Association between Psoralens and Some Ionic Micelles

Sang Chul Shim\* and Young Hee Paik

Department of Chemistry, Korea Advanced Institute of Science and Technology, Seoul 130-650

Received April 2, 1990

The association between psoralens and some micelles is measured by the fluorescence quenching of psoralens by methylviologen ( $MV^{2+}$ ) and bromide ion in some ionic micellar solutions. The association constants were estimated to be  $\approx 10^4$  for all the psoralens studied even though they show different hydrophobicity.

### Introduction

Psoralens(furocoumarins), naturally occurring coumarin derivatives in various plants, have been widely used in the photochemotherapy of various skin diseases<sup>1,2</sup>. The photosensitivity of psoralens is primarily correlated with their photoreactivity toward pyrimidine bases in DNA under the 320–380 nm UV light<sup>3</sup>. Psoralens also cause damage to ribonucleic acids<sup>4</sup>, proteins<sup>5</sup>, and membranes<sup>6</sup>, and are used as molecular probe for nucleic acid structure<sup>7</sup>.

The fluorescence of psoralens in some ionic micellar solutions have been studied in our laboratory as models for biomimetic system<sup>8</sup>. The psoralen molecules are located in the micelle-water interfacial region and show different fluorescence features from that in aqueous solution due to both the low polarity and large viscosity of the micellar interface. The fluorescent behavior of psoralens in micellar solutions, therefore, can be used to determine their partition between the micellar and aqueous phases<sup>9</sup>.

In general, ionic quenchers localized in the aqueous phase fail to quench the fluorescence of substrates incorporated in the micellar phase when the ion and the micellar surface are like-charged and only the substrates in aqueous phase are quenchable. The quenching data, therefore, may be used to determine the association constants between substrates and micelles<sup>9-11</sup>.

In this paper, we would like to report the fluorescence quenching of 8-methoxypsoralen(8MOP), 5-methoxypsoralen(5MOP), 4,5',8-trimethylpsoralen(TMP), and 5,7-dimethoxycoumarin(DMC) by ionic quenchers,  $Br^-$  and methylviologen( $MV^{2+}$ ) in some micellar solutions and the association constants between psoralens and micelles are calculated.

### Experimental

**Materials.** 8MOP, 5MOP, and TMP were obtained from the Aldrich Chemical Company and used without fur-

ther purification. DMC was purified by recrystallization from ethanol. Sodium dodecyl sulfate(SDS,  $NaCH_3(CH_2)_{11}SO_4$ , CMC=8mM, Aldrich) was recrystallized three times from ethanol after washing with ether. Cetyltrimethylammonium bromide(CTAB,  $CH_3(CH_2)_{15}N(CH_3)_3Br$ , CMC=0.94mM, Aldrich) was recrystallized twice from methanol and CTAC(cetyltrimethylammonium chloride, CMC=1.4mM, Aldrich(25%)) was used as received. Methylviologen(1,1'-dimethyl-4,4'-bipyridinium chloride,  $MV^{2+}$ ) was prepared by the method reported<sup>12</sup>. Sodium bromide(Junsei Chemical Co.) and sodium chloride(Kanto Chemical Co.) were used as received. Chromatographic and spectroscopic grade solvents were used for high performance liquid chromatography and emission spectroscopy, respectively. Deionized distilled water was used.

**Methods.** Fluorescence spectra were recorded on an Aminco-Bowman spectrofluorometer with an Amino XY recorder and/or a Perkin-Elmer LS 50 spectrofluorometer at room temperature. To compare the hydrophobicity of psoralens, high performance liquid chromatography was performed on a Waters Associates Model 244 Liquid Chromatograph equipped with Model 6000A solvent delivery system, Model 440 UV absorbance detector(254nm), and Model U6K universal injector. The Lichrosorb RP-18 column and 60% methanol eluent were used. Fluorescence quenching experiments were carried out with  $MV^{2+}$ (0–4 mM) and  $Br^-$ (0–0.1 M) keeping the ionic strength at 0.1 by addition of sodium chloride which does not quench fluorescence. The contribution of surfactants and  $MV^{2+}$  to the ionic strength can be neglected<sup>13</sup>.


### Results and Discussion

**Fluorescence Quenching.** The addition of  $MV^{2+}$  and sodium bromide to the aqueous solutions of psoralens decreased the fluorescence intensity. The quenching data were analyzed by Stern-Volmer equation,

**Table 1.** Stern-Volmer Constants of Psoralen Fluorescence Quenching by MV<sup>2+</sup> and Br<sup>-</sup> in Aqueous and Micellar Solutions.

	$K_{sv}^a$					
	Br <sup>-</sup>				MV <sup>2+</sup>	
	H <sub>2</sub> O	SDS	CTAB	CTAC	H <sub>2</sub> O	SDS
8MOP	0 (0.25) <sup>b</sup>	0	4.0	2.9	0	1100
5MOP	0 (0.34) <sup>b</sup>	0	18	31		
TMP	0.64(1.2) <sup>b</sup>	0	3.1	4.5	80(140) <sup>b</sup>	470
DMC	20 (24) <sup>b</sup>	4.0	30	63	64(87) <sup>b</sup>	2600

<sup>a</sup> Measured at  $[Q] \rightarrow 0$  and ionic strength 0.1. <sup>b</sup> Measured without adjusting the ionic strength.




**Figure 1.** Stern-Volmer plot for fluorescence quenching of DMC (left) and 5MOP(right) by Br<sup>-</sup> in water(●), 10 mM SDS(○), 10mM CTAC(▲) and 5mM CTAB(□) solutions. In all the cases, the ionic strength was 0.1.

$$I_0/I = 1 + K_{sv}[Q] \quad (1)$$

where  $I_0$  and  $I$  denote the fluorescence intensities in the absence and presence of a quencher, respectively. The Stern-Volmer constant  $K_{sv}$  is related to the second order rate constant ( $k_q$ ) for the quenching and fluorescence lifetime ( $\tau_f$ ) of a fluorescing species by  $K_{sv} = k_q \tau_f$ .

The fluorescence quenching by MV<sup>2+</sup> and Br<sup>-</sup> showed linear Stern-Volmer relationships and  $K_{sv}$  values obtained from the slopes are summarized in Table 1. Since the lifetime of the 8MOP singlet state in aerated aqueous solution is 1.28ns<sup>17</sup>, the quenching rate constant ( $k_q$ ) of Br<sup>-</sup> is determined to be  $1.9 \times 10^8 \text{ M}^{-1}\text{s}^{-1}$ . From the lifetime of 7.2ns of DMC<sup>17</sup>, the  $k_q$  values of Br<sup>-</sup> and MV<sup>2+</sup> can be calculated as  $3.3 \times 10^9$  and  $1.2 \times 10^{10} \text{ M}^{-1}\text{s}^{-1}$ , respectively. The fluorescence quenching of 8MOP and 5MOP by MV<sup>2+</sup> was not observed under the experimental conditions ( $[\text{MV}^{2+}] < 4\text{mM}$ ).

The fluorescence quenching of psoralens by ionic quenchers, Br<sup>-</sup> and MV<sup>2+</sup>, are diffusion-controlled and it is greatly affected by the variation of the local concentration in micelles. The Stern-Volmer plots for 5MOP and DMC in micellar solutions are shown in Figure 1. The electrostatically attractive force between the cationic CTAB and CTAC micelle and Br<sup>-</sup> enhances the fluorescence quenching of psoralens embedded in the micellar interior (Stern region),



**Figure 2.** The variations of fluorescence intensity of 5MOP in CTAB(●) and CTAC(□) micellar solutions, plotted according to eq.(2).

whereas the repulsive interaction between the anionic micelle SDS and Br<sup>-</sup> decreases the probability of quenching reaction. Unlike the plots of data taken in aqueous solution, the Stern-Volmer plots in micellar solutions exhibited a large deviation from the linearity. The curvature in Stern-Volmer plot represents, largely, the dependence of the apparent bimolecular quenching constant,  $k_q = K_{sv}/\tau_f$ , on the concentration of quenchers, and mainly arises from the competition of psoralens and quenchers for available sites in the micelles. When the concentration of quencher is low, both psoralens and quenchers condense in the Stern layer of the micelle and the fluorescence quenching is very efficient. However, further addition of quenchers at given concentration of the micelle may displace some psoralens from the micellar phase, resulting in less efficient quenching. Such quenching behavior is frequently observed in the polyelectrolyte<sup>12,18</sup> and micellar solutions.<sup>19</sup> The  $K_{sv}$  values extrapolated to  $[Q]=0$ , i.e. slopes at  $[Q]=0$  are summarized in Table 1. For MV<sup>2+</sup> quencher, the quenching reaction is greatly enhanced in SDS micellar solution.

**Micelle-Psoralen Equilibrium Association Constants.** The fluorescence intensity of psoralens in aqueous solution is quite different from that in the micellar solutions<sup>8</sup>. If the fluorescence quantum yields of psoralens in the micelle and aqueous phase are independent of the surfactant concentration, the equilibrium association constant ( $K_{eq}$ ) for psoralens in the micellar solution can be calculated from the relationship<sup>9</sup>:


$$(I_\infty - I_w^0) / (I_t - I_w^0) = 1 + (K_{eq}[M])^{-1} \quad (2)$$

$I_\infty$ ,  $I_w^0$ , and  $I_t$  are relative emission intensity at 100% solubilization, without surfactant, and at intermediate surfactant concentration, respectively. The variation of fluorescence intensity of 5MOP in micellar solution reported earlier<sup>8</sup> are plotted according to eq.(2)(Figure 2) and the calculated  $K_{eq}$  are summarized in Table 2. The fluorescence intensity of psoralens in SDS solution varied very much in pre-micellar region and  $K_{eq}$  of psoralens in SDS solution could not be determined. By the same reason, that of TMP in CTAB and CTAC solutions could not be calculated.

**Table 2.** Association Constants Between Psoralen and Micelle( $K_{eq}$ ), and Distribution Coefficients Between the Micellar and Aqueous Phase( $a/b$ )

Psoralen	Surfactant	eq. used	$a/b$	$K_{eq}$
8MOP	CTAB	eq.(2)	3.1	$4.5 \times 10^4$
5MOP	CTAB	eq.(2)	3.1	$4.5 \times 10^4$
	CTAC	eq.(2)	6.6	$8.1 \times 10^4$
TMP	CTAB	eq.(5)	1.9*	
	CTAC	eq.(5)	2.9*	
DMC	SDS	eq.(3)	1.9-2.5*	$2.3 \times 10^4$ *
		eq.(5)	2.2*	$2.3 \times 10^4$ *
	CTAB	eq.(5)	1.6*	
		eq.(2)	2.8	$4.2 \times 10^4$


\*: Calculated at ionic strength 0.1


**Figure 3.** DMC fluorescence quenching by  $\text{Br}^-$  in 10mM SDS solution, plotted in the form given by eq. (3). The ionic strength was 0.1.

When a quencher and micellar system have the same sign of charges, the quencher impinging on the micelle surface is repelled instantaneously by the like-charged micelle head groups as a result of electrostatic repulsion and the quenching reaction with the solubilized psoralens in micellar phase rarely occurs. Hence, the quenching effects were attributed solely to the psoralens in the aqueous phase. From the selective quenching of the fluorescence arising from the aqueous phase, Quina and Toscano<sup>10</sup> evaluated the probe distribution between the micelle and aqueous phase by the equation (3)

$$\frac{\Phi_F^0}{\Phi_F^0 - \Phi_F} = \left\{ \frac{a}{b} \frac{\Phi_{Fm}^0}{\Phi_{Fw}^0} + 1 \right\} \left\{ 1 + \frac{1}{K_{sv} \tau_{Fw}^0 [Q]} \right\} \quad (3)$$

$\Phi_F$  is relative fluorescence quantum yield in the presence of the bulk solution quencher  $Q$ ,  $\Phi_F^0$  is the total fluorescence yield in the absence of a quencher,  $\Phi_{Fm}^0$  and  $\Phi_{Fw}^0$  are the fluorescence quantum yields in the micelle and aqueous phase, respectively.  $a$  and  $b$  are the fraction of probe in the micelle and aqueous phase. The ratio of  $a/b$  is related to the equilibrium constant,  $K_{eq}$ , for the association of the probe


**Figure 4.** Fluorescence quenching of DMC in 5 mM CTAB( $\blacktriangle$ ), TMP in 5mM CTAB( $\circ$ ) and 10mM CTAC( $\blacksquare$ ) solutions by  $\text{MV}^{2+}$ , plotted in the form given by eq.(5). In all the cases, 0.1M NaCl was added.

with the micelle by eq. (4).

$$K_{eq} = \frac{a}{b} \left\{ \frac{[D]_t - \text{CMC}}{N} - a[Ps]_t \right\}^{-1} = \frac{a}{b} \left\{ \frac{[D]_t - \text{CMC}}{N} \right\}^{-1} \quad (4)$$

The intercept and slope of the plot of  $\Phi_F^0 / (\Phi_F^0 - \Phi_F)$  vs.  $[Q]^{-1}$  can be used to calculate  $a/b$ . The plot of the experimental data according to eq.(3) for the fluorescence quenching of DMC by  $\text{Br}^-$  is shown in Figure 3. The aggregation number,  $N$ , for SDS in the presence of 0.1 M NaCl is 88<sup>20</sup> and the ratio of  $I_{Fm}^0 / I_{Fw}^0$  for DMC was measured to be 1.5. The equilibrium constant  $K_{eq}$  for DMC in SDS based on eq.(3) and eq.(4) is  $2-2.8 \times 10^4 \text{ M}^{-1}$ . The ratio of intercept/slope ( $K_{sv}$ ) is in reasonable agreement with the value obtained from the Stern-Volmer plot in aqueous solution. Abuin and Lissi<sup>11</sup> have proposed a modified equation(5):

$$(I_0 - I) / I_0 = \alpha K_{sv} [Q] / (1 + K_{sv} [Q]) \quad (5)$$


where  $\alpha$  is the fraction of light emitted from the aqueous phase at  $[Q]=0$ . The fraction of probe in the aqueous phase can be obtained by eq. (6),

$$\% \text{ in the water phase} = 100 \alpha I_{Fm}^0 / I_{Fw}^0 \quad (6)$$

The calculated  $K_{eq}$  for DMC in SDS micelle solution from eq.(5) and eq.(6) ( $2.3 \times 10^4$ ) agreed well with the value from eq. (4). The quenching data of TMP and DMC by  $\text{MV}^{2+}$  in CTAB and CTAC micellar solutions are plotted according to eq.(5) and are shown in Figure 4.

The HPLC experiments were performed(Figure 5) to measure the relative hydrophobicity of psoralens. The solubility of 8MOP, 5MOP, and TMP in water is 23, 5, and 1  $\mu\text{g/ml}$ , respectively. From the solubility and retention time of HPLC, the order of relative hydrophobicity are expected to be  $\text{DMC} < 8\text{MOP} < 5\text{MOP} < \text{TMP}$ . The binding abilities of psoralens to the micelles with hydrophobic interaction are little different in spite of the differences in hydrophobicity and are affected little by the addition of salts.

Column: Lichrosorb RP-18  
Solvent: 60% Methanol



**Figure 5.** HPLC profile of 8MOP, 5MOP, and TMP.

**Acknowledgements.** This investigation was supported by the Korea Science & Engineering Foundation and the Korea Advanced Institute of Science & Technology. We thank professor J.W. Park for the gift of methylviologen.

### References

- J. A. Parrish, T. B. Fitzpatrick, L. Tanenbaum, and M. A. Pathak, *New Eng. J. Med.*, **291**, 1207 (1974).
- J. A. Elliot, Jr., *J. Invest. Dermatol.*, **32**, 311 (1959).
- S. T. Isaacs, C.-K. J. Shen, J. E. Hearst, and H. Rapoport, *Biochemistry*, **16**, 1058 (1977).
- K. Yoshikawa, N. Mori, S. Saskakibara, N. Mizuno, and P. S. Song, *Photochem. Photobiol.*, **29**, 1127 (1979).
- A. Y. Potapenko, M. V. Moshinin, A. A. Krasnowski, and V. L. Sukhoorukov, *Z. Naturf.*, **37C**, 70 (1982).
- P. S. Song and C. N. Ou, *Ann. N. Y. Aca. Sci.*, **346**, 355 (1980).
- M. Sasaki, T. Sadaka, and M. Sukigara, *Chem. Lett.*, **701** (1977).
- S. C. Shim and Y. H. Paik, submitted for publication.
- M. Almgren, F. Grieser, and J. K. Thomas, *J. Am. Chem. Soc.*, **101**, 279 (1979).
- F. H. Quina and V. G. Toscano, *J. Phys. Chem.*, **81**, 1750 (1977).
- D. Stigter, *J. Collid. Interface Sci.*, **47**, 473 (1974).
- J. W. Park and Y. H. Paik, *Bull. Kor. Chem. Soc.*, **6**, 287 (1985).
- Taking the CMC of SDS to be 1.4 mM<sup>14</sup> and the degree of micellar ionization to be about 0.2<sup>15</sup> at the ionic strength 0.1, the concentration of the free sodium ion due to the detergent is 3 mM which is considerably smaller than 0.1. Because the CMC of CTAB and CTAC at the ionic strength 0.1 is smaller than 0.95 and 1.4 mM reported in the absence of salts<sup>16</sup>, respectively, the contribution of these surfactants to the ionic strength can be neglected.
- M. F. Emerson and A. Holtzer, *J. Phys. Chem.*, **71**, 1898 (1967).
- L. Romsted, Ph.D. Thesis, Indiana University, Bloomington, Ind, 1975.
- Ch. Tanford, *The Hydrophobic Effect: Formation of Micelles and Biological Membrane*, Wiley-Interscience, New York, 1973, Chapter 7.
- S. C. Shim, H. K. Kang, S. K. Park, and E. J. Shin, *J. Photochem.*, **37**, 125 (1987).
- C. D. Janah, M. S. Matheson, and D. Meisel, *J. Phys. Chem.*, **83**, 257 (1979).
- D. Meisel, M. S. Matheson, and J. Rabani, *J. Am. Chem. Soc.*, **100**, 117 (1978).
- M. Sasaki, I. Nakasato, H. Sugiura, H. Fujita, and T. Sakata, *Photochem. Photobiol.*, **46**, 551 (1987).

## New Crown Compounds Derived from 1,2-Bis(2-hydroxybenzyl)benzene (II): Bisaryl Crowns

Woo Young Lee\*, Jae-Do Jung, Chang Hee Park, and Wonbo Sim

Department of Chemistry, Seoul National University, Seoul 151-742.

Oee Sook Park

Department of Chemistry, Chungbuk National University, Chungbuk 360-763 Received April 3, 1990

New bisaryl corands (crown ethers) bearing 1,2-dibenzyl- and 1,2-dibenzoylbenzene subunits have been synthesized: The reaction of 1,2-bis(2-hydroxybenzyl)benzene in base with mono-tetrahydropyranyl oligoethylene glycol tosylate, deprotection of the bis-condensation product to give a corresponding diol, tosylation of the free hydroxyls of the diol, and condensation of the ditosylate in base with 1,2-bis(2-hydroxybenzyl)benzene afforded a new type of bisaryl corand(I) of 1,2-dibenzylbenzene system. Oxidation of the benzylic positions of the corands(I) furnished novel aromatic corands(II) containing partly carbonyl functions.

### Introduction

Various bisaryl crown ethers, such as 1, 2, 3 and 4, have been synthesized by many workers, in which two aromatic hydrocarbon subunits are linked symmetrically or unsymmetrically by two ethyleneoxy chains to form a macrocycle.

Although the crowns 1-4 are structurally analogous, their complexing properties vary markedly with the structures of the aromatic subunits and the polyether ring sizes. Dibenzo-18-crown-6(1), synthesized by C. J. Pederson in his early investigations for crown ether synthesis,<sup>1,2</sup> and reasonably selective toward the complexation of potassium ions, is the first