

Nucleoside Recognition by a Fluorescent Macrolactam

Hae-Jo Kim* and Jong-In Hong†,*

Department of Chemistry, College of Natural Sciences, Kyonggi University, Suwon 443-760, Korea. *E-mail: haejkim@kgu.ac.kr

†Department of Chemistry, College of Natural Sciences, Seoul National University, Seoul 151-747, Korea

*E-mail: jihong@snu.ac.kr

Received August 16, 2007

Key Words : Fluorescence, Macrolactam, Nucleoside, Recognition

Molecular recognition of nucleosides or nucleotides is attracting a great deal of interest due to their genetic functions in living organisms.¹ Hydrophilic nature of nucleosides and nucleotides allows only a conformationally well defined receptor to form a hydrogen-bonded, electrostatic or hydrophobic complex with nucleosides or nucleotides in water.² Recently, an anthracene derivative was reported to show a higher affinity toward GTP over ATP owing to cooperative interactions of hydrogen bonding and electrostatic interactions between an imidazolium moiety and a phosphate unit.³

We have developed various sugar receptors with hydrogen-bonding acceptors and donors.⁴ Herein, we report a novel *D*₂-symmetric fluorescent macrolactam. This host possesses not only an aromatic cavity for π - π interaction, but also hydrogen-bonding donors/acceptors in the peripheral site of the macrolactam for effective nucleoside recognition.

Macrolactam host was synthesized via the typical acid chloride coupling method⁵ in which 2,5-dimethyl-*p*-xylyl-diamine was treated with 2,5-dimethoxyterephthaloyl chloride in a high dilute condition⁶ to afford the desired 2:2 macrocyclization product (**H**). The calculated structure shows that the host has a large cavity with dimension of 10.5 Å × 6.9 Å (Fig. 1. left). The global minimum structure clearly indicates that π - π stacking interaction exists between the dimethoxy aryl groups of **H** and the uracil base of uridine with aromatic-aromatic surface distances of 3.56 and 3.55 Å, and one intermolecular H-bonding interaction also exists between the carbonyl group of **H** and 2'-OH group of uridine (Fig. 1. right).⁷

Owing to the characteristic fluorescence property of **H**,⁸ fluorescence titration was carried out in chloroform. Fluorescence emission intensities at $\lambda_{\text{max}} = 384$ nm were recorded after excitation at $\lambda_{\text{ex}} = 331$ nm (Fig. 2). Fluorescence intensities of the host-guest complex increase upon addition of sugars or nucleosides presumably due to the restricted rotation of **H**.⁹ The resulting fluorescence enhancements at 384 nm are shown in the inset of Figure 3. The binding stoichiometry between **H** and guests was also confirmed to be 1:1 by Job's plot (Fig. 4).¹⁰

Curve fitting of the host signals to a 1:1 binding isotherm gives apparent dissociation constants of up to $K_d = 10^{-4}$ M, which are summarized in Table 1.

While the dissociation constants between **H** and anomers of D-glucose were found to be similar (3.99×10^{-4} M for β




Figure 2. UV-vis and fluorescence spectra of **H** in CHCl_3 at 298 K.




Figure 1. Global minimum structures of **H** (left) and its uridine complex (right).




Figure 3. Fluorescence titration of **H** and uridine in CHCl_3 at 298 K. $[\mathbf{H}] = 2.0 \mu\text{M}$.




Figure 4. Job's plot between **H** and D-glucopyranosides at 298 K. $[\mathbf{H}] + [\mathbf{G}] = 2.0 \mu\text{M}$, each in 2.0 mL. Rectangular and circle represent β -glucose and uridine, respectively.

and $5.38 \times 10^{-4} \text{ M}$ for α anomer), the binding affinity of **H** to β -galactose is three times lower than that of β -D-glucose ($1.31 \times 10^{-3} \text{ M}$ for β -D-galactose). This diastereoselectivity for sugars plausibly results from the slight energetic difference in the intermolecular H-bonding patterns due to the varying degree of steric interaction between sugars and **H**. This indicates that geometrical complementarities of H-bonding partners are crucial in hydrogen bond-based molecular recognition system.

It is noticeable that nucleosides, deoxythymidine (d-Thy) and uridine (Uri) show the comparable binding affinities although they have fewer number of hydroxyl groups compared with the pyranosides. Uridine shows much higher binding affinity ($1.72 \times 10^{-4} \text{ M}$) than β -D-glucose. Enhancement in the binding affinity for nucleosides probably results from the presence of π -surface and H-bonding donors and acceptors in the guests.

It is assumed that π - π stacking interaction between **H** and nucleosides plays an important role in host-guest binding. We have chosen several commercially available aromatic guests to test this assumption. While benzene is weakly bound to **H** ($K_d = 4.05 \times 10^{-2} \text{ M}$), the binding affinity of a π

Table 1. Dissociation constants between **H** and guests^a

entry	guest structure	name	K_d (M)
1		β -D-Glucose	$3.99(\pm 0.70) \times 10^{-4}$
2		α -D-Glucose	$5.38(\pm 3.22) \times 10^{-4}$
3		β -D-Galactose	$1.31(\pm 0.58) \times 10^{-3}$
4		Thymidine	$6.44(\pm 5.88) \times 10^{-4}$
5		Uridine	$1.72(\pm 0.23) \times 10^{-4}$


^aFluorescence titration of constant host concentration ($2.0 \mu\text{M}$) in CHCl_3 at 298 K. Fluorescence intensity at $\lambda_{\text{em}} = 384 \text{ nm}$ ($\lambda_{\text{ex}} = 331 \text{ nm}$) was monitored after each addition of guest.

basic guest 1,4-dimethoxybenzene was *c.a.* hundred times enhanced ($5.26 \times 10^{-4} \text{ M}$). The binding affinity of a π acidic guest dimethylterephthalate, however, was too small to determine.

In conclusion, we have developed a novel fluorescent macrolactam as an artificial receptor for nucleosides. The receptor has shown high diastereoselectivity for sugars and even higher affinities for nucleosides due to the intermolecular π - π stacking interaction as well as H-bonds between the macrolactam and sugars/nucleosides.

Experimental

Acid chloride synthesis. To a solution of 400 mg (1.77 mmol) of 2,5-dimethoxyterephthalic acid in 20 mL of dichloromethane was added cat. amount of DMF and 2.0 mL of 2 M oxalic acid chloride in dichloromethane (2 eq. ex., 4.0 mmol). Resulting white suspension was stirred at rt



Scheme 1. Synthetic scheme of macrolactam.

under nitrogen for 5 hrs to afford a yellow clear solution. All volatiles were removed under the reduced pressure, dried in vacuum.

Cyclization. To a solution of p-xylyl diamine (1 eq. 1.77 mmol) and TEA (2 eq ex.) in 500 mL of dichloromethane was dropwise added a solution of above crude 2,5-dimethoxyterephthaloyl chloride in 50 mL of dichloromethane at 0 °C under nitrogen over a period of 2 hrs. Resulting yellow solution was stirred for additional 24 hrs under nitrogen. All volatiles are removed under reduced pressure and purified by column chromatography. Column chromatography on silica gel (CH₂Cl₂:MeOH = 10:1, *R_f* = 0.48) gave a greenish mixture. Additional column chromatography on silica gel (EtOAc, *R_f* = 0.30) gave the desired product, **H** as a white solid in a 4.2% yield.

¹H-NMR (300 MHz, CDCl₃): 8.09 (t, *J* = 6.3 Hz, 4H of NH), 7.68 (s, 4H of ArH_a in 2,5-dimethoxybenzene), 7.06 (s, 4H of ArH in *p*-xylylene), 4.51 (d, *J* = 6.3 Hz, 8H of ArCH₂N), 3.90 (s, 12H of OCH₃), 2.29 (s, 12H of ArCH₃).

UV-vis (CHCl₃): ε_{331nm} = 3997 M⁻¹cm⁻¹, Fluorescence (CHCl₃): λ_{em} = 384 nm (λ_{ex} = 331 nm) in 2.0 μM

Mass (FAB⁺, m-NBA): *m/z* 709 ([M+H], 50%)

Acknowledgement. Financial support from Korea Research Foundation (KRF-2006-312-C00592) is greatly acknowledged.

References

- (a) Yang, F.; Belitsky, J. M.; Villanueva, R. A.; Dervan, P. B.; Roth, M. J. *Biochemistry* **2003**, *42*, 6249. (b) Gearhart, M. D.; Dickinson, L.; Ehley, J.; Melander, C.; Dervan, P. B.; Wright, P. E.; Gottesfeld, J. M. *Biochemistry* **2005**, *44*, 4196.
- (a) Urbach, A. R.; Dervan, P. B. *Proc. Nat'l Aca. Sci.* **2001**, *98*, 4343. (b) Chenoweth, D. M.; Viger, A.; Dervan, P. B. *J. Am. Chem. Soc.* **2007**, *129*, 2216.
- Kwon, J. Y.; Singh, N. J.; Kim, H. N.; Kim, S. K.; Kim, K. S.; Yoon, J. J. *J. Am. Chem. Soc.* **2004**, *126*, 8892.
- (a) Cho, H. -K.; Kim, H. -J.; Lee, K. H.; Hong, J.-I. *Bull. Korean Chem. Soc.* **2004**, *25*, 1714. (b) Lee, D. H.; Kim, H.-J.; Hong, J.-I. *Supram. Chem.* **2007**, *19*, 251.
- Kim, H.-J.; Kim, Y.-H.; Hong, J.-I. *Tetrahedron Lett.* **2001**, *42*, 5049.
- Kim, T. W.; Hong, J.-I. *Bull. Korean Chem. Soc.* **1995**, *16*, 781.
- Conformation searches were performed with the program MacroModel 7.0, Amber* force field in chloroform solvent: Mohamadi, F.; Richards, N. G. J.; Guida, W. C.; Liskamp, R.; Lipton, M.; Caufield, C.; Chang, G.; Hendrickson, T.; Still, W. C. *J. Comput. Chem.* **1990**, *11*, 440.
- (a) Jang, Y. J.; Jun, J. H.; Swamy, K. M. K.; Nakamura, K.; Koh, H. S.; Yoon, Y. J.; Yoon, J. *Bull. Korean Chem. Soc.* **2005**, *26*, 2041. (b) An, B.-K.; Kwon, S.-K.; Park, S. Y. *Bull. Korean Chem. Soc.* **2005**, *26*, 1555.
- (a) Lee, D. H.; Im, J. H.; Lee, J.-H.; Hong, J.-I. *Tetrahedron Lett.* **2002**, *43*, 9637. (b) Mcfarland, S. A.; Finney, N. S. *J. Am. Chem. Soc.* **2002**, *124*, 1178.
- Job, P. *Ann. Chim.* **1928**, *9*, 113.