

Revisiting the Water Binding of Small Cavitands: The Role of Benzene Hydrogen Bonding

Chaesang Ihm, Seungjoo Cho,[†] and Kyungsoo Paek^{*}

Department of Chemistry, Soongsil University, Seoul 156-743, Korea. *E-mail: kpaek@ssu.ac.kr
[†]Computational Science Center, Korea Institute of Science and Technology, Seoul 130-650, Korea
 Received August 16, 2007

Key Words : Cavitand, Water, Anion recognition, Benzene hydrogen bonding

The solvent of biological systems, water, is hard to study in molecular recognition field due to its high polarity and small size. The strong hydrogen bonding interaction of water with themselves or other polar molecules makes it very hard to isolate and observe a single molecule in solution.

Small cavitands have been designed to accommodate small molecules such as alcohols.¹ Recently, we have reported the synthesis and water-specific binding property of C_{2v} cavitands **1-3** at low temperature.²


The CPK molecular model study as well as the experimental ¹H NMR study showed that these C_{2v} cavitands can not accommodate MeOH, EtOH, CH₃CN, CH₃CHO, CH₃NO₂, CH₄, or NH₄⁺, but can accommodate H₂O due to its small cavity partially blocked by protons H_a of the bridging resorcinolic units (Figure 1(a)). When the temperature of the

solutions of cavitands **1**, **2**, and **3** in water-saturated CD₂Cl₂ or CDCl₃ decreased to -70 °C, the intensity of free H₂O peak was decreased due to water-freezing. On the contrary, the peak intensity of complexed water was unchanged or even slightly increased. This phenomenon implies that the energy barrier for decomplexation of H₂O is substantially high compared to that for complexation at -70 °C due to the solvophobic driving force.

The distinct peaks of free and complexed guests enable the direct calculation of K_a , which decreased in this order: cavitand **3** > cavitand **1** > cavitand **2**. This phenomena was explained as follows; A COOCH₃ group of host **3** increased the hydrophilicity to the gate of its cavity and H₂O molecules could be easily gathered and enter through the gate better than those of host **1** or **2**.²

Recently molecular recognitions based on sp² C-H...anion interactions have been reported.³ As the molecular mechanics-optimized structure of H₂O@**1** (Fig. 1(b) and (c)) shows, the hydrogen bonding interactions between two C-H_a and OH₂ seems to be significant in the water-specific binding of these cavitands. To observe the effects of benzene hydrogen bonding various C_{2v} cavitands with *p*-X group to H_a hydrogen were synthesized and studied.

To increase the solubility of C_{2v} cavitands by incorporating long alkyl feet the synthetic route shown in Scheme 2 was followed (Scheme 2).⁴ From tetrabromide **11**, tetrakisester **12**, tetrakis(hydroxymethyl)cavitand **13**, and then tetrakis(bromomethyl)cavitand **14** were obtained subsequently. The coupling of tetrakis(bromomethyl)cavitand **14** with 5-X-resorcinol (X = substituent) in K₂CO₃/DMF



Scheme 1. C_{2v} cavitands.





Figure 1. (a) X-ray crystal structure of cavitand **1**, (b), (c) Molecular mechanics-optimized structure of H₂O@**1** (CFF95 Force-field by Cerius2[®]) (b: top view, c: side view (the front wall was removed for clarity)).



Scheme 2. Synthesis of C_{2v} cavitands 4-10 (R = (CH₂)₆CH₃).

gave cavitands 4-10 in good yields (22%-35%). These C_{2v} cavitands have been fully characterized by NMR, FAB+ Mass spectra and elemental analyses.

Figure 2 shows the ¹H NMR spectra of cavitant 7 without or with 10 eq of NBu₄⁺ X⁻ in CDCl₃ at 298 K. The chemical shifts changes upon anion complexation of H_a and H_b from 5.21 and 4.41 ppm for free 7, respectively, imply the qualitative interactions between these protons and anions (H_a and H_b are illustrated in Scheme 1). The patterns of chemical shifts of these two protons according to the anions are similar; F⁻ (5.29, 4.43 ppm) ~ CN⁻ (5.32, 4.42 ppm) > Cl⁻ (5.27, 4.42 ppm) >> Br⁻ (5.21, 4.40 ppm) ≈ I⁻ (5.21, 4.40 ppm). These results show that F⁻ and CN⁻ best fit to the cavity and Cl⁻ fits rather improperly, but Br⁻ and I⁻ cannot fit the cavity due to their large size. But the quantitative evaluations of thermodynamic parameters for anion complexation were difficult due to the interference by water.

Table 1 shows the association constants and binding energies for H₂O@cavitant by ¹H NMR experiment and *ab*




Figure 2. ¹H NMR spectra of cavitant 7 with or without 10 eq of NBu₄⁺ X⁻ in CDCl₃ (298 K, 400 MHz). (a) free 7, (b) NBu₄⁺F⁻@7, (c) NBu₄⁺Cl⁻@7, (d) NBu₄⁺Br⁻@7, (e) NBu₄⁺I⁻@7, (f) NBu₄⁺CN⁻@7.

initio calculations. As expected the affinity of water complexation mainly depends on the nature of substituent (X) of cavitant. Cavitant 6 (X = H) gave K_a (M⁻¹) = 103 and cavitands 7-10 with the better electron withdrawing group compared to H gave K_a (M⁻¹) = 211-379. But cavitands 4 and 5 with electron donating X group gave K_a (M⁻¹) = 76 and 74, respectively. These results strongly support that the hydrogen bonding interaction between sp² C-H...OH₂ plays an important role in the water complexation of these C_{2v} cavitands. The calculated binding energies in the gas phase were consistent with the experimental results in solution, *i.e.*, with electron-withdrawing substituents, the binding energy becomes stronger, while with electron-donating ones the energy becomes smaller.

In conclusion, these C_{2v} container hosts 4-10 showed the specific binding properties for H₂O in CDCl₃ at low temperature due to their complementarity to water, the solvophobic interaction of water, and the hydrogen bonding interaction

Table 1. The association constants (K_a) and binding energies (-ΔG°) for H₂O@cavitant

Cavitant	K _a (M ⁻¹) ^a		-ΔG° (kcal mol ⁻¹)	
	Exp. ^b	Calcd ^c	Exp. ^b	Calcd ^c
4	76		1.9	9.1
5 (2)	74(61)		1.9(1.8)	9.3
6 (1)	103(71)		2.0(1.9)	9.5
7 (3)	230(137)		2.4(2.2)	10.0
8	232		2.4	9.9
9	211		2.2	10.0
10	379		2.6	10.0
15 (X = NO ₂)	-		-	10.5

^aestimated Error ± 10%. ^b-50 °C in water-saturated CD₂Cl₂ (1-3) or CDCl₃ (4-10) using 400-MHz ¹H NMR Spectrometer. ^c*ab initio* calculations using B3LYP/6-31G** in the gas phase

between sp^2 C-H and O-H₂. The presence of hydrogen bonding interaction between sp^2 C-H and O-H₂ was confirmed by observing the relationships between their binding strength and the electronic effect of X groups.

Experimental

General coupling reaction between tetrakis(bromo-methyl)cavitand 14 and 5-X-resorcinol; Tetrakis(bromo-methyl)cavitand **14** (500 mg, 0.38 mmol) and bridging reagent (0.77 mmol) were dissolved in 20 mL of dry, degassed DMF. This solution was added dropwise over 1 hour to a stirred mixture of 30 mL of DMF and 1.07 g of K₂CO₃ heated to 60 °C. After 1 day, the reaction mixture was cooled to room temperature. K₂CO₃ was filtered and washed with CH₂Cl₂ (20 mL). The filtrate was partitioned between CH₂Cl₂ (30 mL) and 3 N HCl (100 mL). The organic phase was washed with H₂O, brine, and then dried over MgSO₄. The organic phase was filtered and concentrated under reduced pressure. The residue was purified by silica gel column chromatography and the concentrate of the best portions was poured into a solvent to give a precipitate of pure cavitand.

Cavitand 4: 5-Methoxyresorcinol (107 mg) was used as a bridging reagent and a mixture of hexane:EtOAc = 8:1 was used as eluent for column chromatography. EtOH was used as a recrystallization solvent to give pure cavitand **4** as a white solid (108 mg, 22%). mp >300 °C (dec). MALDI-TOF: m/z 1319 [(M + Na + NaOH)⁺ 30%], 1280 [(M + Na)⁺ 100%]. Elemental analysis calcd for C₇₈H₉₆O₁₄: C, 74.49; H, 7.69. found: C, 74.12; H, 7.77. ¹H NMR (400 MHz, CDCl₃, 25 °C): δ = 7.16 (s, 4H, ArH), 6.39 (d, 4H, *J* = 2.0 Hz, α to OCH₃), 5.94 (d, 2H, *J* = 8.0, noncyclic outer OCH₂O), 5.42 (d, 4H, *J* = 8.0, outer ArCH₂O), 5.05 (m, 4H, noncyclic methine + cyclic outer OCH₂O), 4.63 (t, 2H, *J* = 8.0, cyclic methine), 4.55 (br-s, 2H, γ to OCH₃), 4.38 (m, 6H, noncyclic inner OCH₂O + inner ArCH₂O), 3.76 (s, 6H, OCH₃), 3.53 (d, 2H, *J* = 6.8, cyclic inner OCH₂O), 2.34, 2.13 (m, 8H CH₂(CH₂)₅CH₃), 1.52-1.27 (m, 40H, CH₂(CH₂)₅CH₃), 0.91 (m, 12H, CH₂(CH₂)₅CH₃). ¹³C NMR (CDCl₃, 100 MHz, 25 °C): δ = 163.38, 162.14, 155.79, 150.58, 139.93, 138.10, 125.19, 120.57, 111.76, 105.88, 102.54, 96.42, 70.50, 55.95, 37.47, 37.32, 32.29, 32.25, 31.43, 30.29, 30.15, 30.11, 29.88, 29.71, 28.48, 28.40, 28.27, 23.11, 23.07, 14.53, 14.51.

Cavitand 5: Orcinol (96 mg) was used as a bridging reagent and a mixture of hexane:CH₂Cl₂ = 1:5 was used as eluent for column chromatography. CH₃CN was used as a recrystallization solvent to give pure cavitand **5** as a white solid (130 mg, 28%). mp > 300 °C (dec). MALDI-TOF: m/z 1288 [(M + Na + NaOH)⁺ 30%], 1247 [(M + Na)⁺ 100%]. Elemental analysis calcd for C₇₈H₉₆O₁₂: C, 76.44; H, 7.90. found: C, 75.85; H, 7.95. ¹H NMR (400 MHz, CDCl₃, 25 °C): δ = 7.16 (s, 4H, ArH), 6.65 (s, 4H, α to CH₃), 5.96 (d, 2H, *J* = 8.0, noncyclic outer OCH₂O), 5.42 (d, 4H, *J* = 12.0, outer ArCH₂O), 5.07 (t, *J* = 8.0, 2H, noncyclic methine), 4.93 (d, 2H, *J* = 8.0, cyclic outer OCH₂O), 4.73 (br-s, 2H, γ

to OCH₃), 4.63 (t, 2H, *J* = 8.0, cyclic methine), 4.42 (d, 2H, *J* = 8.0, noncyclic inner OCH₂O), 4.33 (d, 4H, *J* = 8.0, inner ArCH₂O), 3.42 (d, 2H, *J* = 4.0, cyclic inner OCH₂O), 2.29 (s, 6H, CH₃), 2.34, 2.13 (m, 8H CH₂(CH₂)₅CH₃), 1.52-1.24 (m, 40H, CH₂(CH₂)₅CH₃), 0.89 (m, 12H, CH₂(CH₂)₅CH₃). ¹³C NMR (CDCl₃, 100 MHz, 25 °C): δ = 161.29, 155.83, 150.56, 143.43, 139.92, 138.10, 125.22, 120.50, 120.47, 116.13, 102.53, 96.45, 70.51, 37.46, 37.33, 32.28, 32.25, 31.42, 30.29, 30.15, 29.88, 29.78, 29.71, 28.48, 28.40, 28.27, 23.10, 23.06, 21.73, 14.53.

Cavitand 6: Resorcinol (85 mg) was used as a bridging reagent and a mixture of hexane:CH₂Cl₂ = 1:2 was used as eluent for column chromatography. EtOH was used as a recrystallization solvent to give pure cavitand **6** as a white solid (162 mg, 35%). mp > 300 °C (dec). MALDI-TOF: m/z 1261 [(M + Na + NaOH)⁺ 30%], 1221 [(M + Na)⁺ 100%]. Elemental analysis calcd for C₇₆H₉₂O₁₂: C, 76.22; H, 7.74. found: C, 75.67; H, 7.78. ¹H NMR (400 MHz, CDCl₃, 25 °C): δ = 7.38 (t, 2H, *J* = 8.0 resorcinol) 7.17 (s, 4H, ArH), 6.84 (d, 4H, *J* = 2.4, resorcinol), 5.98 (d, 2H, *J* = 8.0, noncyclic outer OCH₂O), 5.45 (d, 4H, *J* = 12.0, outer ArCH₂O), 5.05 (t, *J* = 8.0, 2H, noncyclic methine), 4.97 (br-s, 2H, resorcinol), 4.91 (d, 2H, *J* = 8.0, cyclic outer OCH₂O), 4.66 (t, 2H, *J* = 8.0, cyclic methine), 4.44 (d, 2H, *J* = 8.0, noncyclic inner OCH₂O), 4.34 (d, 4H, *J* = 12.0, inner ArCH₂O), 3.32 (d, 2H, *J* = 8.0, cyclic inner OCH₂O), 2.34, 2.13 (m, 8H CH₂(CH₂)₅CH₃), 1.52-1.24 (m, 40H, CH₂(CH₂)₅CH₃), 0.89 (m, 12H, CH₂(CH₂)₅CH₃). ¹³C NMR (CDCl₃, 100 MHz, 25 °C): δ = 161.63, 155.90, 155.17, 150.63, 140.03, 138.12, 132.88, 125.16, 120.53, 119.79, 119.41, 102.45, 96.39, 70.71, 37.49, 37.38, 32.30, 32.27, 31.44, 31.21, 30.30, 29.89, 29.72, 28.66, 28.49, 28.28, 23.11, 23.08, 14.55.

Cavitand 7: Methyl 3,5-dihydroxybenzoate (129 mg) was used as a bridging reagent and a mixture of Hexane:EtOAc = 8:1 was used as eluent for column chromatography. EtOH was used as a recrystallization solvent to give pure cavitand **7** as a white solid (128 mg, 25%). mp >300 °C (dec). MALDI-TOF: m/z 1377 [(M + Na + NaOH)⁺ 30%], 1337 [(M + Na)⁺ 100%]. Elemental analysis calcd for C₈₀H₉₆O₁₆: C, 73.15; H, 7.37. found: C, 72.55; H, 7.38. ¹H NMR (400 MHz, CDCl₃, 25 °C): δ = 7.50 (s, 4H, α to COOCH₃) 7.18 (s, 4H, ArH), 5.98 (d, 2H, *J* = 8.0, noncyclic outer OCH₂O), 5.50 (d, 4H, *J* = 12.0, outer ArCH₂O), 5.21 (br-s, 2H, γ to COOCH₃), 5.07 (t, *J* = 8.0, 2H, noncyclic methine), 4.95 (d, 2H, *J* = 4.0, cyclic outer OCH₂O), 4.65 (t, 2H, *J* = 8.0, cyclic methine), 4.38 (m, 6H, noncyclic inner OCH₂O + inner ArCH₂O), 3.40 (d, 2H, *J* = 8.0, cyclic inner OCH₂O), 2.35, 2.13 (m, 8H CH₂(CH₂)₅CH₃), 1.52-1.26 (m, 40H, CH₂(CH₂)₅CH₃), 0.89 (m, 12H, CH₂(CH₂)₅CH₃). ¹³C NMR (CDCl₃, 100 MHz, 25 °C): δ = 165.98, 161.43, 155.88, 150.51, 140.02, 138.28, 134.94, 128.97, 124.73, 123.27, 120.91, 120.70, 102.42, 96.45, 70.79, 52.81, 37.51, 37.31, 32.27, 32.25, 30.27, 30.14, 30.11, 29.88, 29.78, 29.71, 28.45, 28.35, 28.24, 23.09, 23.06, 14.52, 14.51.

Cavitand 8: 2-Bromoresorcinol (145 mg) was used as a bridging reagent and a mixture of hexane:CH₂Cl₂ = 1:1 was

used as eluent for column chromatography. EtOH was used as a recrystallization solvent to give pure cavitand **8** as a white solid (152 mg, 29%). mp >300 °C (dec). MALDI-TOF: m/z 1417 [(M + Na + NaOH)⁺, 30%], 1377 [(M + Na)⁺, 100%]. Elemental analysis calcd for C₇₆H₉₀Br₂O₁₂: C, 67.35; H, 6.69. found: C, 66.79; H, 6.64. ¹H NMR (400 MHz, CDCl₃, 25 °C): δ = 7.17 (s, 4H, ArH), 7.00 (s, 4H, α to Br), 5.92 (d, 2H, J = 8.0, noncyclic outer OCH₂O), 5.45 (d, 4H, J = 12.0, outer ArCH₂O), 5.08 (m, 4H, noncyclic methine + cyclic outer OCH₂O), 4.89 (br-s, 2H, γ to Br), 4.64 (t, 2H, J = 8.0, cyclic methine), 4.35 (m, 6H, noncyclic inner OCH₂O + inner ArCH₂O), 3.52 (d, 2H, J = 8.0, cyclic inner OCH₂O), 2.35, 2.13 (m, 8H CH₂(CH₂)₅CH₃), 1.52-1.26 (m, 40H, CH₂(CH₂)₅CH₃), 0.89 (m, 12H, CH₂(CH₂)₅-CH₃). ¹³C NMR (CDCl₃, 100 MHz, 25 °C): δ = 161.85, 155.78, 150.51, 140.05, 138.26, 124.76, 124.68, 123.69, 120.76, 117.98, 102.39, 96.51, 70.88, 37.48, 37.28, 32.27, 32.23, 31.34, 30.26, 30.12, 29.87, 29.70, 28.44, 28.33, 28.23, 23.09, 23.05, 14.53, 14.50.

Cavitand 9: 2-Chlororesorcinol (112 mg) was used as a bridging reagent and a mixture of hexane:CH₂Cl₂ = 1:1 was used as eluent for column chromatography. CH₃CN was used as a recrystallization solvent to give pure cavitand **9** as a white solid (104 mg, 31%). mp > 300 °C (dec). MALDI-TOF: m/z 1327 [(M + Na + NaOH)⁺, 80%], 1289 [(M + Na)⁺, 100%]. Elemental analysis calcd for C₇₆H₉₀Cl₂O₁₂: C, 72.08; H, 7.16. found: C, 71.50; H, 7.19. ¹H NMR (400 MHz, CDCl₃, 25 °C): δ = 7.17 (s, 4H, ArH), 6.85 (s, 4H, α to Br), 5.93 (d, 2H, J = 8.0, noncyclic outer OCH₂O), 5.45 (d, 4H, J = 12.0, outer ArCH₂O), 5.08 (m, 4H, noncyclic methine + cyclic outer OCH₂O), 4.86 (br-s, 2H, γ to Br), 4.64 (t, 2H, J = 8.0, cyclic methine), 4.35 (m, 6H, noncyclic inner OCH₂O + inner ArCH₂O), 3.52 (d, 2H, J = 8.0, cyclic inner OCH₂O), 2.35, 2.13 (m, 8H CH₂(CH₂)₅CH₃), 1.52-1.26 (m, 40H, CH₂(CH₂)₅CH₃), 0.89 (m, 12H, CH₂(CH₂)₅CH₃). ¹³C NMR (CDCl₃, 100 MHz, 25 °C): δ = 161.78, 155.78, 150.50, 140.03, 138.26, 137.37, 124.68, 120.76, 120.59, 117.47,

102.41, 96.50, 70.83, 37.48, 37.28, 32.27, 32.23, 31.34, 30.26, 30.12, 29.87, 29.70, 28.44, 28.33, 28.23, 23.09, 23.05, 14.52, 14.50.

Cavitand 10: 1,2-Dihydroxybenzotrile (104 mg) was used as a bridging reagent and a mixture of hexane:EtOAc = 8:1 was used as eluent for column chromatography. CH₃CN was used as a recrystallization solvent to give pure cavitand **10** as a white solid (118 mg, 25%). mp > 300 °C (dec). MALDI-TOF : m/z 1309 [(M + Na + NaOH)⁺, 80%], 1269 [(M + Na)⁺, 100%]. ¹H NMR (400 MHz, CDCl₃, 25 °C): δ = 7.28 (s, 4H, α to CN), 7.12 (s, 4H, ArH), 5.94 (d, 2H, J = 8.0, noncyclic outer OCH₂O), 5.53 (d, 4H, J = 12.0, outer ArCH₂O), 5.35 (br-s, 2H, γ to Br), 5.15 (m, 4H, noncyclic methine + cyclic outer OCH₂O), 4.66 (t, 2H, J = 8.0, cyclic methine), 4.0 (d, 4H, J = 12.0, inner ArCH₂O), 4.33 (d, 2H, J = 8.0, noncyclic inner OCH₂O), 3.59 (d, 2H, J = 8.0, cyclic inner OCH₂O), 2.37, 2.15 (m, 8H CH₂(CH₂)₅CH₃), 1.55-1.26 (m, 40H, CH₂(CH₂)₅CH₃), 0.89 (m, 12H, CH₂(CH₂)₅-CH₃). ¹³C NMR (CDCl₃, 100 MHz, 25 °C): δ = 161.31, 155.38, 149.99, 139.50, 137.94, 123.83, 122.83, 120.48, 117.21, 115.88, 101.79, 95.88, 70.34, 37.17, 36.83, 31.85, 31.82, 30.86, 29.81, 29.68, 29.46, 29.28, 27.98, 27.79, 22.68, 22.64, 14.11, 14.09.

Acknowledgment. The financial support from CBMH (Yonsei University), Seoul R&BD program, and Soongsil University is acknowledged.

References

- (a) Ihm, H.; Hwang, S.; Paek, K. *Tetrahedron Lett.* **2004**, *45*, 9119. (b) Paek, K.; Cho, J. H. *Tetrahedron Lett.* **2001**, *42*, 1927. (c) Paek, K.; Ihm, C.; Ihm, H. *Tetrahedron Lett.* **1999**, *40*, 4697.
- Ihm, C.; In, Y.; Park, Y.; Paek, K. *Org. Lett.* **2004**, *6*, 369.
- In, S.; Cho, S. J.; Lee, K. H.; Kang, J. *Org. Lett.* **2005**, *7*, 3993 and references are therein.
- Bryant, J. A.; Blanda, M. T.; Vincenti, M.; Cram, D. J. *J. Am. Chem. Soc.* **1991**, *113*, 2167.