

BULLETIN

OF THE
KOREAN CHEMICAL SOCIETY

VOLUME 17, NUMBER 6
JUNE 20, 1996

BKCS 17(6) 491-572
ISSN 0253-2964

Communications

The Cause of Dispersion Phenomenon in Grunwald-Winstein Plots: Solvolyses of Cyclopentyl Tosylate in Aqueous Binary Mixtures

In Sun Koo, Kiyull Yang, Keumduck Kang, Hyuck Keun Oh*, and Ikchoon Lee**

*Department of Chemical Education,
Geongsang National University, Chinju 660-701, Korea*

**Department of Chemistry,
Chonbuk National University, Chunju 560-756, Korea*

***Department of Chemistry, Inha University,
Incheon 402-751, Korea*

Received February 23, 1996

Dispersion into separate lines in the correlation of the specific rates of solvolysis of a substrate in various binary mixtures was documented¹⁻³ in early treatments using the Grunwald-Winstein equation [eqn. (1)].

$$\log(k/k_0) = mY + c \quad (1)$$

In equation (1), k and k_0 are specific rates of solvolysis of a substrate in a given solvent, of solvent ionizing power Y , and in 80% ethanol respectively, m is the sensitivity to changes in Y values, and c is a residual (constant) term. Equation (1) is commonly written without the intercept (c), which is not required for typical interpretations, but c is often considered "hidden" adjustable parameter in the correlations.⁴

In general, dispersion effects in unimolecular solvolyses^{5,6} make smaller contribution to the overall linear free energy relationship (LFER) than solvent nucleophilicity effects in bimolecular solvolyses. It was suggested² that a second term which is governed by the sensitivity l to solvent nucleophilicity N , should be added to equation (1) for bimolecular solvolyses. The resulting equation (2) is often referred to as the extended Grunwald-Winstein equation.

$$\log(k/k_0) = mY + lN \quad (2)$$

It was suggested that a Y scale based on 2-adamantyl tosylate solvolysis^{7,8,9} (Y_{OTs}) would be the most appropriate, and N_{OTs}

values can be obtained from equation (3).

$$N_{OTs} = \log(k/k_0)_{MeOTs} - 0.3Y_{OTs} \quad (3)$$

The cause of dispersion in Grunwald-Winstein plots using equation (1) is when the leaving group in the solvolysis of a given substrate differs from the leaving group present in the standard substrate used to establish the Y scale being used in the correlation.¹⁰

Another cause of dispersion phenomenon due to not only the resonance stabilization within benzylic carbocations but also appreciable crowding in the vicinity of the reaction center.¹¹

Rate constants for solvolyses of cyclopentyl tosylate are reported for aqueous binary mixtures with acetone, ethanol and methanol (Table 1). In order to eliminate dispersion phenomenon factor to variation of leaving group in Grunwald-Winstein plots, we tried to correlate $\log k$ with Y_{OTs} using k values obtained in this work. The Grunwald-Winstein plots of first-order rate constants for cyclopentyl tosylate with Y_{Cl} (based on 2-adamantyl chloride) show large dispersions (Figure 1). However, the plot of first-order rate constants for cyclopentyl tosylate with Y_{OTs} (based on 2-adamantyl to-

Table 1. Rate Constants (k) Solvolyses of Cyclopentyl Tosylate in Aqueous Alcohol and Acetone Binary Mixtures at 25 °C

v/v %	MeOH	EtOH	Acetone
	$k \times 10^4$		
100	0.115	0.0325	
90	0.242	0.103	
80	0.515	0.254	0.0288
70	1.07	0.496	0.112
60	2.20	0.907	0.321
50	4.56	1.93	0.919
40	9.17	4.35	2.48
30	19.3	12.0	7.69
20	36.8	32.7	21.9
10	69.1	69.3	59.3
H ₂ O	101	101	101




Figure 1. Logarithms of first-order rate constants for solvolyses of cyclopentyl tosylate at 25 °C vs. Y_{Cl} .




Figure 3. Plot of $\log(k/k_0)$ for cyclopentyl tosylate against $0.70Y_{OTS} + 0.91N_{OTS}$ ($r=0.996$).




Figure 2. Logarithms of first-order rate constants for solvolyses of cyclopentyl tosylate at 25 °C vs. Y_{OTS} .

shown to give considerable improvement when it is added as an lN term to original Grunwald-Winstein correlations of the solvolyses of cyclopentyl tosylate. The m value is changed slightly after removal of dispersion, but the associated standard errors are substantially reduced. The correlation coefficient of 0.989 in the absence of the lN term was improved to 0.996 with the lN term.

Therefore, this shows the importance of solvent nucleophilicity compared to solvent ionizing power for the solvolysis of cyclopentyl tosylate. This study has showed that the magnitude of l and m values associated with a change of solvent composition is able to predict the transition state variation to more product like transition state (asymmetric TS, $[N^{\delta+} \cdots C \cdots L^{\delta-}]^{\ddagger}$), where bond breaking is much more progressed and bond formation is slightly more advanced, upon increasing solvent ionizing power or solvent nucleophilicity.

The S_N2 reaction via product like transition state has appeared to cause the dispersions in the present Grunwald-Winstein correlations.

Acknowledgment. This work was supported by Korea Science and Engineering Foundation.

ylate) still shows dispersions (Figure 2).

Cyclopentyl tosylate is neither conjugated systems nor aromatic rings adjacent reaction center in the substrate. Therefore, it can not be explained by phenomenon of the dispersion cause by differential solvation effect on the stabilized reaction center by conjugation with adjacent conjugated system or aromatic rings. Therefore such phenomenon can be explained as dispersion effect caused by other specific solvent effect or change of reaction mechanisms according to the variation of solvent composition. In order to examine the cause of this dispersion phenomenon, we tried to correlate extended Grunwald-Winstein plots of the rate data in Table (1). The extended Grunwald-Winstein plot shows fairly good correlation ($m=0.70$, $l=0.91$, $r=0.996$) for cyclopentyl tosylate solvolysis (Figure 3).

The nucleophilicity parameter (N) has previously been

References

1. Winstein, S.; Grunwald, E. *J. Am. Chem. Soc.* **1948**, *70*, 846.
2. Winstein, S.; Grunwald, E.; Jones, H. W. *J. Am. Chem. Soc.* **1951**, *73*, 2700.
3. Fainberg, A. H.; Winstein, S. *J. Am. Chem. Soc.* **1956**, *78*, 2770.
4. Bentley, T. W.; Bowen, C. T.; Morten, D. H.; Schleyer, P. v. R. *J. Am. Chem. Soc.* **1981**, *103*, 5466.
5. Winstein, S.; Fainberg, A. H.; Grunwald, E. *J. Am. Chem. Soc.* **1957**, *79*, 4146.
6. Fainberg, A. H.; Winstein, S. *J. Am. Chem. Soc.* **1957**, *79*, 1597.
7. Bentley, T. W.; Llewellyn, G. *Prog. Phys. Org. Chem.* **1990**, *17*, 121.

8. Schadt, F. L.; Bentley, T. W.; Schleyer, P. v. R. *J. Am. Chem. Soc.* **1976**, *98*, 7667.
9. Bentley, T. W.; Schleyer, P. v. R. *J. Am. Chem. Soc.* **1976**, *98*, 7656.
10. Fainberg, A. H.; Winstein, S. *J. Am. Chem. Soc.* **1957**, *79*. (a) 1602. (b) 1608.
11. Liu, K.-T.; Sheu, H.-C. *J. Org. Chem.* **1991**, *56*, 3021.


A New Route to a Key Chiral Intermediate of Thienamycin from *trans*-Ethyl Crotonate

Guncheol Kim*[†] and Youngha Ryu

Hanhyo Institutes of Technology,
San 6 Daeyadong, Siheungshi,
Kyunggido 429-010, Korea

Received February 23, 1996

Thienamycin,¹ a β -lactam antibiotic exhibiting broad antibiotic activity, and its derivatives are continuing targets for research and development in the area of antibiotics.² Among various intermediates^{2b,c} to those compounds, acid chloride **1** is unique in that the corresponding amides afford the β -lactam ring, azetidinones, *via* a complete intramolecular S_N2 type cyclization.³ The intermediate **1** has been prepared as a chiral form from *D*-*allo*-threonine⁴ or as a racemic form from *trans*-crotonic acid.³ Considering the high cost of nonracemic threonine, we counted that any route to chiral intermediate **1** from cheap crotonates would be quite compatible.




Scheme 1.

In this brief communication, we describe a concise route to **1** from *trans*-ethyl crotonate employing asymmetric dihydroxylation (ADH).⁵ Although structural transformations after ADH reactions from α,β -unsaturated esters to acetoxy bromo esters have been reported,⁶ to our best knowledge, any route to **1** or to acetoxy bromo acyl chloride derivatives has not been developed. We followed a known procedure⁶ using (DHQD)₂PHAL⁷ to obtain *threo*-2,3-dihydroxy ester in 94% yield. And hydrolysis of the ester using LiOH in MeOH afforded the corresponding acid **2**⁸ quantitatively. The dihydroxy acid **2** was transformed with HBr in acetic acid to provide *erythro*- β -acetoxy- α -bromo acid **3**⁸ in 91% yield. Finally, the desired acyl chloride **1** was readily prepared by treating **3** with SOCl₂ in methylene chloride in a quantitative yield.

[†]Present address: Department of Chemistry, College of Natural Sciences, Chungnam National University, Taejon 305-764, Korea

With no necessity of purification, the four step sequence from the crotonate afforded the intermediate **1** in good yields. This reaction sequence would be applicable in general for the preparation of analogue acid chlorides. And the intermediate **1** was converted to a known azetidinone **4** for comparison using dimethoxybenzylaminomalonate.⁹ The spectral data of **4** (¹H NMR, ¹³C NMR) matched well with those in the literature,⁹ though the optical purity is lower ($[\alpha]_D^{24} + 30.9^\circ$ ($c=2.0$, EtOH), lit.⁹ $[\alpha]_D^{24} + 39.5^\circ$ ($c=2.03$, EtOH)).



Reagents: a. (DHQD)₂PHAL, K₂OsO₂(OH)₄, NMO, *t*-BuOH; LiOH, MeOH
b. HBr, AcOH, 45°C c. SO₂Cl₂, CH₂Cl₂

In conclusion, this concise route employing practical procedures suggests a compatible pathway to the key intermediate, and the optimization for optical purity is left as further work.

References

1. Kahan, J. S.; Kahan, F. M.; Goegelman, R.; Currie, S. A.; Jackson, M.; Stapley, E. O.; Miller, A. K.; Hendlin, D.; Mochales, S.; Hernandez, S.; Woodruff, H. B.; Birnbaum, J. *J. Antibiot.* **1979**, *32*, 1. Albers-Schönberg, G. *et al. J. Am. Chem. Soc.* **1978**, *100*, 6491.
2. For recent reviews see: (a) Palomo, C. Recent Advances in the Synthesis of PS-5 and PS-6 Antibiotics and Related Compounds. *Recent Progress in the Chemical Synthesis of Antibiotics*; Lukacs, G.; Ohno, M. Eds; Spinger-Verlag: **1990**, 565. (b) Nagahara, T.; Kametani, T. *Heterocycles* **1987**, *25*, 729. (c) Kametani, T. *Heterocycles* **1982**, *17*, 463.
3. Shiozaki, M.; Hiraoka, T. *Tetrahedron Lett.* **1980**, *21*, 4473.
4. Shiozaki, M.; Ishida, N.; Hiraoka, T.; Yanagisawa, H. *Tetrahedron Lett.* **1980**, *21*, 4473.
5. For recent reviews see: (a) Kolb, H. C.; VanNieuwenhze, M. S.; Sharpless, K. S. *Chem. Rev.* **1994**, *94*, 2483. (b) Lohray, B. B. *Tetrahedron Asymm.* **1992**, *3*, 1317.
6. Fleming, P. R.; Sharpless, K. B. *J. Org. Chem.* **1991**, *56*, 2869.
7. Sharpless, K. B.; Amberg, W.; Bennani, Y. L.; Crispino, G. A.; Hartung, J.; Jeong, K.-S.; Kwong, H.-L.; Morikawa, K.; Wang, Z.-M.; Xu, D.; Zhang, X.-L. *J. Org. Chem.* **1992**, *57*, 2768.