

Study on FeCl₃ Induced Rearrangement Reaction of Bicyclic Acetal Compound

Si-Min Kim, Yun Hee Maeng, and Jong-Gab Jun*

Department of Chemistry, Hallym University, Chunchon 200-702, Korea


Received September 21, 2001

Keywords : Bicyclic acetal, Ferric chloride, Diacetate, Lewis acid.


Lewis acid induced C-O bond cleavage has been well documented because it is not only important but also variable depending on the structure of starting materials.¹ In our laboratory, we explored Lewis acid systems for the specific transformation of bicyclic acetal **1** to other important structures in unique one-flask reactions.² Recently, we reported an interesting transformation reaction of acetal **1** to monoacetate **2**, diacetate **3** and enone **4** as 37%, 16% and 16% yield, respectively by using AcCl-NaI (Scheme 1).³ We proposed the transformation mechanism (Scheme 2) and selectively obtained the monoacetate **2** in 85% yield as a single product by using MgBr₂-Ac₂O-NaOAc system.⁴ We now wish to report the production of the diacetate **3** selectively.

Ether cleavage and acylation with ferric chloride/acetic anhydride have been reported.⁵ On the other hand, ferric chloride on silica gel has been used in dehydration of alcohols,⁶ cleavage of acetals⁷ and in coupling of phenol ethers.⁸ Also, ferric chloride in acetic acid has been used in acetylation of alcohols, ethers and acetals.⁹


We have extended the use of FeCl₃ as a Lewis acid to the one pot conversion of bicyclic acetal to the corresponding



Scheme 1




Scheme 2




Scheme 3

diketone by rearrangement reactions. Sharma⁹ used FeCl₃ in AcOH for the acetylation of acetal and we adopted this method for the acetylation of bicyclic acetal. The acetal **1** was refluxed overnight with FeCl₃ (0.3 equiv)-AcOH (3 equiv) in CH₂Cl₂ to produce the mixture of 1,5-diketone **5** (54%) and cyclohexenone **6** (7%) as shown in Scheme 3. The mechanism for the 1,5-diketone **5** should be similar to AlCl₃ case.¹⁰ The C (5)-O (6) bond cleavage followed by 1,2-hydride shift *via* an epoxide intermediate produced the diketone **5**, and further reaction proceeded to afford the cyclohexenone **6** as aldol product (Scheme 4). The addition of NaI (1 equiv) in the reaction produced only the diketone in 68% yield. The reaction mechanism of NaI in the reaction was not exactly known, but it prohibited the aldol condensation as shown in TMSCl-NaI system.²¹

The substitution of AcOH to acetic anhydride in the reaction produced the expected diacetate **3** (Scheme 5). Thus, the acetal **1** was refluxed overnight with FeCl₃ (0.3 equiv)-Ac₂O (3 equiv) in CH₂Cl₂ and yielded the diacetate **3** in 84% yield



Scheme 4



Scheme 5

as a single product. Addition of NaI (1 equiv) in the reaction produced only the monoacetate **2** in 60% yield. From the results, we can propose the role of NaI in the reaction as a diminishing the nucleophilicity of acetate anion, but increasing the basicity.

In conclusion, the diacetate **3** was selectively obtained from the acetal **1** by using FeCl₃ with Ac₂O in CH₂Cl₂, while, the monoacetate **2** was also obtained selectively by the addition of NaI in the reaction. But, the best way to make the monoacetate will be the use of MgBr₂-Ac₂O-NaOAc system.

Experimental Section

All chemicals used were purchased from commercial sources and used as received unless otherwise stated. NMR spectra were recorded at Varian Gemini-400 MHz FT-NMR for ¹H and 100 MHz for ¹³C, with the chemical shifts (δ) reported in parts per million (ppm) relative to TMS and the coupling constants (J) quoted in Hz. CDCl₃ was used as a solvent and an internal standard. Infrared spectra were recorded on a Shimadzu IR-435 spectrometer. GC-MS analyses were performed using a HP-5890/JMS-AM 150, JEOL. Flash chromatography was carried out using silica gel Merck 60 (230-400 mesh). Thin-layer chromatography (TLC) was performed on DC-Plastikfolien 60, F254 (Merck, layer thickness 0.2 mm) plastic-backed silica gel plates with visualization by UV light (254 nm) or by treatment with *p*-anisaldehyde.

Procedure by using FeCl₃-AcOH. To a solution of FeCl₃ (50 mg, 0.19 mmol) and AcOH (0.11 mL, 1.92 mmol) in methylene chloride (10 mL) under nitrogen atmosphere was added bicyclic acetal **1** (0.10 g, 0.64 mmol) and refluxed for 15 hr. The reaction was quenched by the addition of 10% aqueous sodium hydroxide solution (10 mL). The organic product was extracted with diethyl ether, washed with brine, dried and concentrated. The residue was purified by silica gel column chromatography (Hexane/EtOAc, 2 : 1) to give the 1,5-diketone **5** (54 mg, 54%, R_f 0.38) and the cyclohexenone **6** (7 mg, 7%, R_f 0.69).^{2f}

Procedure by using FeCl₃-AcOH-NaI. To a solution of FeCl₃ (50 mg, 0.19 mmol), NaI (100 mg, 0.64 mmol) and AcOH (0.11 mL, 1.92 mmol) in methylene chloride (10 mL) under nitrogen atmosphere was added bicyclic acetal **1** (0.10 g, 0.64 mmol) and refluxed for 15 hr. The reaction was quenched by the addition of 10% aqueous sodium hydroxide solution (10 mL). The organic product was extracted with diethyl ether, washed with brine, dried and concentrated. The residue was purified by silica gel column chromatography (Hexane/EtOAc, 2 : 1) to give 7-methyloctane-2,6-dione (**5**) (68 mg, 68%, R_f 0.38).]: ¹H NMR (CDCl₃): δ 2.58 (1H, m, H-7), 2.49 (2H, t, J = 7 Hz, COCH₂), 2.46 (2H, t, J = 7 Hz, COCH₂), 2.13 (3H, s, CH₃CO), 1.83 (2H, quintet, J = 7 Hz, CH₂) 1.09 (6H, d, J = 7 Hz, 2 x CH₃); ¹³C NMR (CDCl₃): δ 214.2 (s), 208.4 (s), 42.6 (t), 40.8 (d), 39.0 (t), 29.8 (q), 18.2 (q, 2 x Me), 17.8 (t); IR (neat): 2963, 1709 (br, C=O), 1465, 1410, 1379, 1174, 1089 cm⁻¹. Ms m/z 156 (M⁺), 141,

123, 113, 85(base), 71, 55; HRMS calcd for C₉H₁₆O₂ (M⁺) 156.1150, found 156.1148.

Procedure by using FeCl₃-Ac₂O. To a solution of FeCl₃ (30 mg, 0.01 mmol) and Ac₂O (0.10 mL, 0.96 mmol) in methylene chloride (10 mL) under nitrogen atmosphere was added bicyclic acetal **1** (0.05 g, 0.32 mmol) and refluxed for 15 hr. The reaction was quenched by the addition of 10% aqueous sodium hydroxide solution (10 mL). The organic product was extracted with diethyl ether, washed with brine, dried and concentrated. The residue was purified by silica gel column chromatography (Hexane/EtOAc, 2 : 1) to give 6,7-diacetoxy-7-methyl-2-octanone (**3**) (42 mg, 84%, R_f 0.32): ¹H NMR (CDCl₃): δ 5.15 (1H, m, H-6), 2.43 (2H, m, COCH₂), 2.12 (3H, s, CH₃CO), 2.08 (3H, s, OCOCH₃), 1.94 (3H, s, OCOCH₃), 1.7-1.5 (4H, m, CH₂CH₂), 1.43 (3H, s, CH₃), 1.40 (3H, s, CH₃); ¹³C NMR (CDCl₃): δ 208.4 (s), 170.5 (s), 170.0 (s), 82.4 (s), 76.2 (d), 42.7 (t), 29.8 (q), 28.1 (t), 22.2 (q), 22.1 (t), 22.0 (q), 20.8 (q), 19.7 (q); IR (neat): 1690 (br, C=O), 1351, 1220, 1130, 1010 cm⁻¹; Ms m/z 157 (M⁺-2CH₂CO-OH), 141, 115, 97, 71, 59, 43 (base); HRMS calcd for C₉H₁₇O₂ (M⁺-2CH₂CO-OH) 157.1229, found 157.1201.

Procedure by using FeCl₃-Ac₂O-NaI. To a solution of FeCl₃ (30 mg, 0.01 mmol), NaI (50 mg, 0.32 mmol) and Ac₂O (0.10 mL, 0.96 mmol) in methylene chloride (10 mL) under nitrogen atmosphere was added bicyclic acetal **1** (0.05 g, 0.32 mmol) and refluxed for 15 hr. The reaction was quenched by the addition of 10% aqueous sodium hydroxide solution (10 mL). The organic product was extracted with diethyl ether, washed with brine, dried and concentrated. The residue was purified by silica gel column chromatography (Hexane/EtOAc, 2 : 1) to give 6-acetoxy-7-methyl-7-octen-2-one (**2**) (30 mg, 60%, R_f 0.32): ¹H NMR (CDCl₃): δ 5.14 (1H, t, J = 5.5 Hz, H-6), 4.93 (1H, br s, C=CH), 4.87 (1H, br s, C=CH), 2.43 (2H, t, J = 7 Hz, COCH₂), 2.12 (3H, s, CH₃CO), 2.04 (3H, s, OCOCH₃), 1.70 (3H, br s, CH₃C=C), 1.7-1.5 (4H, m, CH₂CH₂); ¹³C NMR (CDCl₃): δ 208.3 (s, C=O), 170.2 (s, ester C=O), 142.8 (s, C=CH₂), 112.8 (t, CH₂=C), 76.8 (d, CHOAc), 43.0 (t, COCH₂), 31.9 (t, CH₂CHOAc), 29.8 (q, CH₃CO), 21.1 (q, CH₃COO), 19.4 (t, CH₂CH₂CH₂), 18.0 (q, CH₃C=C); IR (neat): 1695 (br, C=O), 1351, 1225, 1012 cm⁻¹; Ms m/z 156 (M⁺-CH₂CO), 138, 95, 81, 71, 58, 43 (base); HRMS calcd for C₉H₁₆O₂ (M⁺-CH₂CO) 156.1150, found 156.1151.

Acknowledgment. This research was supported by the Hallym University, 2001.

References

- Bhatt, M. V.; Kulkarni, S. U. *Synthesis* **1983**, 249 and references cited therein.
- (a) Bjorklund, M.; Jun, J.-G.; Mundy, B. P. *Tetrahedron Lett.* **1985**, 26, 3895; (b) Jun, J.-G.; Suh, S.; Shin, D. G. *J. Chem. Soc. Perkin Trans. 1* **1989**, 1349; (c) Jun, J.-G.; Shin, H. S. *Synth. Commun.* **1993**, 23, 1871; (d) Jun, J.-G.; Shin, H. S. *Tetrahedron Lett.* **1992**, 33, 4593; (e) Jun, J.-G.; Ha, T. H.; Kim, D.-W. *Tetrahedron Lett.* **1994**, 35, 1235; (f) Jun, J.-G.; Ha, T. H.; Mundy, B. P.; Bartelt, K. E.; Bain, R. S.; Cardellina II, J. H. *J. Chem. Soc. Perkin*

- Trans. 1* **1994**, 2643; (g) Ha, T. H.; Jun, J.-G. *Bull. Korean Chem. Soc.* **1996**, 17, 289; (h) Jun, J.-G.; Ha, T. H. *J. Heterocyclic Chem.* **1997**, 34, 325; (i) Jun, J.-G. *J. Heterocyclic Chem.* **1997**, 34, 633; (j) Kim, A.-R.; Lee, I.-S.; Jun, J.-G. *Synth. Commun.* **2001**, 31, 853.
3. Jun, J.-G.; Lee, D. W.; Mundy, B. P. *Synth. Commun.* **1998**, 28, 2499.
4. Jun, J.-G.; Lee, D. W. *Synth. Commun.* **2000**, 30, 73.
5. (a) Kasturi, T. R.; Abraham, E. M. *J. Chem. Soc., Perkin Trans. 1* **1973**, 2468; (b) Gross, R. S.; Watt, D. S. *Synth. Commun.* **1987**, 17, 1749; (c) Fraga, B. M.; Cabrera, I.; Garcia, V. P.; Guillermo, R.; Perales, A. *Tetrahedron* **1997**, 53, 16177.
6. (a) Ganem, B.; Smale, V. R. *J. Org. Chem.* **1974**, 39, 3728; (b) Keinan, E.; Mazur, Y. *J. Org. Chem.* **1978**, 43, 1020.
7. Tal, D. M.; Keinan, E.; Mazur, Y. *Tetrahedron* **1981**, 37, 4327.
8. Kim, K. S.; Song, Y. H.; Lee, B. H.; Hanhn, C. S. *J. Org. Chem.* **1986**, 51, 404.
9. Sharma, G. V. M.; Mahalingam, A. K.; Nagarajan, M.; Ilangovan, A.; Radhakrishna, P. *Synlett* **1999**, 1200.
10. Jun, J.-G.; Shin, H. S.; Kim, S. H. *J. Chem. Soc. Perkin Trans. 1* **1993**, 1815.
-