

Triterpenoid Saponins from *Elsholtzia bodinieri*

Hao-bin Hu,* Xu-dong Zheng, Huai-sheng Hu, and Yu-feng Jian[†]

College of Chemistry and Chemical Engineering, [†]Department of Biology, Longdong University, Qingyang 745000, P. R. China

*E-mail: huhaobin_88@yahoo.com.cn; hhb-88@126.com

Received April 30, 2007

A new oleanane-type triterpenoid estersaponin, bodinierin C (**1**), along with two known saponins, mazusaponin I (**2**) and ciwujianoside C (**3**), were isolated from the water-soluble part of the root barks of *Elsholtzia bodinieri*. The structure of bodinierin C was characterized by spectroscopic means and chemical hydrolysis as 3β -*O*-caffeoyl-23-hydroxyechinocystic acid 28-*O*- α -*L*-rhamnopyranosyl-(1 \rightarrow 4)- β -*D*-glucopyranosyl-(1 \rightarrow 6)- β -*D*-glucopyranosyl ester. The known compounds were identified by comparing their spectral data with those of authentic samples or data reported in the literature. All compounds were firstly isolated from *Elsholtzia bodinieri* family.

Key Words : *Elsholtzia bodinieri*, Labiatae, 3β -*O*-Caffeoyl-23-hydroxyechinocystic acid 28-*O*- α -*L*-rhamnopyranosyl-(1 \rightarrow 4)- β -*D*-glucopyranosyl-(1 \rightarrow 6)- β -*D*-glucopyranosyl ester, Bodinierin C

Introduction

The genus *Elsholtzia* belongs to the Labiatae family and has approximately 40 species worldwide, which are mainly distributed in East Asia. Thirty-three species, fifteen varieties and five forms of the genus *Elsholtzia* are found in China. Among these, some are used as medicines, some are taken as food and some are source of honey manufacture. The genus *E. wild* generally possesses plentiful volatile oil, which exerts strong inhibition of central nervous system and takes on definite analgesic effect. It shows antibiotic and anti-inflammatory effects as well. As a species of the genus *Elsholtzia*, *E. bodinieri* is an annual herbaceous plant, widely distributed in the mountainous regions of the west and southwest district of China (Chinese name "Dongzisu"),¹ which has been mainly used as a traditional Chinese folk drug for the treatment of eczema, enteritis, diarrhea, bacillary dysentery and cold, and are also known to have anticancer and antibacterial effects.² However, at the best of our knowledge, the active principles of this plant are unknown except for few constituents.³⁻⁷ Therefore, as a continuation of our efforts to pursue the active natural products from *E. bodinieri*, three oleanane-type triterpenoid saponins were isolated by repeated column chromatography and preparative TLC from the *n*-BuOH fraction of the ethanolic extract of *E. bodinieri* gathered in Gansu province of China. Their structures were elucidated as bodinoside C (**1**) (3β -*O*-caffeoyl-23-hydroxyechinocystic acid 28-*O*- α -*L*-rhamnopyranosyl-(1 \rightarrow 4)- β -*D*-glucopyranosyl-(1 \rightarrow 6)- β -*D*-glucopyranosyl ester), mazusaponin I (**2**)⁸ and ciwujianoside C (**3**),⁹ respectively. Among these compounds, **1** was a new one, and all compounds were firstly isolated from the genus *Elsholtzia*. We report herein the isolation and structural elucidation of bodinoside C using chemical and spectral evidence.

Results and Discussion

Compound **1** was obtained as a pale yellow amorphous

powder from MeOH and responded positively to the Molisch and Liebermann-Burchard tests for triterpene glycoside. The molecular formula was established as C₅₇H₈₄O₂₂ from the HRFAB-MS (positive) ion peak at *m/z* 1143.5338 [M+Na]⁺ (C₅₇H₈₄O₂₂Na, calcd. for 1143.5352), corresponding to 16 degrees of unsaturation. The positive-ion FAB-MS also displayed 975 [M+H-146]⁺, 813 [M+H-146-162]⁺ and 651 [M+H-146-162-162]⁺, which showed the presence of one terminal 6-deoxysugar and two inter hexose units in a linear linkage. It exhibited UV maxima at 220, 237, 298 and 325 nm, suggesting the presence of strong conjugation in the molecule, a bathochromic shift of 47 nm by adding NaOH indicated the presence of free phenolic hydroxy, and a distinct bathochromic shift with AlCl₃ but again returning to normal value after adding HCl, indicated the presence of catechol group. Its IR spectrum (KBr) of **1** indicated the presence of hydroxyl groups (3450-3100 cm⁻¹), α,β -unsaturated ester moiety (1694 and 1270 cm⁻¹), trisubstituted double bond (1634 cm⁻¹), aromatic rings (1595 and 1514 cm⁻¹) and glucoside functionalities (1086, 1070, 1036 cm⁻¹). The broad band decoupled ¹³C-NMR spectrum showed 57 carbon signals, of which 30 were assigned to the pentacyclic triterpene moiety, 9 to the caffeic acid moiety and 18 to the saccharide portion. The DEPT spectrum displayed the presence of twelve quaternary, twenty-six methine, twelve secondary methene and seven primary methyl carbons in the molecule. The analysis of NMR spectra by the aid of DEPT technique demonstrated the presence of a caffeoyl group [including two *trans*-conjugated olefinic protons at δ_{H} 6.32 (1H, d, *J* = 16.0 Hz) and 7.55 (1H, d, *J* = 16.0 Hz) were connected directly to carbons at δ_{C} 116.0 and 144.8, an ester carbonyl (δ_{C} 167.8), three aromatic methine protons of an typical ABX type at δ_{H} 7.08 (1H, d, *J* = 1.5 Hz), 6.88 (1H, d, *J* = 8.0 Hz) and 7.10 (1H, dd, *J* = 8.0, 1.5 Hz) were linked to three aromatic tertiary carbons at δ_{C} 115.4, 114.3 and 122.2, another three aromatic quaternary carbons at δ_{C} 127.1, 144.1, 146.6], a triaccharide chain [three anomeric proton signals at δ_{H} 6.18 (1H, d, *J* =

8.0 Hz), 4.95 (1H, d, $J = 8.0$ Hz), 5.82 (1H, d, $J = 1.5$ Hz) and one doublet methyl at δ_{H} 1.66 (1H, d, $J = 6.0$ Hz), corresponding to three anomeric carbon and methyl signals at δ_{C} 95.7, 104.7, 102.8 and 18.8]. On alkaline hydrolysis, **1** yielded caffeic acid, *D*-glucose and *L*-rhamnose (in the molar ratio of 1:2:1), respectively, which was compared with authentic sample by *co*-TLC and *co*-PC. Moreover, The ^1H and ^{13}C -NMR of **1** exhibited six angular methyl groups [δ_{H} 1.32, 1.18, 1.10, 1.65, 0.98 and 1.08, corresponding to δ_{C} 15.6, 16.7, 17.7, 27.2, 33.3 and 24.8] and one double bond [δ_{C} 123.1 and 143.9, a broad triplet at δ_{H} 5.41(1H, br t)], characteristic of a typical Δ^{12} -oleanene skeleton.^{10,11} The signals at δ_{H} 4.10 (1H, d, $J = 11.0$ Hz)/3.68 (1H, d, $J = 11.0$ Hz) and δ_{C} 66.1 were assigned to H₂-23 and C-23, respectively, suggesting that a hydroxyl group was linked at C-23 on the basis of the chemical shifts of C-23 and C-24.¹² This conclusion was also drawn from the correlations of H-23 with C-3/C-4/C-24 and H-24 with C-3/C-4/C-5/C-23 in the HMBC spectrum as well as the observed NOE interactions between H-23 and H-3/H-5, H-24 and H-25 in the ROESY spectrum. In addition, the signals at δ_{H} 5.06 (1H, m) and δ_{C} 74.2 were assigned to H-16 and C-16, which suggested that another hydroxyl group was linked at C-16¹³ with α -configuration on account of the chemical shifts of C-16 and H-16, along with the correlations of H-16 with C-14/C-17/C-18/C-22/C-28 and H-16 with H-18 in the HMBC and ROESY spectrum. By comparison of the spectroscopic data of the aglycone of **1** with echinocystic acid¹⁴ and 3 β , 23, 28-trihydroxy-12-oleanene 3 β -caffeate isolated previously from *Hibiscus syriacus*,¹⁵ the aglycone of **1** was suggested to be 23-hydroxyechinocystic acid with a caffeoyl linkage at C-3, which was confirmed by the correlations of H-3 with C-2/C-4/C-23/C-24/C-9 in the HMBC spectrum, and also consistent with the observed downfield shifts of C-3 (δ_{C} 80.9) and H-3 (δ_{H} 5.02) with respect to the corresponding signal in hederagenin 28-*O*- α -*L*-rhamnopyranosyl-(1 \rightarrow 4)- β -*D*-glucopyranosyl-(1 \rightarrow 6)- β -*D*-glucopyranoside.¹⁶ The β -configuration of 3-*O*-caffeoyl group was determined from the chemical shifts and coupling constants of H-3 (5.02, dd, $J =$

5.5, 11.0 Hz),^{17,18} and further evidenced by the observed NOE interactions between H-3 and H-1/H-5/H-23 in the ROESY spectrum. These spectral features and physico-chemical properties suggested **1** to be 3 β -*O*-caffeoyl-23-hydroxyechinocystic acid with three sugar moieties.

Comparison of NMR data of the sugar moieties with literature values¹⁶ revealed that the glucoses and rhamnose were present in pyranoside form. The relative stereochemistry of each monosaccharide was determined as β -glucopyranose and α -rhamnopyranose based on the characteristic $^3J_{\text{H-1,H-2}}$ coupling constants (8.0 and 1.5 Hz)²⁰ and ^{13}C -NMR data. The HMBC correlations (Fig. 2) between H-1'''' of the terminal rhamnopyranosyl unit and C-4''' of the centre glucopyranosyl unit, H-4''' of the centre glucopyranosyl unit and C-1'''' of the terminal rhamnopyranosyl unit, H-1''' of the centre glucopyranosyl unit and C-6'' of the inner glucopyranosyl unit, together with H-6'' of the inner glucopyranosyl unit and C-1''' of the centre glucopyranosyl unit, suggested the linkage of α -*L*-rhamnopyranosyl-(1 \rightarrow 4)- β -*D*-glucopyranosyl-(1 \rightarrow 6)- β -*D*-glucopyranosyl. Further supporting information came from the observed NOE interactions of H-6''/H-1'''' and H-4'''/H-1'''' in ROESY experiment (Fig. 3), together with the FAB-MS ion peaks at m/z 975 [$\text{M}+\text{H}-146$]⁺ (loss of a terminal rhamnose unit) and 813 [$\text{M}+\text{H}-146-162$]⁺ and 651 [$\text{M}+\text{H}-146-162-162$]⁺ (loss of two inner glucose units successively). Easy hydrolysis of the compound **1** with base confirmed that sugar was attached by an ester linkage. The exact position of the trisaccharide chain at C-28 of the aglycone was established from the HMBC correlations between the H-1 (δ_{H} 6.18) of ester-linked β -glucopyranosyl unit and the C-28 (δ_{C} 176.3) of aglycone, this was also supported by the typical upfield (*ca* 4.4) *O*-glycosylation shift of C-28 compared to the shift of a free carboxylic acid²¹ and the anomeric carbon signal (δ_{C} 95.7) of β -glucopyranosyl unit, indicating the presence of sugar ester. From the foregoing evidences, the structure of compound **1** was established as 3 β -*O*-caffeoyl-23-hydroxyechinocystic acid 28-*O*- α -*L*-rhamnopyranosyl-(1 \rightarrow 4)- β -*D*-glucopyranosyl-(1 \rightarrow 6)- β -*D*-glucopyranosyl ester, named




Figure 1. The structures of compound 1-3.




Figure 2. The key HMBC correlations of compound 1.




Figure 3. The key ROESY correlations of compound **1**.

bodinierin C. To our knowledge, **1** has not been reported previously from any plant source.

The known compounds **2** and **3** were identified by analysis and comparison of their spectral data obtained with literature values.

Experimental Section

General procedures. Melting points were measured on a Chinese X-4 melting point apparatus (uncorrected); Optical

rotation was determined on a Perkin-Elmer 341 automatic polarimeter; IR spectra (KBr disks) were obtained on Alpha-Centari FT-IR spectrometer; UV spectra were run on a Shimadzu UV-300 spectrophotometer (double beam); NMR spectra were scanned on Bruker AM-500 spectrometer (chemical shifts in δ downfield from TMS internal standard) operating 500 and 125 MHz for ^1H and ^{13}C respectively, and FAB-MS spectra on VG Autospect 3000 spectrometer; TLC was performed on silica gel GF₂₅₄ plates (0.50 mm thickness); Spots were visualized under UV light or by exposure to I_2 vapours and/or by spraying (analyt. TLC) with 10% H_2SO_4 in EtOH, followed by heating for a few minute; Separation and purification were performed by column chromatography on silica gel (100-200, 200-300 mesh). All solvents were distilled before use.

Plant materials. The root barks of *E. bodinier* were collected in August 2004 from Ziwuling mountainous area of Gansu Province in China, and identified by Fu-shun Liu (Department of Biology, Longdong University, China). A voucher specimen (No. 107083) of the plant is previously deposited at the Herbarium of the Botany Department, Longdong University, Qingyang, 745000, China.

Extraction and isolation. The air-dried root barks of *E. bodinier* (2.5 kg) were extracted in soxhlet successively with 75% EtOH for 3 days. The extracts were evaporated to dryness. The residue (155 g) was suspended in warm water

Table 1. ^1H and ^{13}C -NMR data of compound **1** (500/125 MHz, pyridine- d_5)^a

No.	δ_{H}	δ_{C} (DEPT)	No.	δ_{H}	δ_{C} (DEPT)
1	1.67(1H, m)/1.48(1H, m)	38.3(CH ₂)	30	1.08(3H, s)	24.8(CH ₃)
2	2.19/2.08(brd, 12.3)	24.2(CH ₂)	1'	–	127.1(C)
3	5.02(1H, dd, 5.5, 11.0)	80.9(CH)	2'	7.08 (1H, d, 1.5)	115.4(CH)
4	–	43.7(C)	3'	–	144.1(C)
5	1.68(1H, m)	48.1(CH)	4'	–	146.6(C)
6	1.58(1H, m)/1.49(1H, m)	18.4(CH ₂)	5'	6.88(1H, d, 8.0)	114.3(CH)
7	2.10(1H, m)/1.87(1H, m)	33.5(CH ₂)	6'	7.10(1H, dd, 8.0, 1.5)	122.2(CH)
8	–	40.1(C)	7'	7.55(1H, d, 16.0)	144.8(CH)
9	1.68(1H, m)	47.4(CH)	8'	6.32(1H, d, 16.0)	116.0(CH)
10	–	37.1(C)	9'	–	167.8(C)
11	2.02(1H, m)/1.98(1H, m)	24.2(CH ₂)	28-O-Glycosyl moieties		
12	5.41(1H, br t)	123.1(CH)	Glc-1''	6.18(1H, d, 8.0)	95.7(CH)
13	–	143.9(C)	2''	4.09(1H, t, 8.0)	74.1(CH)
14	–	42.3(C)	3''	4.17(1H, m)	78.6(CH)
15	1.74(1H, dd, 3.5, 14.9)/1.51(1H, dd, 2.6, 14.9)	36.4(CH ₂)	4''	4.23(1H, m)	71.5(CH)
16	5.06(1H, m)	74.2(CH)	5''	4.08(1H, m)	78.0(CH)
17	–	49.4(C)	6''	4.66(1H, m)/4.30(1H, m)	69.2(CH ₂)
18	3.41(1H, dd, 4.2, 13.5)	41.8(CH)	Glc-1'''	4.95(1H, d, 8.0)	104.7(CH)
19	1.27(1H, m)/2.33(1H, t, 13.7)	47.4(CH ₂)	2'''	3.96(1H, t, 8.0)	75.5(CH)
20	–	30.8(C)	3'''	4.12(1H, m)	76.5(CH)
21	1.98(1H, m)/1.21(1H, m)	36.1(CH ₂)	4'''	4.36(1H, t, 8.0)	78.5(CH)
22	1.95(1H, m)/1.82(1H, dd, 4.7, 13.4)	32.2(CH ₂)	5'''	3.66(1H, m)	77.1(CH)
23	4.10(1H, d, 11.0)/3.68(1H, d, 11.0)	66.1(CH ₂)	6'''	4.20(1H, m)/4.10(1H, m)	61.3(CH ₂)
24	1.32(3H, s)	15.6(CH ₃)	Rha-1''''	5.82(1H, d, 1.5)	102.8(CH)
25	1.18(3H, s)	16.7(CH ₃)	2''''	4.64(1H, m)	72.5(CH)
26	1.10(3H, s)	17.7(CH ₃)	3''''	4.53(1H, dd, 9.0, 2.0)	72.6(CH)
27	1.65(3H, s)	27.2(CH ₃)	4''''	4.31(1H, m)	74.0(CH)
28	–	176.3(C)	5''''	4.91(1H, m)	70.5(CH)
29	0.98(3H, s)	33.3(CH ₃)	6''''	1.66(1H, d, 6.0)	18.8(CH ₃)

^aSignals were assigned by ^1H -NMR, ^{13}C -NMR, ^1H - ^1H COSY, ROESY, HMBC, 90° and 135° DEPT.

and extracted with water-saturated *n*-BuOH. The organic layer was concentrated to obtain a residue (50 g) which was redissolved in MeOH (300 mL). Addition of Et₂O gave a flocculent precipitate which was washed with Et₂O (three times), yielded a crude saponin mixture (30 g) which was subjected to silica gel column chromatography (100-200 mesh) with EtOAc-MeOH-H₂O (10:5:1 → 100% MeOH, v/v/v) in increasing polarity and combined by monitoring with TLC to give ten fractions (A-J). Fraction G (2 g) was repeatedly subjected to silica gel column chromatography (200-300 mesh), then with MCI-gel CHP-20 to afford **2** (12 mg). Fraction I (560 mg) was separated successively over Sephadex LH-20 (MeOH-H₂O 8:2) and silica gel column (CHCl₃-MeOH-H₂O 6:4:1 → 10:5:1) to obtain **3** (11 mg). Fraction J (1.2 g) was further purified by preparative TLC and developed with Me₂CO-MeOH (1:8, v/v) as development to provide compound **1** (15 mg).

Bodinerin C (1): Pale yellow amorphous powder from MeOH, C₅₇H₈₄O₂₂, mp. 208-210 °C, [α]_D²⁰ -19.2° (c 0.41, MeOH); UV $\lambda_{\max}^{\text{MeOH}}$: 220, 237, 298 and 325 nm; (+NaOH): 372 nm; (+AlCl₃): 382 nm; IR ν_{\max}^{KBr} : 3450-3100, 2921, 1723, 1694, 1634, 1595, 1514, 1464, 1270, 1086, 1070, 1036, 940 cm⁻¹; HRFAB-MS (positive-ion mode): *m/z* 1143.5338 [M+Na]⁺; FAB-MS: *m/z* 975 [M+H-146]⁺, 813 [M+H-146-162]⁺ and 651 [M+H-146-162-162]⁺; ¹H and ¹³C-NMR see Table 1.

Mazosaponin I (2): White powder from MeOH, C₂₆H₄₄O₁₁, mp. 189-191 °C, [α]_D²⁰ -32.5° (c 0.25, MeOH); UV $\lambda_{\max}^{\text{MeOH}}$: 210 nm; ¹³C-NMR (125 MHz, pyridine-*d*₅): δ_{C} 38.8 (C-1), 26.7 (C-2), 88.6 (C-3), 39.6 (C-4), 55.1 (C-5), 18.6 (C-6), 33.2 (C-7), 40.2 (C-8), 48.3 (C-9), 37.2 (C-10), 24.2 (C-11), 123.1 (C-12), 144.3 (C-13), 42.2 (C-14), 28.9 (C-15), 27.9 (C-16), 46.7 (C-17), 44.5 (C-18), 81.2 (C-19), 35.5 (C-20), 29.1 (C-21), 33.1 (C-22), 28.1 (C-23), 16.8 (C-24), 15.6 (C-25), 17.7 (C-26), 24.9 (C-27), 177.2 (C-28), 28.6 (C-29), 24.7 (C-30), α -Ara: 107.6 (C-1), 72.7 (C-2), 74.4 (C-3), 69.6 (C-4), 66.8 (C-5), β -Glc: 95.8 (C-1), 73.8 (C-2), 78.5 (C-3), 70.7 (C-4), 78.1 (C-5), 69.3 (C-6), β -Glc: 105.1 (C-1), 75.2 (C-2), 78.3 (C-3), 71.4 (C-4), 78.4 (C-5), 62.7 (C-6); FAB-MS (negative-ion mode): *m/z* 927 [M-H]⁻, 603 [M-H-2Glc]⁻, 471 [M-H-2Glc-Ara]⁻. ¹H-NMR and IR data were consistent with those reported in the literature.⁸

Ciwujianoside C (3): White powder from MeOH, C₅₃H₈₆O₂₁, mp. 211-213 °C (decompose), UV $\lambda_{\max}^{\text{MeOH}}$ 210 nm; ¹³C-NMR (125 MHz, pyridine-*d*₅): δ_{C} 38.7 (C-1), 26.5 (C-2), 87.9 (C-3), 39.5 (C-4), 55.8 (C-5), 18.1 (C-6), 33.6 (C-7), 39.6 (C-8), 47.8 (C-9), 36.5 (C-10), 23.1 (C-11), 122.5 (C-12), 143.8 (C-13), 41.8 (C-14), 26.6 (C-15), 23.6 (C-16), 46.9 (C-17), 41.6 (C-18), 45.9 (C-19), 30.7 (C-20), 33.5 (C-21), 32.3 (C-22), 28.2 (C-23), 16.9 (C-24), 15.5 (C-25), 17.6 (C-26), 25.6 (C-27), 176.2 (C-28), 32.8 (C-29), 23.1 (C-30), α -Ara: 106.9 (C-1), 72.5 (C-2), 74.2 (C-3), 69.1 (C-4), 66.3 (C-5), β -Glc: 95.9 (C-1), 73.6 (C-2), 78.1 (C-3), 70.4 (C-4), 76.5 (C-5), 69.1 (C-6), β -Glc: 104.9 (C-1), 75.1 (C-2), 76.3 (C-3), 78.4 (C-4), 77.8 (C-5), 61.1 (C-6), α -Rha: 102.2 (C-1), 72.3 (C-2), 72.7 (C-3), 73.6 (C-4), 70.1 (C-5),

18.2 (C-6); FAB-MS (negative-ion mode): *m/z* 1057 [M-H]⁻, 587 [M-H-2Glc-Rha]⁻, 455 [M-H-2Glc-Rha-Ara]⁻. ¹H-NMR and IR data were identical with those of literature.⁹

Alkaline hydrolysis of compound 1: Compound **1** (5 mg) was treated with an aqueous solution of Ba(OH)₂ (0.30 mol/L) for 4 hr at 100 °C. The reaction mixture was cooled, then adjusted to PH = 7 with 10% H₂SO₄. The BaSO₄ was removed by filtration and the solution was extracted with Et₂O (twice), then the ether layer was concentrated under reduced pressure to give a light-yellow solution of caffeic acid, which was identified by *co*-TLC with authentic samples using bromphenol blue as chromogenic reagent. The aqueous layer was partitioned between *n*-BuOH/H₂O, and the aqueous layer was concentrated *in vacuo*. The sugar were identified by *co*-PC with authentic samples using [*n*-BuOH:HOAc:H₂O (4:1:5, v/v/v)] as developing solvent and 0.9% aniline-oxalate solution as color developing reagent (*D*-glucose *R_f*: 0.19, brown; *L*-rhamnose *R_f*: 0.37, brown).

Acknowledgements. To Fu-shun Liu for identifying the plant samples, and to the Education Department of Gansu Province of China (Grant No. 0610-05) and the Longdong University (Grant No. XYZK0503) for financial support.

References

1. Delectis Florae Reipublicae Popularis Sinicae Agendae Academiae Sinicae Edita. *Flora Reipublicae Popularis Sinicae*; Science Press: Beijing, 1990; Vol. 60, p 336.
2. Jiangsu New Medical College. *Dictionary of Traditional Chinese Drug*; Shanghai Science and Technology Press: Shanghai, 1985; Vol. 1, p 489.
3. Zhu, W. M.; He, H. P.; Wang, S. Z.; Zuo, G. Y.; Hao, X. J. *Chin. Chem. Lett.* **2000**, *13*, 253.
4. Li, R. T.; Li, J. T.; Wang, J. K. *Helv. Chim. Acta* **2005**, *88*, 252.
5. Zhao, D. B.; Yang, Y. X.; Zhang, W.; Liu, X. H.; Zhai, C. P.; Wang, H. Q. *J. Chin. Med. Mat.* **2005**, *28*, 94.
6. Hu, H. B.; Liu, J. X.; Zheng, X. D. *Chin. Tradit. Herb Drugs* **2006**, *37*, 18.
7. Hu, H. B.; Jian, Y. F.; Zheng, X. D.; Cao, H. *Bull. Korean Chem. Soc.* **2007**, *28*, 351.
8. Yaguchi, E.; Miyase, T.; Ueno, A. *Phytochemistry* **1995**, *39*, 185.
9. Shao, C. J.; Kesai, R.; Xu, J. D.; Tanaka, O. *Chem. Pharm. Bull.* **1988**, *36*, 601.
10. Ikuta, A.; Itokawa, H. *Phytochemistry* **1988**, *27*, 2813.
11. Conrad, J.; Vogler, B.; Klaiber, I.; Roos, G.; Walter, U.; Kraus, W. *Phytochemistry* **1998**, *48*, 647.
12. Mahato, S. B.; Kundu, A. P. *Phytochemistry* **1994**, *37*, 1517.
13. Zhang, Y. J.; Yang, C. R. *Phytochemistry* **1994**, *36*, 997.
14. Niranjana, P. S.; Shashi, B. M. *Phytochemistry* **1994**, *37*, 1425.
15. Yun, B. S.; Ryoo, I. J.; Lee, I. K.; Park, K. H.; Choung, D. H.; Han, K. H.; Yoo, I. D. *J. Nat. Prod.* **1999**, *62*, 764.
16. Li, X. C.; Wang, D. Z.; Wu, S. G. *Phytochemistry* **1990**, *29*, 595.
17. Teng, R. W.; Zhong, H. M.; Chen, C. X.; Wang, D. Z. *Chin. J. Magn. Reson.* **1999**, *16*, 389.
18. Teng, R. W.; Wang, D. Z.; Chen, C. X. *Chin. Chem. Lett.* **2000**, *11*, 337.
19. He, Z. D.; Qiao, C. F.; Han, Q. B.; Wang, Y.; Ye, W. C.; Xu, H. X. *Tetrahedron* **2005**, *61*, 2211.
20. Agrawal, P. K. *Phytochemistry* **1992**, *31*, 3307.
21. Haddad, M.; Miyamoto, T.; Laurens, V.; Lacaille-Dubois, M. A. *J. Nat. Prod.* **2003**, *66*, 372.