

A Single Amino Acid Substitution Alters Substrate Sequence Specificity of the Yeast Protein Tyrosine Phosphatase YPTP1

Dongki Yang, Misook Choi, Junseo Park, Hwang-Keun Jun, Hee-Young Choi, Sun-Myoung Lee, Jae-Seung Kang,[†] and Hyeongjin Cho*

Division of Chemistry and Physics, College of Natural Sciences

[†]Department of Microbiology, College of Medicine, Inha University, Incheon 402-751, Korea

Received August 17, 1999

Protein tyrosine phosphatases (PTPases) constitute a family of enzymes that catalyze the hydrolysis of a phosphate moiety from a phosphotyrosyl residue of a protein.¹⁻⁵ Because of the biological importance of the reversible tyrosine phosphorylation, much research has been focused on the PTPases since its first identification in 1988.⁶ More than 70 PTPases have been identified in various organisms including human to yeasts.⁷ Much progress has been achieved in the study of the mechanism of catalysis.⁴ However, *in vivo* substrates and biological roles of most of the PTPases still remains to be elucidated.

PTPases recognize short phosphotyrosyl(pY)-peptides as substrates and the sequence specificities of several PTPases toward pY-peptides have been investigated.⁷⁻¹³ We previously reported that the lowest K_M substrate for the yeast PTPase, YPTP1, was DADEpYDA which is characterized by the multiple acidic residues at the N-terminal side of pY.⁸ Similar trends were also found in many other PTPases including PTP1B,⁹ HPTP β ,¹⁰ T-cell PTPase,¹¹ PTP-5 from bovine brain and PTP-1 from rat brain,¹² *Yersinia* PTPase and rat PTP1.¹³

In the case of PTP1B, the sequence specificity was explained on the molecular basis by Jia *et al.*¹⁴ X-ray crystallographic analysis of inactive PTP1B complexed with a phosphotyrosyl peptide, DADEpYL-NH₂, revealed that Arg-47 is responsible for defining the sequence specificity of PTP1B; Arg-47 forms salt bridges with the carboxyl side chains of Asp and Glu at -1 and -2 position of N-terminal side of pY (pY-1 and pY-2 positions) of the peptide. It is noteworthy that rat brain PTP-1, *Yersinia* PTPase, and T-cell PTPase, which share a preference for pY-peptides with acidic amino acid residues N-terminal to the pY, contain Arg at the equivalent position of Arg-47 of PTP1B.^{15,16}

YPTP1, on the other hand, has Val-59,¹⁷ not Arg, at the equivalent position still exhibiting sequence specificity similar to the PTPases mentioned above. HPTP β does not have Arg at the position, either. It is Gln in HPTP β .¹⁵ Diverse amino acid residues appear in PTPases at the position equivalent to Arg-47 of PTP1B; they are Ser, Lys, Val, Ile, Gln, Thr and Ala.⁵

To investigate if Val-59 is an important determinant for the peptide substrate recognition by YPTP1, we substituted Val-59 with Asp or His by site-directed mutageneses and per-

formed kinetic experiments. These mutations that change the hydrophobic side chain of Val to amino acid side chains with a negative or a positive charge might alter the preferences of YPTP1 for the peptide substrate sequences. Previously we subcloned the *YPTP1* gene in the *E. coli* expression vector pT7-7 (pT7-*YPTP1*) and used it for the overexpression of the wild-type YPTP1.⁸ To generate the mutants, YPTP1 (V59D) and YPTP1 (V59H), we modified pT7-*YPTP1* plasmid by the "overlap extension method"¹⁸ based on polymerase chain reaction (PCR) technique. The wild-type and mutant PTPases were purified to > 90% purity by the combination of affinity chromatography and conventional chromatography techniques as described previously for wild-type YPTP1.^{8,19,20} A few milligrams of wild-type or mutant enzymes are generally obtained from 1 liter of *E. coli* culture. Specific activities of purified mutant enzymes toward p-nitrophenyl phosphate are 30-40 $\mu\text{mol} \cdot \text{min}^{-1} \cdot \text{mg}^{-1}$ which are essentially indistinguishable with that of wild-type YPTP1.^{10,19}

Kinetic experiments were performed with wild-type and the two mutant phosphatases (Table 1).⁸ Chemically synthesized 5 mer pY-peptides with sequences AXApYA (X is a variable amino acid) were used as substrates.^{10,21} Eventhough the negatively charged residues at pY-2 and pY-1 positions are both necessary for high affinity recognition by YPTP1, to avoid complication, the -1 position is fixed to Ala and only the -2 position of pY was changed sequentially.

The first substrate examined in this study was ADApYA. Toward this substrate wild-type YPTP1 exhibited a K_M value of 34 μM , 8.5-fold higher than that of the lowest K_M substrate DADEpYDA ($K_M = 4 \mu\text{M}$).⁸ Because of the importance of the negative charge at pY-2 position of the peptide, it is anticipated that the introduction of a positively charged amino acid to that position would reduce the affinity for wild-type YPTP1 and this was truly the case; $K_M = 159 \mu\text{M}$

Table 1. K_M values with Wild-Type and Mutant Recombinant Yeast YPTP1 (unit: μM)

Line	Peptides	YPTP1		
		wild-type	V59D mutant	V59H mutant
1	H ₂ N-ADApYA-OH	34	220	280
2	H ₂ N-ARApYA-OH	159	92	290
3	Ac-ADApYA-NH ₂	43	162	198
4	Ac-AHApYA-NH ₂	156	127	> 400

*To whom correspondence should be addressed. E-mail: hcho@inha.ac.kr, Phone: +82-32-860-7683, Fax: +82-32-867-5604

for ARApYA (Line 2).

To determine if Val-59 is an important determinant for the recognition of the pY-2 residue of peptide substrates, we substituted Val-59 of YPTP1 with the negatively charged Asp residue. YPTP1(V59D) mutant showed 7-fold reduced affinity toward ADAPYA (Line 1) and 2-fold increased affinity toward ARApYA (Line 2). Collectively, ADAPYA was a better substrate for wild-type YPTP1 and ARApYA was better for the mutant enzyme.

When the N-termini of the peptides were acetylated and the C-termini were amidized to eliminate the effect of the charge at both ends, the peptide substrates showed similar trends. Ac-ADAPYA-NH₂ gave K_M values not much different from those obtained with the unmodified version of the peptide with the same sequence for the wild-type and mutant enzymes (Line 1 and 3). Results obtained with Ac-AHApYA-NH₂ (Line 4) was also comparable with those obtained with ARApYA (Line 2). These results implicate that Val-59 of YPTP1 is involved in the recognition of the pY-2 residue of the pY-peptide substrate.

Substitution of Val-59 with His in YPTP1, in agreement with the results described above, was not favorable for the interaction of the enzyme with Ac-AHApYA-NH₂ (Line 4). YPTP1(V59H), however, did not differentiate Asp and Arg at pY-2 position of the peptide (Line 1 and 2). This result, together with the fact that V59H mutant exhibits low affinity toward any substrates tested, indicates that electronic interaction is not a prevailing factor in this case.

It may be possible to speculate that the peptide substrates bind to YPTP1 in a peptide sequence-dependent manner; *i.e.* positively charged Arg residue of H₂N-ARApYA-OH interacts favorably with the negatively charged Asp-59 residue of YPTP1(V59D) thereby exhibiting a low K_M; Asp residue of H₂N-ADAPYA-OH, however, interacts with a residue other than Val-59 of YPTP1 for high affinity binding to wild-type YPTP1. We constructed a plasmid encoding V59R mutant for extended study but unfortunately this mutant protein is not expressed in an active form in *E. coli*. Unambiguous explanation of the structure-specificity relationship of YPTP1 would be possible by extensive studies of X-ray crystal structures of YPTP1 and its mutants complexed with pY-peptides, and these studies are in progress.

In summary, we modified the sequence specificity of YPTP1 for the pY-peptide substrates by the single amino acid substitution. Especially V59D mutation reversed the specificity of the wild-type enzyme. This study also demonstrates that Val59 plays an important role in the recognition of the pY-peptide substrates.

Acknowledgment. The authors wish to acknowledge the financial support of the Korea Research Foundation made in the program year of 1997.

References

- Walton, K. M.; Dixon, J. E. *Annu. Rev. Biochem.* **1993**, *62*, 101.
- Zhang, Z.-Y.; Dixon, J. E. *Adv. Enzymol.* **1994**, *68*, 1.

- Tonks, N. K.; Neel, B. G. *Cell* **1996**, *87*, 365.
- Zhang, Z. Y. *Curr. Top. Cell. Regul.* **1997**, *35*, 21.
- Huijsduijnen, R. H. *Gene* **1998**, *225*, 1.
- Tonks, N. K.; Diltz, C. D.; Fischer, E. H. *J. Biol. Chem.* **1988**, *263*, 6731.
- Wu, L.; Buist, A.; Hertog, J.; Zhang, Z.-Y. *J. Biol. Chem.* **1997**, *272*, 6994.
- Kwon, M.; Oh, M.; Han, J.; Cho, H. *J. Biochem. Mol. Biol.* **1996**, *29*, 386.
- Zhang, Z.-Y.; Thieme-Seffler, A. M.; Maclean, D.; McNamara, D. J.; Dobrustin, E. M.; Sawyer, T. K.; Dixon, J. E. *Proc. Natl. Acad. Sci. U.S.A.* **1993**, *90*, 4446.
- Cho, H.; Krishnaraj, R.; Itoh, M.; Kitas, E.; Bannwarth, W.; Saito, H.; Walsh, C. T. *Protein Sci.* **1993**, *2*, 977.
- Ruzzene, M.; Donella-Deana, A.; Marin, O.; Perich, J. W.; Ruzza, P.; Borin, G.; Calderan, A.; Pinna, L. A. *Eur. J. Biochem.* **1993**, *211*, 289.
- Hippen, K. L.; Jakes, S.; Richards, J.; Jena, B. P.; Beck, B. L.; Tabatabai, L. B.; Ingebristen, T. S. *Biochemistry* **1993**, *32*, 12405.
- Zhang, Z.-Y.; Maclean, D.; McNamara, D. J.; Sawyer, T. K.; Dixon, J. E. *Biochemistry* **1994**, *33*, 2285.
- Jia, Z.; Barford, D.; Flint, A. J.; Tonks, N. K. *Science* **1995**, *268*, 1754.
- Krueger, N. X.; Streuli, M.; Saito, H. *EMBO J.* **1990**, *9*, 3241.
- Guan, K. L.; Dixon, J. E. *Science* **1990**, *249*, 553.
- Guan, K.; Deschens, R. J.; Qiu, H.; Dixon, J. E. *J. Biol. Chem.* **1991**, *266*, 12964.
- Directed Mutagenesis*; McPherson, M. J., Ed.; IRL Press: Oxford, U.K., 1991; p 217.

Construction of Val59 → Asp Mutant Gene of YPTP1

The Val-59 to Asp mutation utilized four oligonucleotide primers (CH-44, 46, 50, and 52). CH-44 is a mismatched primer in which the codon GTT for Val is converted to GAT for Asp. CH-50 has the complementary sequence of CH-44. CH-46 and CH-52 are flanking primers 1240 and 1040 base pairs away from the mutation site in pT7-YPTP1.

Three rounds of PCR was performed to prepare the mutant construct. The first round of PCR was performed using CH-44 and CH-46 with pT7-PTP1 plasmid as a template to obtain the PCR product 1. The second round of PCR utilizing primers CH-50 and CH-52 produced the PCR product 2. The third round of PCR utilized the double-stranded PCR products 1 and 2 as templates and CH-46 and CH-52 as primers and the amplified PCR product was ligated with pCRTM2.1 vector (Promega). The ligation product was then digested with Ksp I and Nco I and the 715bp fragment was inserted into the Ksp I/Nco I site of pT7-PTP1 to complete the construction. The MluI/PstI fragment of the mutant construct was then replaced with the same part of wild-type pT7-PTP1 to avoid any undesired mutation. Primers; CH-44 (5'-gCT AgA AAC AgA TAC gAT AAC ATT ATg CCg TAT gAg-3', nucleoties 981-1016 of YPTP1 gene), CH-46 (5'-CgA TAC CgA TAg CAC CTT ggA A-3', antisense of nucleoties 2200-2221 of YPTP1), CH-50(5'-CTC ATA Cgg CAT AAT gTT ATC gTA TCT gTT TCT AgC-3'), CH-52(5'-gAg CgA ACg ACC TAC ACC gAA C-3', nucleoties 896-917 of pT7-7 vector).

Construction of Val59 → His Mutant Gene of YPTP1

The same strategy was used to prepare the construct for the V59H mutant except that the mutant primers CH-56 and CH-57 were used instead of CH-44 and CH-50. In CH-56, the codon GTT for Val is replaced with CAT for Asp. Flanking primers were the same. Primers; CH-56 (5'-C gAT gCT AgA AAC AgA TAC CAT AAC ATT ATg CCg TAT gAg-3'), CH-57 (5'-CTC ATA Cgg CAT AAT

- gTT ATg gTA TCT gTT TCT AgC ATC g-3').
19. Park, S.; Cho, H. *Bull. Korean Chem. Soc.* **1997**, 18, 691.
 20. Han, J.; Kwon, M.; Cho, H. *J. Biochem. Mol. Biol.* **1998**, 31, 135.
 21. Kitas, E. A.; Knorr, R.; Trzeciak, A.; Bannwarth, W. *Helv. Chim. Acta* **1991**, 74, 1314.
-