

Theoretical Studies on the Methanolysis of a Cephalosporin; Mimicking Acylation of the Active Site Serine of D-Ala-D-Ala Transpeptidases

Keeyung Nahm

Specialty Chemicals, R & D Center, LUCKY LTD., P. O. Box 10, Science Town, Taejeon 305-343.

Received July 10, 1991


Methanolysis of a β -lactam ring of a cephalosporin was simulated with AM1 semiempirical quantum mechanical calculation. The tetrahedral intermediate **TD1** from an O-protonated cephalosporin and a methanol transfers the proton intramolecularly to the C-4 carboxylate to generate an oxyanion, *i.e.*, second tetrahedral intermediate **TD2**, which undergoes the amide bond cleavage without further protonation on the N-5. For this cleavage a low-energy barrier **TS2** was located. According to the energy diagram, tetrahedral intermediates easily undergo ring cleavage even without the protonation on the amide nitrogen.

Introduction

The active site of glycopeptide D-Ala-D-Ala transpeptidases of bacteria is now known to have a serine unit and a generally positively charged area,¹ which is similar to those of the serine proteases.² It is widely believed that the addition of the hydroxy group of the serine to the amide of a β -lactam ring initiates the acting mode of β -lactam antibiotics and finally the serine at the active site is acylated with β -lactam antibiotics, which reduce or terminate the activities of the D-Ala-D-Ala transpeptidase.² As a result, the formation of cross linkages in the peptidoglycans of bacteria can not occur and bacteria will finally burst out because of the weak cell wall.² The major difference between the D-Ala-D-Ala peptidase and the classical serine protease family is that the former has not a well-known Asp-His-Ser triad, which is an essential active site of the serine protease family to activate the serine hydroxy group.³ Almost all known penicillin-binding-proteins do not have even the histidines around the active sites.^{1c,4}

The chemical processes involved in the acting mode of β -lactam antibiotics are very similar to the hydrolysis of an amide which is well studied experimentally and theoretically.⁵ In solution phase, the protonated carbonyl of an amide is attacked by a hydroxide to form a tetrahedral intermediate which is further cleaved to a carboxylate and an amine. Existence of this tetrahedral intermediate is very controversial,⁶ but in solution these intermediates have been detected experimentally.⁷

Antibiotic activities of β -lactam antibiotics, such as penicillins and cephalosporins, have been thought to be originated from strain of the four-membered ring and reduced amide resonance.^{2b,8} However, it is now generally accepted that the biological or chemical reactivity of β -lactam antibiotics is not determined solely by the strain or the reduced amide resonance.⁹ The studies on the serine protease reveal that the active site serine is more reactive to the substrates or the substrate-like than other serines in the enzymes.^{2a} Therefore, the amide bond in the β -lactam ring or the hydroxyl group or the serine should be activated to form a tetrahedral intermediate either by protonation of the amide bond (hydrogen bonding donor) or by deprotonation of the



serine hydroxyl group (hydrogen bonding acceptor). The candidates for the protonating agents are His and Lys, and those for the deprotonating are His and Glu, etc. Another major feature of the antibiotic, the carboxylate at C-3 in penicillins and at C-4 in cephalosporins has been studied actively and has been postulated to participate in transpeptidase inhibition or β -lactamase catalysis by acting as a counterpart of an electrostatic anchor, such as Lys.¹⁰

However, it is of great difficulty to monitor the whole actual mode. Here, we report the results of the semiempirical calculation on the hypothetical methanolysis of a cephalosporin by using AMPAC program.¹¹ We describe the protonated cephalosporins, formation of tetrahedral intermediates with methanol, and the breakdown of the intermediates.

Calculation methods.

Semiempirical molecular orbital calculation has been performed with the AMPAC package and AM1 parameter has been used throughout this calculation.¹¹ The cephalosporin (1) studied here has an acetylamino group at C-7 but no leaving substituent at C-3. The NH proton of the C-7 acetylamino group is located above the β -lactam ring. The attacking direction of the methoxide anion could be either *cis* or *trans* to the C₆-S₁ bond, but in this simulation study only the *trans* direction was studied considering the transpeptidase enzyme reaction (Scheme 1). Every atoms was set free for the full

Table 1. Heat of Formation of the Protonated Cephalosporins (in kcal/mol)


	H_f	Length				Torsional angle		
		O ₉ -C ₈	C ₈ -N ₅	N ₅ -C ₆	O ₉ -C ₁₀	C ₃ -C ₄	C ₈ -N ₅ -C ₆ -C ₄	C ₆ -S ₁ -C ₂ -C ₃
Cepha (2)	-41.14	1.274	1.380	1.476	3.274	1.352	146.4	-43.2
Cepha (3)	-41.69	1.198	1.553	1.508	3.052	1.340	119.2	-42.2
Cepha (4)	-96.99	1.219	1.438	1.470	3.310	1.358	138.6	-39.8

Table 2. Geometric Informations of Reactants, Intermediates, Products, and Transition States (bond lengths in angstrom, angles in degree)

	Length					Torsional angle	
	O ₉ -C ₈	C ₈ -N ₅	N ₅ -C ₆	O ₉ -C ₁₀	C ₃ -C ₄	C ₈ -N ₅ -C ₆ -C ₄	C ₆ -S ₁ -C ₂ -C ₃
Cephalosporin (1)	1.219	1.422	1.468	3.416	1.354	143.4	-35.8
TD1	1.390	1.504	1.467	3.313	1.355	133.2	-39.0
TD2	1.288	1.584	1.454	3.322	1.364	137.4	-38.1
TS1	1.339	1.535	1.465	3.021	1.359	129.8	-43.7
TS2	1.265	1.811	1.436	3.470	1.368	136.5	-35.1
Prod (1)	1.231	3.027 ^a	1.402	5.246 ^a	1.377	-	-29.0
Prod (2)	1.230	3.028 ^a	1.421	5.151 ^a	1.354	-164.4 ^b	-28.3

^acompletely cleaved bond. ^b ω (H-N₅-C₆-C₄).

optimization, and the "PRESICE" key word was used. Vibrational frequencies of all intermediates and products were checked to see all positive frequencies and all transition structures were confirmed by observing only one negative frequency.



Results and Discussion

Protonated Cephalosporins. A calculated cephalosporin **1** has a similar structure of a β -lactam ring and a dihydrothiazine ring to those determined by crystallographic methods.¹² The C-4 carbon was deviated from the plane of N₅-C₆-C₈ by 36.4°. The chain of O₉-C₈-N₅-C₄-C₃ was composed by the bonds of 1.219, 1.421, 1.397, and 1.354 Å, respectively. A normal peptide bond has an amide (O=C-N-C) of 1.24, 1.32, and 1.46 Å.^{2a} It is clear that cephalosporin **1** has a less conjugated amide bond. Especially the C-N bond length of **1** is similar to a C-N single bond (1.49 Å) because of the β -lactam ring strain. The enamine conjugation in the C₃-C₄-N₅ bond is believed to shorten the C₄-N₅ bond, which should decrease the electron density at N-5. This also should decrease the amide resonance on the β -lactam ring.

Three protonated cephalosporins were considered. The first one is an O-protonated cephalosporin (**2**) which has two conformation depending on the direction of the proton. The second one is an N-protonated (**3**) and the last one is a protonated carboxylic cephalosporin (**4**). Although the last one is most stable, we excluded this form from further studies

because it seems to be equivalent to the unprotonated **1** in the amide methanolysis. The calculated heat of formation of an O-protonated, an N-protonated, and a carboxylic acid form are -41.14, -41.69, and -96.99 kcal/mol, respectively. The detailed parameters are written on Table 1.

Methanolysis of the O-Protonated Cephalosporin.

A proposed scheme for the methanolysis of the O-protonated cephalosporin **2** was drawn in the Scheme 1. The total charge density around the ring carbonyl carbon of the O-protonated cephalosporin, **2**, was calculated to be more positive compared to the unprotonated **1** by +0.632. Not surprisingly, the proton on the carbonyl oxygen was located toward the C-4 carboxylate because of electrostatic attraction, which has been maintained throughout the methanolysis. The geometrical differences between the unprotonated and the protonated are as follows; on protonation, the carbonyl bond length is elongated by 0.055 Å, but the C₈-N₅ bond is shortened by 0.042 Å.


Tetrahedral intermediate, **TD1**, is expected to be formed from **2** and a methoxide anion. Inside the peptidase enzymes, formation of **TD1** would be a lot more complicated; the hydroxy group of the active serine will attack the ring amide bond with assistance from the surroundings, which should lower the activation energy for the **TD1** formation. The C₈-N₅ bond in **TD1** is 1.504 Å, which is not an amide bond any more, and the O₉-C₈ bond is an almost normal hydroxyl bond (1.390 Å). The proton on the O-9 of **TD1** is directing toward the C-4 carboxylate, and in this conformation one of the lone electron pairs of the O-9 is in the antiperiplanar position of the C₈-N₅ bond.

The second tetrahedral intermediate, **TD2**, which has an O-9 anion and a protonated C-4 carboxylic group, were located. The shorter O₉-C₈ bond (1.288 Å) compared to that of the **TD1** indicates that the electron on the O-9 was moving to the N-5 through the O₉-C₈-N₅ bond, as a consequence,

Table 3 Heat of Formation and Reaction of Reactants, Intermediates, Products, and Reactions (in kcal/mol)

Intermediates and reactions	H_f or H_{rxn}	ΔE_{act}
CH ₃ OH	-57.03	
CH ₃ O ⁻	-38.50	
Cephalosporin (1)	-127.28	
TD1	-200.60	
TD2	-198.71	
Prod(1)	-217.32 ^a	
Prod(2)	-217.18 ^b	
TS1	-191.80	
TS2	-196.73	
CH ₃ OH + 1 → TD1	-16.29	
TD1 → TD2	+1.89	+8.80
TD2 → Prod(1)	-18.61	+1.98
Prod(1) → Prod(2)	+0.24	


^a $\omega(N_5-C_6-C_7-C_8) = -70.9^\circ$. ^b $\omega(N_5-C_6-C_7-C_8) = -72.2^\circ$.

**Figure 1.** Various protonated cephalosporins.

the C₈-N₅ bond was elongated to 1.584 Å.

The approaching direction of the methanol (a serine equivalent) in both TD1 and TD2 was fixed only to the β-side (*trans* to the sulfur) as shown in Figure 1, which is well known from crystallographic studies.^{1c} However, in solution phase, the methanol should approach to either the α or β side.^{2b} The position of the methoxy group obeys the stereoelectronic effect; *i.e.*, either one of the nonbonding electron pairs is located to the antiperiplanar position of the C₈-N₅ bond and therefore the methyl has the staggered position to the bond.

TD2 could be formed from TD1 by many ways; *i.e.*, protonation and deprotonation by external catalysts. However, TD1 and TD2 are similar in energies (TD1 has a slightly lower energy compared to TD2 by 1.9 kcal/mol) and the C-4 carboxylate group can participate in the proton transfer from TD1 to TD2. In that sense, we located the transition state, TS1, between two tetrahedral intermediates. It has a proton partially attached between the oxygens of an oxide and a carbo-

**Figure 2.** Various reactants, intermediates, and transition structures.

xylate, and is located above TD1 by 8.8 kcal/mol. The corresponding C₈-N₅ and C₈-O₉ bonds are 1.535 and 1.339 Å, respectively, and their values are somewhat inbetween those of TD1 and TD2. Our calculation results show that the C-4 carboxylic group in TS1 behaves as an internal catalyst. Recent experimental work also shows that a protonated tetrahedral intermediate (TD1) should convert an unprotonated tetrahedral intermediate (TD2) to undergo methanolysis.¹³

In the amide bond cleavage in the intermediate TD2 to yield a product (Prod1), a transition state (TS2) was located and this process was calculated to have a barrier of 1.98 kcal/mol. The angles of <C₈-C₇-C₆ and <C₇-C₆-N₅ are 93.3° and 95.9°, respectively, and the β-lactam ring strain has been relieved by 5°. The torsional angle of $\omega(C_6-S_1-C_2-C_3)$ was calculated to be -35.1° in this transition state. The bond length of C₈-N₅ is 1.811 Å, which is longer than those of 2, TD1, and TD2 by 0.2-0.3 Å. Otherwise, TS2 and TD2 are very similar in geometry; for examples, the torsional




Figure 3. Energy diagram for the methanolysis of the O-protonated cephalosporin (units in kcal/mol, not scaled).

angles of $\omega(C_4-N_5-C_6-C_7)$ and $\omega(S_1-C_6-C_7-C_8)$ are -140.0° and -120.9° for **TS2**, and -141.5° and -119.4° for **TD2**. The similarity in geometry of **TS2** with **TD2**, not with **Prod1**, is also expected from the low energy barrier for the conversion to **Prod1**. It should be emphasized that the amide bond cleavage of **TD2** can occur without protonation on N-5, which is the major difference from the ordinary amide hydrolysis.^{5c} Almost all known cephalosporins have leaving groups on C-3 (C_3-CH_2-X) and they may not need protonation on N-5 to repel the leaving group (X) during the methanolysis with a serine residue in DD-prptidases.

Prod1 has a planar ester bond around C-8. The dihydrothiazine ring was rotated away from the methoxy group on C-8 to have $C_8-C_7-C_6-C_5$ torsional angle of -70.9° , which is well correlated with crystallographic data.^{1c} Charge densities on N-5 were changed from -0.206 in **1** to -0.308 in **TS2**, and finally to -0.438 in **Prod1**. Those on C-3 also were changed from -0.304 in **1** to -0.344 in **TS2**, and to -0.489 in **Prod1**. The negative charge created by the amide bond cleavage is observed to be stabilized by being distributed over the C-3 carbon, which is one of the major differences between the cephalosporin methanolysis and the normal amide methanolysis.^{9a} There are some previous studies on the leaving group effect at the C-3 position.¹⁴ From our charge density study on **Prod1** also shows that when there is a leaving group at the C-3 position, the leaving group will be easily detached from the C-3 position.

Prod2 is expected to be the final product in the methanolysis and has a $\omega(C_8-C_7-C_6-N_5)$ torsional angle of -72.2° . Even though there is another conformer which has slightly higher energy (by 3.6 kcal/mol) than **Prod2** and has the torsional angle of 42.8° , it was not considered because only **Prod2** fits the cavity suggested by the crystallographic work.^{1c}

In conclusion, the proton on the β -lactam amide oxygen can be removed by the C-4 carboxylate group to give the cleavage-ready tetrahedral intermediate **TD2**, which has an oxyanion. The amide cleavage has been calculated to have a low activation energy, and the negative charge on the N_5 of **Prod1** stabilized well by the $C_3=C_4$ double bond.

The whole processes involved in the deactivation of the transpeptidase by β -lactam is not clear yet and it will take more time to figure out the exact mechanism. Because even modern X-ray crystallography and kinetic studies can not pro-

vide the exact pictures of acting modes of drugs, it is important to consider the experimental results in conjunction with theoretical data

Acknowledgement. I am indebted very much to Dr. Hunseung Oh. I thank also to Dr. Yong-Zu Kim and Dr. Sung Chun Kim for their helpful discussions.

References

- (a) J. A. Kelly, P. C. Moews, J. R. Knox, J. M. Frere, and J. M. Ghuysin, *Science*, **218**, 479-481 (1982); (b) J. A. Kelly, J. R. Knox, P. C. Moews, G. J. Hite, J. B. Bartolone, H. Zhao, B. Joris, J. M. Frere, and J. M. Ghuysen, *J. Biol. Chem.*, **260**, 6449-6458 (1985); (c) J. A. Kelly, J. R. Knox, H. Zhao, J.-M. Frere, and J. M. Ghuysen, *J. Mol. Biol.*, **209**, 281-295 (1989).
- (a) L. Stryer, *Biochemistry*; 3rd ed, Freeman: New York, 1988; pp 195-198; (b) D. B. Boyd, in *Chemistry and Biology of β -lactam Antibiotics*; ed. R. B. Morin and M. Gorman, Vol. 1, Academic Press: New York, NY, 1982.
- (a) B. W. Matthews, P. B. Sigler, R. Henderson, and D. M. Blow, *Nature (London)*, **214**, 652 (1967); (b) D. M. Blow, J. J. Birktoft, and B. S. Hartley, *Nature (London)* **221**, 337 (1969); (c) D. M. Blow, *Acc. Chem. Res.*, **9**, 145 (1976); (d) R. M. Stroud, *Sci. Am.*, **231**(1), 24 (1974); (e) H. Tsukada and D. M. Blow, *J. Mol. Biol.*, **184**, 703 (1985).
- (a) R. P. Ambler, in *Beta-Lactamases*; ed. by J. M. T. Hamilton-Miller and J. T. Smith; Academic Press, New York, 1979; pp 99-125. (b) B. Joris, J. M. Ghuysen, G. Dive, A. Renard, O. Dideberg, P. Charlier, J. M. Frere, J. A. Kelly, J. C. Boyington, P. C. Moews, and J. R. Knox, *Biochem. J.*, **250**, 313-324 (1988).
- Experimentally, see (a) T. C. Bruice and S. J. Benkovic, *Bioorganic Mechanisms*; Wiley: New York, 1966; (b) R. J. E. Talbot, in *Comprehensive Chemical Kinetic*; C. H. Bamford and C. F. H. Tipper eds., Elsevier: Amsterdam, 1972. Theoretically, for examples, see (c) S. J. Weiner, U. C. Singh, and P. A. Kollman, *J. Am. Chem. Soc.*, **107**, 2219-2229 (1985); (d) D. Arad, R. Langridge, and P. A. Kollman, *ibid.* **112**, 491-502 (1990); (e) I. Lee, C. K. Kim, and H. S. Seo, *Tetrahedron*, **42**, 6627-6633 (1986).
- O. I. Asubiojo and J. I. Brauman, *J. Am. Chem. Soc.*, **101**, 3715-3724 (1979).
- (a) M. L. Bender, *Chem. Rev.*, **60**, 53 (1960); (b) S. L. Johnson, *Adv. Phys. Org. Chem.*, **5**, 237 (1967); (c) M. Cocivera, *et al.*, *J. Am. Chem. Soc.*, **98**, 1573, 7366, 7371, 7362 (1976).
- S. N. Rao and R. A. More O'Ferrall, *J. Am. Chem. Soc.*, **112**, 2729-2735 (1990).
- (a) M. I. Page, *Adv. Phys. Org. Chem.*, 165 (1987); (b) M. I. Page, P. Webster, S. Bogdan, B. Tremier, and L. Ghosez, *J. Chem. Soc., Chem. Commun.*, 1039 (1986).
- (a) O. Herzberg, J. Moul, *Science*, **236**, 694-701 (1987); (b) L. M. Ellerby, W. A. Escobar, A. L. Fink, C. Mitchinson, and J. A. Wells, *Biochemistry*, **29**, 5797-5806 (1990).
- M. J. S. Dewar, E. G. Zeolish, E. F. Healy, and J. J. P. Stewart, *J. Am. Chem. Soc.*, **107**, 3902-3909 (1985); (b) AMPAC, Version 2.1, QCPE 506, Indiana University.
- For X-ray structure of cephalosporins, see R. M. Sweet, L. F. Dahl, *J. Am. Chem. Soc.*, **92**, 5489-5507 (1970).

13. K. J. Davis and M. I. Page, *J. Chem. Soc., Chem. Commun.*, 1448-1450 (1990).
 14. For examples, see, (a) M. I. Page, *Acc. Chem. Res.*, **17**,

144-151 (1984); (b) D. B. Boyd, *J. Org. Chem.* **50**, 886-888 (1985).

Theoretical Studies on the Gas-Phase Wittig-Oxy-Cope Rearrangement of Deprotonated Diallyl Ether¹

Chang Kon Kim, Ikchoon Lee*, Hai Whang Lee, and Bon-Su Lee

Department of Chemistry, Inha University, Incheon 402-751. Received July 18, 1991



The Wittig-oxy-Cope rearrangements of deprotonated diallyl ether, I, $\text{CH}_2=\bar{\text{C}}\text{H}-\text{CH}-\text{O}-\text{CH}_2-\text{CH}=\text{CH}_2$, have been investigated theoretically by the AM1 method. A two step mechanism forming a Wittig product ion, II, $(\text{CH}_2=\text{CH})(\text{CH}_2=\text{CH}-\text{CH}_2)\text{CHO}^-$, through a radical-pair intermediate was found to provide the most favored reaction pathway in the Wittig rearrangement. The subsequent oxy-Cope rearrangement from species II also involves a two step mechanism through a biradicaloid intermediate. The intramolecular proton transfer in I (to form $\text{CH}_2=\text{CH}-\text{CH}_2-\text{O}-\bar{\text{C}}\text{H}-\text{CH}=\text{CH}_2$) is a higher activation energy barrier process compared to the Wittig and oxy-Cope rearrangements and is considered to be insignificant. These results are in good agreement with the condensed-phase as well as gas-phase experimental results.

Introduction

The Wittig rearrangement (WR)² has been extensively studied as one of the better known carbanion rearrangement in the condensed phase. Two-step mechanisms (reaction pathways *a* and *b*) involving either of the intermediate shown in Scheme 1 are proposed for the rearrangement.³ However the possibility of a concerted one-step process (reaction pathways *c*) can not be entirely ruled out.⁴ Various experimental results seem to support the two-step mechanism involving a radical pair, pathway *b*, rather than that involving an ionic intermediate, pathway *a*.⁵ Some of the condensed-phase experimental results in favor of the radical-pair mechanism are: (i) the migratory aptitude of substituents R^2 is in the order of free-radical stabilities,⁶ allyl \cong benzyl > methyl > ethyl > phenyl, (ii) partial racemization of R^2 is observed,⁷ (iii) when ketyl radicals and R radicals from different precursors were brought together, similar products resulted.⁸ Despite these experimental evidence in support of the radical-pair intermediate pathway, *b*, the radical mechanism is unable to account for all the reaction products and a concerted mechanism, *c*, is suggested as a possible alternative.⁴ It is also known that when R^2 is an allyl group, the [2,3] sigmatropic rearrangement can take place.⁹

The Wittig rearrangement can also occur in the gas-phase.¹⁰ Eichinger *et al.*,^{10a} have shown in their gas-phase studies on the Wittig rearrangement of deprotonated diallyl ether that (i) the proton-transfer reaction I to I' (Scheme 2) is insignificant, (ii) I undergoes facile reaction leading to 1,2- and 1,4-rearrangement products, (iii) the Wittig ions rearrange further by an oxy-Cope mechanism (OCR)¹¹ (Scheme 2).

In this work, we explore MO theoretically the most possi-



ble reaction pathway for the Wittig rearrangement of deprotonated diallyl ether (Scheme 1) and the mechanism of the subsequent oxy-Cope rearrangement of the Wittig product ion, II (Scheme 2).

Computation

The AM1 procedure¹² implemented in AMPAC package¹³ was used throughout in this work. The AM1 method accommodates some electron correlation effect through its parametrization¹⁴ and it requires 2-3 orders of magnitude lesser computing time than even those using the relatively low level ab initio (3-21G) method.^{12,14} It has been reported that AM1 gives good results for reactions of anionic species by giving

*Author for correspondence