

Viscosity and Diffusion of Small Normal and Isomeric Alkanes: An Equilibrium Molecular Dynamics Simulation Study

Choong Do Yoo, Soon-Chul Kim,[†] and Song Hi Lee^{*}

*Department of Chemistry, Kyungsoong University, Busan 608-736, Korea. *E-mail: shlee@ks.ac.kr*

[†]Department of Physics, Andong National University, Andong 760-749, Korea

Received February 28, 2008

Key Words : Diffusion, Viscosity, Normal alkanes, Isomeric alkanes, Molecular dynamics simulation

Constitutional isomers¹ have different physical properties that have the same carbon number but different structures. The difference may not be large, but they are found to have different melting points, boiling points, densities, indexes of refraction, and so forth. For example, a branched chain isomer has a lower boiling point than a straight chain isomer. Thus normal pentane has a boiling point of 36°C, isopentane with a single branch 28°C, and neopentane with two branches 9.5°C. This effect of branching on boiling point is observed within all families of organic compounds. It is reasonable that branching should lower the boiling point: with branching the molecular shape tends to approach to that of a sphere and as this happens the surface area decreases, as the result of that the intermolecular forces become weaker and are overcome at a lower temperature.

The branching effect on the dynamic properties of liquid alkanes, such as self-diffusion constant, viscosity, and thermal conductivity, is one of the most interesting phenomena. For liquid butane, the experimentally observed behavior² tells us that viscosity increases with branching. For liquid pentane, hexane, and heptane, however, branching decreases the viscosity, for example, 0.289 and 0.273 cp at 273.15 K, and 0.240 and 0.223 cp at 293.15 K for normal pentane and isopentane, 0.326 and 0.306 cp at 293.15 K for normal hexane and isohexane, and 0.409 and 0.384 cp at 293.15 K for normal heptane and isoheptane,³ respectively. These experimental results, except for liquid butane, indicate that as the molecular shape tends to approach that of a sphere and the surface area tends to decrease with branching, the intermolecular forces become weaker and the viscosity of alkane isomers decreases.

In the present note, we report equilibrium molecular dynamics (MD) simulations for the systems of small normal and isomeric alkanes - normal butane and isobutane, normal pentane and isopentane, and normal hexane and isohexane. The primary study goal is to analyze the diffusion and viscosity dynamics of small normal and isomeric alkanes at different temperatures.

Molecular Models and MD Simulation Methods

For small normal alkanes, we have chosen 3 systems - normal butane (C₄H₁₀), normal pentane (C₅H₁₂), and normal hexane (C₆H₁₄), and the corresponding isomeric alkane

systems are isobutane (C₄H₁₀), isopentane (C₅H₁₂), and isohexane (C₆H₁₄). Each simulation was carried out in the NVT ensemble; the number of n-alkane was N=100 and the lengths of cubic simulation boxes were obtained from the experimental densities⁴ for given temperatures of 248, 273, 293 and 298 K. The usual periodic boundary condition in the x-, y-, and z-directions and the minimum image convention for pair potential were applied. Gaussian isokinetics was used to keep the temperature of the system constant.⁵

We used a united atom (UA) model for n-alkanes, that is, methyl and methylene groups are considered as spherical interaction sites centered at each carbon atom. This model was used in the previous simulation studies.⁶⁻¹⁰ Here, we briefly describe the salient features of the model. The interaction between the sites on different n-alkane molecules and between the sites separated by more than three bonds in the same n-alkane molecule was described by a Lennard-Jones (LJ) potential. All the sites in a chain have the same LJ size parameter $\sigma_i \equiv \sigma_{ii} = 3.93$ Å, and the well depth parameters were $\epsilon_i \equiv \epsilon_{ii} = 0.94784$ kJ/mol for interactions between the end sites and $\epsilon_i = 0.39078$ kJ/mol for interactions between the internal sites. The Lorentz-Berthelot combining rules [$\epsilon_{ij} \equiv (\epsilon_i \epsilon_j)^{1/2}$, $\sigma_{ji} \equiv (\sigma_i + \sigma_j)/2$] were used for interactions between an end site and an internal site. A cut-off distance of $2.5\sigma_i$ was used for all the LJ interactions.

Initially the bond-stretching was described by a harmonic potential, with an equilibrium bond distance of 1.54 Å and a force constant of 1882.8 kJ/mol/Å². The bond bending interaction was also described by a harmonic potential with an equilibrium angle of 114° and a force constant of 0.079187 kJ/mol/degree². The torsional interaction was described by the potential developed by Jorgensen *et al.*¹¹:

$$U_{torsion}(\phi) = a_0 + a_1 \cos \phi + a_2 \cos^2 \phi + a_3 \cos^3 \phi, \quad (1)$$

where ϕ is the dihedral angle, and $a_0 = 8.3973$ kJ/mol, $a_1 = 16.7862$ kJ/mol, $a_2 = 1.1339$ kJ/mol, and $a_3 = -26.3174$ kJ/mol. For the time integration of the equations of motion, we adopted Gear's fifth-order predictor-corrector algorithm¹² with a time step of 0.5 femto-second for all the systems. Later the bond-stretching was switched to a constraint force which keeps intramolecular nearest neighbors at a fixed distance. The advantage for this change is to increase the time step as 5 femto-seconds with the use of RATTLE algorithm.¹³ After a total of 1,000,000 time steps (5 nano-

seconds) for equilibration, the equilibrium properties were then averaged over 5 blocks of 200,000 time steps (1 nano-seconds). The configurations of all the molecules for further analyses were stored every 10 time steps (0.05 pico second) which is small enough for the tick of any time auto-correlation functions.

Self-diffusion constant (D) can be obtained through two routes: the Green-Kubo formula from velocity auto-correlation functions (VAC):

$$D = \frac{1}{3} \int_0^{\infty} \langle \mathbf{v}_i(t) \cdot \mathbf{v}_i(0) \rangle dt. \quad (2a)$$

and the Einstein formula from mean square displacements (MSD):

$$D_s = \frac{1}{6} \lim_{t \rightarrow \infty} \frac{d \langle |\mathbf{r}(t) - \mathbf{r}(0)|^2 \rangle}{dt}. \quad (2b)$$

Shear viscosity is calculated by a modified Green-Kubo formula for a better statistical accuracy^{14,15}:

$$\eta = \frac{V}{kT} \int_0^{\infty} dt \sum_i \langle P_{i\alpha\beta}(0) P_{i\alpha\beta}(t) \rangle, \quad (3)$$

where $P_{i\alpha\beta}$ is the $\alpha\beta$ component of the molecular stress tensor, \mathbf{P}_i , of particle i by translational motion:

$$P_{i\alpha\beta}(t) = \frac{1}{V} \left[m v_{i\alpha}(t) v_{i\beta}(t) + \sum_{j \neq i} r_{ij\alpha}(t) f_{ij\beta}(t) \right], \quad (4)$$

where $\alpha\beta = xy, xz, yx, yz, zx, \text{ or } zy$.

Results and Discussion

We compare the results of self-diffusion constants of normal alkanes and isomeric alkanes. In Figure 1 the log-log plot of self-diffusion constant (D) versus molecular mass (M) is shown for several temperatures. Calculated self-diffusion constants of liquid isobutane are larger than those of liquid normal butane over four different temperatures while the opposite is observed for liquid pentane and




Figure 1. A log-log plot of D (10^{-6} cm²/sec) vs. M (g/mol). From top, $T = 298$ (\blacktriangle), 293 (\blacklozenge), 273 (\blacksquare), and 248 K (\bullet), respectively. Black: normal alkanes and white: isomeric alkanes.

hexane. The crosses of the slope lines in Figure 1 represent this. The slopes are almost linear for given temperatures, which indicates that the behavior of D vs. M is well described by $D \sim M^{-\alpha}$. The obtained exponents are between 1.25 and 1.36 for liquid normal alkanes and between 1.45 and 1.56 for liquid isomeric alkanes. At the molecular weight of the Rouse regime (the polymer chain dynamics of unentangled chains is commonly described by the Rouse), D of normal alkanes also show power law behaviors. For example, D of normal alkanes for normal octane to polyethylene of the molecular weight of several thousands was reported that the exponents α is in the range of 1.75-2.72 depending on temperature.¹⁶⁻¹⁸ Our previous MD simulation studies¹⁹ for normal C₁₂-C₄₄ at $T = 273$ - 473 K are another examples. The obtained exponents are between 1.6 and 2.4. Apparently the exponent in liquid alkanes decreases with temperature.

Viscosities of liquid normal and isomeric alkanes obtained from the modified Green-Kubo relation, Eq. (3), underestimate the experimental values: comparing in the unit of cp, 0.289/0.193 at 273.15 K and 0.240/0.147 at 293.15 K for normal pentane, 0.273/0.133 at 273.15 K and 0.223/0.106 at 293.15 K for isomeric pentane, 0.401/0.297 at 273.15 K and 0.326/0.226 at 293.15 K for normal hexane, and 0.376/0.266 at 273.15 K and 0.306/0.182 at 293.15 K for liquid isomeric hexane, respectively. The difference is severe in isomeric alkanes. Figure 2 shows the log-log plot of viscosity (η) versus molecular mass (M). Calculated viscosities of liquid isomeric alkanes are always smaller than those of liquid normal alkanes over four different temperatures. The slopes are almost linear for given temperatures, which indicates that the behavior of η vs. M is well described by $\eta \sim M^{\beta}$. The obtained exponents are between 2.32 and 2.47 for liquid normal alkanes and between 1.77 and 1.98 for liquid isomeric alkanes. The experimental results for normal alkanes and linear polyethylene¹⁷ show that η is also well described by the power law: $\eta \sim M^{1.8}$ at low molecular weight ($M < 5$ kg/mol) and $\eta \sim M^{3.6}$ at high molecular weight ($M > 5$ kg/




Figure 2. A log-log plot of η vs. M (g/mol). From top, $T = 248$ (\bullet), 273 (\blacksquare), 293 (\blacklozenge), and 293 K (\blacktriangle), respectively. Black: normal alkanes and white: isomeric alkanes.

mol) at 448 K. Our previous MD simulation studies¹⁹ for normal C₁₂-C₄₄ at T = 273-473 K gives the obtained exponents between 2.0 and 3.7. Assumably the exponent in liquid normal alkanes decreases with temperature in the same direction to the exponent in the log-log plot of self-diffusion constant (D) versus molecular mass (M) discussed above.

Experimental measurements show that the viscosity of liquid isomeric butane is higher than that of liquid normal butane at 298 K (0.166/0.162 cp) and also for gas state (76.14/75.43 mp).⁴ However, branching decreases the viscosity for liquid pentane, hexane, and heptane as discussed above. The simulation results underestimate the experimental values for both liquid isomeric and normal butanes (0.0632/0.0836 cp), and show that branching decreases the viscosity in contradiction to the experimental trend. A similar result for the viscosity of liquid butanes was reported in the non-equilibrium MD simulation using UA model for liquid butanes.²⁰

Usually the behavior of self-diffusion constant (D) in long chains of alkanes is the opposite to that of viscosity (η). For liquid butane, calculated self-diffusion constants (D) of normal alkanes are smaller than those of isomeric alkanes and the opposite is observed for calculated viscosities (η). However, exceptionally both calculated self-diffusion constants (D) and viscosities (η) of liquid normal pentane and hexane are larger than those of liquid isomeric alkanes. From Figures 1 and 2, the increase in the exponent of the log-log plot of self-diffusion constant (D) versus molecular mass (M) from normal alkanes to isomeric alkanes (1.3 \rightarrow 1.5, averagely) agrees with the decrease in the exponent of the log-log plot of viscosity (η) versus molecular mass (M) (2.4 \rightarrow 1.9, averagely) and the product of two exponents are almost constant (3.1 vs. 2.9).

The temperature dependence of the calculated diffusion constants of liquid n-alkanes over the whole temperatures considered is suitably described by an Arrhenius plot, $D = D_0 \exp(-E_D/RT)$, as shown in Figure 3, where D_0 is the pre-




Figure 3. Arrhenius plot of D (10^{-6} cm²/sec) vs. 1/T. From top, C₄ (●), C₅ (■), and C₆ (◆), respectively. Black: normal alkanes and white: isomeric alkanes.




Figure 4. Arrhenius plot of η (cp) vs. 1/T. From top, C₆ (◆), C₅ (■), and C₄ (●), respectively. Black: normal alkanes and white: isomeric alkanes.

exponential factor, RT has the usual meaning, and E_D is the activation energy of normal alkane diffusion. The value of the activation energy is a direct measure of how fast the self-diffusion changes with temperature. The activation energies obtained from the slope of the least square fit are 1.84, 1.91, and 1.97 kcal/mol for liquid normal alkanes and 1.90, 1.95, and 2.05 kcal/mol for liquid isomeric alkanes, respectively. The activation energies of normal alkane diffusion are smaller than those of isomeric alkane diffusion. Our previous MD simulation studies¹⁹ for normal C₁₂-C₄₄ at T = 273-473 K gives the activation energies of 2.83, 3.52, 3.91, and 4.06 kcal/mol for C₁₂, C₂₀, C₃₂, and C₄₄, respectively. E_D is small for small alkanes.

We also show the temperature dependence of the calculated viscosities of liquid n-alkanes over the temperatures which is also suitably described by an Arrhenius plot, $\eta = \eta_0 \exp(E_\eta/RT)$, as shown in Figure 4: where η_0 is the pre-exponential factor and E_η is the activation energy of n-alkane viscosity. The activation energies obtained from the slope of the least square fit are 2.03, 2.11, and 2.18 kcal/mol for liquid normal alkanes and 1.73, 1.91, and 1.97 kcal/mol for liquid isomeric alkanes, respectively. The activation energies of normal alkane viscosity are larger than those of isomeric alkane viscosity, which is opposite to that of diffusion. Our previous MD simulation studies¹⁹ for normal C₁₂-C₄₄ at T = 273-473 K gives the activation energies of 2.33, 3.29, 4.63, and 5.46 kcal/mol for C₁₂, C₂₀, C₃₂, and C₄₄, respectively. E_η is also small for small alkanes. It was experimentally reported for n-alkanes and linear polyethylene¹⁷ that the activation energy increases with chain length and at the highest molecular weight tested (M ~ 4.4 kg/mol) the activation energy reaches 6.6 kcal/mol, which is similar to the average value found for NBS (National Bureau Standard) 1482-4 (6.7 kcal/mol) and the values reported by others^{21,22} for high molecular weight linear polyethylene (6.1-6.9 kcal/mol). As chain length n increases the increment of E_η decreases, and it is expected to approach an asymptotic value as n increases to the higher values.

In summary, self-diffusion constants (D) of liquid isobutane obtained in this study are larger than those of liquid normal butane over four different temperatures while the opposite is observed for liquid pentane and hexane. Viscosities (η) of liquid normal alkanes are larger than those of liquid isomeric alkanes, which agrees with the experimental result except liquid butane. The usual opposition in the trends of self-diffusion constant and viscosity is not applied to small liquid alkanes.

Acknowledgments. This work was supported by Grant No. R01-2003-000-10327-0 from the Basic Research Program of the Korea Science and Engineering Foundation Grant. This research is a partial fulfillment of the requirements for the degree of Ph. D. of Science for CDY at Department of Chemistry, Graduate School, Kyungshung University.

References

1. Graham Solomon, T. W. *Organic Chemistry*, 4th ed.; John Wiley and Sons: New York, 1988.
2. Diller, D. E.; van Poolen, L. J. *Int. J. Thermophysics* **1985**, *6*, 43.
3. Weast, R. C.; Astle, M. J. *CRC Handbook of Chemistry and Physics*, 63rd ed.; CRC press: Boca Ranton, 1982-1983.
4. Gallant, R. W. *Physical Properties of Hydrocarbons*; Gulf Pub.: Houston, 1968.
5. Evans, D. J.; Hoover, W. G.; Faylor, B. H.; Moran, B.; Ladd, A. J. *C. Phys. Rev.* **1983**, *A28*, 1016. (b) Simmons, A. J. D.; Cummings, P. T. *Chem. Phys. Lett.* **1986**, *129*, 92.
6. Siepmann, J. I.; Karaborni, S.; Smit, B. *Nature (London)* **1993**, *365*, 330.
7. Smit, B.; Karaborni, S.; Siepmann, J. I. *J. Chem. Phys.* **1995**, *102*, 2126.
8. Mundy, C. J.; Siepmann, J. I.; Klein, M. L. *J. Chem. Phys.* **1995**, *102*, 3376.
9. Cui, S. T.; Cummings, P. T.; Cochran, H. D. *J. Chem. Phys.* **1996**, *104*, 255.
10. Cui, S. T.; Gupta, S. A.; Cummings, P. T.; Cochran, H. D. *J. Chem. Phys.* **1996**, *105*, 1214.
11. Jorgensen, W. L.; Madura, J. D.; Swenson, C. J. *J. Am. Chem. Soc.* **1984**, *106*, 6638.
12. Gear, C. W. *Numerical Initial Value Problems in Ordinary Differential Equation*; Prentice-Hall: Englewood Cliffs, 1971.
13. Andersen, H. J. *Comput. Phys.* **1984**, *52*, 24.
14. Lee, S. H. *Bull. Kor. Chem. Soc.* **2007**, *28*, 1371.
15. Min, S. H.; Son, C. M.; Lee, S. H. *Bull. Kor. Chem. Soc.* **2007**, *28*, 1689.
16. Fleisher, G. *Polym. Bull. (Berlin)* **1983**, *9*, 152.
17. Pearson, D. S.; Ver Strate, G.; von Meerwall, E.; Schilling, F. C. *Macromolecules* **1987**, *20*, 1133.
18. Von Meerwall, E.; Beckman, S.; Jang, J.; Mattice, W. L. *J. Chem. Phys.* **1998**, *108*, 4299.
19. Lee, S. H.; Chang, T. *Bull. Kor. Chem. Soc.* **2003**, *24*, 1590.
20. Lee, S. H.; Cummings, P. T. *Mol. Sim.* **1996**, *16*, 229.
21. Mendelson, R. A.; Bowles, W. A.; Finer, F. L. *J. Polym. Sci., Part A-2*, **1970**, *8*, 105.
22. Raju, V. R.; Smith, G. G.; Marin, G.; Knox, J. R.; Graessley, W. W. *J. Polym. Sci., Polym. Phys. Ed.* **1979**, *17*, 1183.