

Rat Liver β -Glucuronidase; Its Purification and Inhibition Studies

Han-Seung Jeong and Chul-Hak Yang†

Department of Chemistry, College of Natural Sciences, Seoul National University (Received July 30, 1985)

β -Glucuronidase (EC 3.2.1.31) which hydrolyzes D-glucuronate from β -D-glucuronide was purified from rat liver, using ammonium sulfate fractionation, DEAE-cellulose chromatography, Concanavalin-A Sepharose 4B chromatography and gel filtration on Sephadex G-200. This enzyme has the molecular weight of 280,000 daltons by gel filtration and 75,000 daltons by SDS-polyacrylamide gel electrophoresis. As its function is reverse of detoxification in the liver, the inhibition of the enzyme was tested with extracts of several food products and medicinal herbs, some are known as anti-cancer agents. Among them, *Panax ginseng* and *Cortnellus shiiae* inhibited the enzyme competitively and the K_i values were 9.22×10^{-2} and 0.102 mg/ml, respectively. These inhibitors strongly bound to DEAE-cellulose. The negatively charged amino acids, L-aspartate and L-glutamate, inhibited the enzyme, and K_i value of L-aspartate was 0.80 mM. The interaction between β -glucuronidase and p-nitrophenyl- β -D-glucuronide was found to involve ionic forces by the effect of ionic strength on the kinetic constant, V_{max}/K_m . It was inferred from these findings that cationic group at the active center of the enzyme is probably involved in attacking the substrate.

Introduction

β -Glucuronidase (EC 3.2.1.31) is a hydrolytic enzyme which catalyzes β -D-glucuronide + H₂O \longrightarrow Alcohol + D-glucuronate. It is distributed in animal tissues, plants and bacteria, and has been purified and characterized from human liver,¹ bovine liver,² mouse kidney,³ ox spleen,⁴ rabbit,⁵ rat liver,⁶⁻⁸ and rat uterus.⁹

It was reported that β -glucuronidase is complexed into different subcellular structures, the lysosomes and microsomes,¹⁰ which is derived from the same structural gene, *Gus*, on chromosome 5 and the two forms arise by post-translational processing.¹¹ Though they are heterogeneous after isoelectric focusing over the range PI 5.4-6.0,⁷ the two forms differ only a little in both carbohydrate and amino acid compositions.⁸

But there is also increasing interest on the role of glucuronidase in the destruction of host material at the invading edge of many tumors.^{12,13} Benzo(a)pyrene is a carcinogenic hydrocarbon commonly found in the environment and its metabolic formation of a diol epoxide is highly mutagenic¹⁴⁻¹⁶ and binds very actively to nucleic acids.^{17,18} The water soluble conjugates of benzo(a)pyrene, such as the glucuronides, have been generally viewed as detoxification products.¹⁹ The benzo(a)pyrene glucuronides are not entirely detoxification product but converted by glucuronidase to carcinogens at the sites distal to their formation.²⁰ (Kinoshita & Gelboin, 1978).

In this study, we found that *Panax ginseng* and *Cortnellus shiiae* contained some inhibitors of β -glucuronidase. Previously Saccharo-1,4-lactone was known to be powerful competitive inhibitor of β -glucuronidase and inhibitor was isolated from porcine sublingual gland (Sakamoto, 1973).²¹

Materials and Methods

Materials. Rats (300-500g) were obtained from animal breeding center of Seoul National University. p-Nitrophenyl- β -D-Glucuronide, α -methyl-D-mannoside, L-arginine and all marker proteins for molecular weight determination were purchased from Sigma Chemical CO. DEAE-Cellulose was from Whatman Ltd., Sephadex G-200 and Concanavalin-A

Sepharose 4B were obtained from Pharmacia Fine Chemicals. *Panax ginseng* was obtained from the Office of Monopoly and other herbs were purchased at the market. L-Cysteine and L-glutamate were purchased from Merck and other amino acids from Eastman Kodak Co.

All chemicals were of the highest purity available.

Enzyme Assay. β -Glucuronidase assays were performed by the method of Harris *et al.*²² with some modification. The assay mixture consisted of 0.5 ml of 0.2 mM p-nitrophenyl- β -D-glucuronide in 0.2 M sodium acetate buffer, pH 4.5, and 0.1 ml of enzyme solution. The mixture was incubated for 1 hr at 37°C. The reaction was terminated by the addition of 0.5 ml of 2.0 N sodium hydroxide solution. The color developed was measured by reading the absorbance at 400 nm. The amount of p-nitrophenol formed was determined from standard curve. One unit of the enzyme activity was defined as the amount of enzyme that release 1 μ mole of p-nitrophenol from p-nitrophenyl- β -D-glucuronide per hour under the condition specified above.

The protein was determined by the method of Lowry *et al.*²³ with crystallized bovine serum albumin as standard.

Enzyme Purification

Step I, Preparation of Crude Extract. About 10g of rat liver was minced with razor and homogenized with pestle in 2 volume (w/v) of 50 mM Tris-HCl buffer, pH 7.2, for 10 min. The pellet was resuspended in 2 volumes of the buffer, and homogenization and centrifugation were repeated. The supernatants were pooled.

Step II, (NH₄)₂SO₄ Fractionation. The crude extract was fractionated by adding solid ammonium sulphate to 70% saturation over a period of 20 min with constant stirring with magnetic stirrer. After 20 min setting the mixture was centrifuged at 30,000 g for 10 min and the pellet was dissolved in 30 ml of 50 mM Tris-HCl buffer, pH 7.2, and dialyzed against 2 l of the same buffer with three changes.

Step III, DEAE-Cellulose Chromatography. The dialyzed sample was loaded on DEAE-Cellulose column (3 \times 10 cm)

pre-equilibrated with 50 mM Tris-HCl buffer, pH 7.2. The column was washed with 150 ml of the same buffer and eluted with 500 ml of the same buffer and eluted with 500 ml of a linear gradient of 0-0.5 M NaCl in the buffer at the rate of 30 ml/hr, and 3 ml fractions were collected. The enzyme active fractions were pooled and concentrated to 10 ml with Dioflo PM 10 filter.

Step IV, Concanavalin-A Chromatography. The concentrated sample was applied to a column of Concanavalin-A Sepharose 4B (1.5 \times 7 cm) pre-equilibrated with 50 mM CaCl₂, 0.4 M NaCl. The column was washed with 100 ml of the above buffer and the enzyme activities were eluted with 100 ml of 0-0.6 M linear gradient α -methyl-D-mannoside in 50 mM Tris-HCl buffer, pH 7.2, at the rate of 5 ml/hr. β -Glucuronidase active fractions were pooled and concentrated to 5 ml by Diaflo PM 10 filter.

Step V, Sephadex G-200 Chromatography. The pooled sample from the previous step was applied to a column of Sephadex G-200 (2.6 \times 30 cm) pre-equilibrated with 50 mM Tris-HCl buffer, pH 7.2. The column was eluted with this buffer at the rate of 2.5 ml/hr and the void volume was measured by loading Blue Dextran 2000 as marker. β -Glucuronidase active fractions were pooled and used to inhibition studies.

Inhibition Studies. *Panax ginseng* extracts was diluted with 10 fold (w/v) with 50 mM Tris-HCl buffer, pH 7.2. *Cornellus shiike* was homogenated with pestle in the same buffer, after boiling for 5 min, centrifuged 4,000 g \times 5 min (Kokusan) and the supernatant was used. The inhibitor was mixed with 0.9 ml of 0.2 M sodium acetate buffer, pH 4.5 and 0.1 ml of purified enzyme. After the pre-incubation at 37°C for 10 min, 0.5 ml of 0.2 mM of p-nitrophenyl- β -D-glucuronide in the same buffer was added to the reaction tube. Enzyme reaction was carried out for 1 hr until 0.5 ml of 2.0 N NaOH was added as terminator. The color developed was detected at 400 nm. For determination of K_i values, the concentrations of substrate 4.0, 2.0, 1.0 and 0.5 mM were used and the incubation time was 5 min.

Effect of Amino Acid. The enzyme was pre-incubated with 1.0 and 10 mM of each amino acid in standard assay mixture without the substrate for 10 min at 37°C. Then added 0.2 mM substrate, the reaction mixture was incubated for 1 hr. For L-aspartate, the K_i value was calculated as in the case of inhibition studies.

Effect of Salt. Enzyme reactions were carried out in 10, 50 and 100 mM of NaCl, KCl, CaCl₂ and MgSO₄ to check the ionic

strength effect on enzyme activity. To test the suppression by salt, 80 μ g/ml of *ginseng* extract was added to the assay mixture. After incubation for 1 hr at 37°C, the activities were compared to the control which didn't contain any additional salt but inhibitor.

Results and Discussion

Purification of β -Glucuronidase. Overall purification was 430 fold with 36% recovery of the enzyme. The entire purification scheme is summarized in Table 1.

DEAE-cellulose column provided 10.3 fold increase in β -glucuronidase activity and showed double peaks at about 0.1 M NaCl with a little tailing. When the DEAE-Sephacyl was used, there were double peaks and tailing, too, but a little lower efficiency. The profile of DEAE-cellulose chromatography is shown in Figure 1.

The double peaks of the DEAE-cellulose chromatography means there are a few different types of β -glucuronidase. Plapp & Cole²⁴ showed multiple peaks of the DEAE-cellulose chromatography and described that the pattern was due to the difference in amount of some material bound to the protein. Now it is known that the six intracellular forms of the enzyme is L forms in the lysosomal fraction, a X form and four M forms in microsome, and the M forms are complexes containing 1, 2, 3, or 4 molecules of egasyn for each β -glucuronidase X tetramer.^{11, 25}




Figure 1. DEAE-Cellulose Chromatography. The column was eluted with 50 mM Tris-HCl buffer, pH 7.2 and developed with linear gradient of 0-0.5 M NaCl. Its dimension was 3 \times 10 cm and 3 ml fractions were collected. ●—●; protein. ○—○; β -glucuronidase.

TABLE 1: Purification of Rat Liver β -glucuronidase. A unit of Enzyme Activity was Defined as the Amount that Release 1 μ mole of p-nitrophenol per hour at 37°C under the Assay Condition Described in the text

Procedure	Total Protein (mg)	Total Activity (units)	Specific Activity (unit/mg protein)	Yield (%)	Purification Ratio
Homogenate	4020	76.8	0.0191	100	1
30,000 g Supernatant	2370	70.7	0.0298	92	1.56
70% (NH ₄) ₂ SO ₄ Fractionation	1680	53.2	0.0317	69	1.66
DEAE-Cellulose	141	46.2	0.328	60	17.2
Concanavalin-A Sepharose 4B	21.4	37.5	1.75	49	109
Sephadex G-200	3.30	27.5	8.33	36	436

With Concanavalin-A Sepharose affinity chromatography, the enzyme activity was eluted as a single peak at about 0.3 M α -methyl-D-mannoside. It provided 6.3 fold increase in activity and the pattern of the column is shown in Figure 2.

Gel filtration on Sephadex G-200 provided 4.0 fold increase in activity. The profile of this step is shown in Figure 3. After the gel filtration on Sephadex G-200, the enzyme showed one band on polyacrylamide gel, so we could set about the inhibition studies.

Inhibition Studies. Among the several food products and medicinal herbs, known as anti-cancer agent, *Panax ginseng* and *Cortnellus shiiae* showed more inhibition which are shown in Figure 4 and 5. The K_m value of β -glucuronidase was 0.735 mM.

In the case of inhibition by *Panax ginseng*, as the protein concentration increased 32, 80, 160 $\mu\text{g/ml}$, so the K_m values 1.07, 1.50 and 2.10 mM, respectively. Each lines of Lineweaver-Burk plot had the common V_{max} value. So we concluded that the inhibition by *Panax ginseng* is competitive.

The inhibition by *Cortnellus shiiae* showed the same pattern. When the protein concentrations were 38, 75 and 190

$\mu\text{g/ml}$, the K_m values were 1.25, 1.44 and 2.10 mM, respectively. Each lines of Lineweaver-Burk plot had the common V_{max} value, too, so the inhibition by *Cortnellus shiiae* is deduced to be competitive.

Since the inhibitors bind to the catalytic site of the enzyme or to other potential sites for a reaction to take place, in either case a dead-end complex (EI) would be formed.

The K_i values of *P. ginseng* and *C. shiiae* extracts were 9.22×10^{-2} and 0.102 mg/ml, respectively (Figure 6). But this result doesn't show that the former is more effective inhibitor than the latter. As we didn't use the purified materials, merely using the extracts of putative inhibitors, there might be any possibility that the inhibitor proteins from the two sources are the same and there is more quantity in *P. ginseng* components than in *C. shiiae*.

When the extract of *P. ginseng* was loaded on DEAE-cellulose column, the inhibitors were strongly bound to DEAE-cellulose, indicating that they have strong anionic charges. So it is anticipated that the inhibitions are resulting from interaction between the anionic inhibitors and cationic group at the active center of β -glucuronidase. That this cationic group of




Figure 2. Concanavalin-A Sepharose 4B Affinity Chromatography. The column was eluted with 50 mM Tris-HCl buffer, pH 7.2 and developed with linear gradient of 0–0.6 M of α -methyl-D-mannoside in 50 mM Tris-HCl buffer, pH 7.2 containing 0.4 M NaCl. The column size was 1.5 \times 7.0 cm. ●—●; protein. ○—○; β -glucuronidase.




Figure 3. Gel Filtration on Sephadex G-200. The column was eluted with 50 mM Tris-HCl buffer, pH 7.2. The dimension of the column was 2.6 \times 30 cm. The void volume was measured by loading Blue Dextran 2000 as marker previous to apply the sample. ●—●; protein. ○—○; β -glucuronidase.




Figure 4. Lineweaver-Burk plot of β -glucuronidase in the presence of *Panax ginseng*. Enzyme activity was measured as described in the text and inhibitor concentrations and K_m values of each cases are as follows. ●; control $K_m = 0.735$ mM, □; 32 $\mu\text{g/ml}$ $K_m = 1.07$ mM, ○; 80 $\mu\text{g/ml}$ $K_m = 1.50$ mM, ■; 160 $\mu\text{g/ml}$ $K_m = 2.10$ mM.




Figure 5. Lineweaver-Burk plot of the enzyme in the presence of *Cortnellus shiiae* extract. Enzyme activities were measured as described in the text. Inhibitor concentrations and K_m values of each line are as follows. ●; control $K_m = 0.735$ mM, □; 38 $\mu\text{g/ml}$ $K_m = 1.25$ mM, ○; 75 $\mu\text{g/ml}$ $K_m = 1.44$ mM, ■; 190 $\mu\text{g/ml}$ $K_m = 2.10$ mM.

the β -glucuronidase probably belong to an imidazolium group had been reported by Wang (1972). Table 2 shows the effect of ionic strength in the reaction mixture. The double reciprocal plot of enzyme activities in different NaCl concentrations is shown in Figure 7 and the K_m and V_{max} values are in Table 3. When the inhibition reactions were tested in 10–30 mM of salt solution, such as $CaCl_2$, NaCl, KCl and $MgSO_4$, there were not such drastic suppression as Wataru Sakamoto had reported (1973). Instead, the ionic strength affects to enzyme activity as some mixed inhibition. The equation relating the rate of an enzyme reaction at low substrate concentration to the ionic strength of the medium on the basis of the primary kinetic effect is

$$\log(k_2/K_m) = \log(k_2/K_m)_0 + 1.02 z_A z_B I$$

where Z_A and Z_B are the charges on the active site and on the




Figure 6. Secondary Plots for Inhibitions. Upper; Inhibition by *Panax ginseng* extract $K_i = 9.22 \times 10^{-2}$ mg/ml Lower; Inhibition by *Cortnellus shiike* extract $K_i = 0.102$ mg/ml.

TABLE 2: Effect on Enzyme Activity and Suppression by the Ionic Strength. Suppression tests were carried out in 80 μ g/ml of *P. ginseng* extract. Activities are in % Compared to Control not Containing any Additional Salt, Respectively

Salt	Without Inhibitor	With Inhibitor
NaCl	10 mM	98
	50 mM	89
	100 mM	83
KCl	10 mM	97
	50 mM	91
	100 mM	83
$CaCl_2$	10 mM	97
	50 mM	84
	100 mM	38
$MgSO_4$	10 mM	96
	50 mM	79
	100 mM	30

interacting substrate, respectively, I is the ionic strength, and $(k_2/K_m)_0$ is the second order rate constant at zero ionic strength (Mintel & Westley, 1966). A plot of $\log(k_2/K_m)$ versus the square root of ionic strength should yield a straight line with a slope approximately equal to $Z_A Z_B$. We have used V_{max} in place of k_2 due to the uncertainty of the enzyme concentration used. From experiments employing sodium chloride and substrate concentrations, $\log(V_{max}/K_m)$ for β -glucuronidase has been plotted against I , giving a straight line with a slope of -1.7 (Figure 8). The magnitude of this value must be regarded as approximate, since the law of Debye-Huckel is strictly valid only at low salt concentrations. The direction of the effects, however, indicates that charge neutralization occurs when glucuronides react with the enzyme.

Another evidence that the active site of β -glucuronidase is charged is the result of amino acid effects. We tested with nine amino acids and found that *L*-aspartate and *L*-glutamate in-




Figure 7. Ionic Strength Effect on K_m and V_{max} . From 50 mM sodium acetate buffer, pH 4.5, the ionic strength was gradually increased by adding NaCl to 0.1, 0.2 and 0.3 M, respectively. In each ionic strength, the enzyme activities were measured with the variation of substrate concentrations; 0.5, 1.0, 2.0 and 4.0 mM. ●; No NaCl ($I = 0.05$), □; 0.1 M NaCl ($I = 0.15$), ○; 0.2 M NaCl ($I = 0.25$), ■; 0.3 M NaCl ($I = 0.35$). K_m and V_{max} values are in Table 3.

TABLE 3: K_m and V_{max} Values from NaCl Inhibition Reactions. All the Parameters Presented were Obtained from Figure 10

Concentration (M)	I	K_m (mM)	$10 \times V_{max}$ (OD ₄₀₀ /min)	$\sqrt{\log(10 \times V_{max}/K_m)}$
0	0.05	0.735	18.6	0.22
0.1	0.15	0.775	15.2	0.39
0.2	0.25	0.870	10.6	0.50
0.3	0.35	0.952	7.7	0.59




Figure 8. Ionic strength effect. The logarithm of $10 \times V_{\max}/K_m$ was plotted against the square root of ionic strength. This plot is based on Table 3.




Figure 9. Lineweaver-Burk plots on the enzyme in the presence of *L*-aspartate. Upper; Secondary plot for inhibition to find K_i value. The K_i is 0.80 mM. Lower: Enzyme activities were measured as the same method of the tests to find the K_m values. ●; control ▲; 1.0 mM *L*-aspartate. ○; 2.0 mM ■; 5.0 mM. K_m and V_{\max} values are in Table 3.

hibited the enzyme activity (Table 4). When the reaction was performed with 10 mM of *L*-aspartate at 37°C for 1 hr, the remained activity was only 7%. The inhibition pattern of *L*-

TABLE 4: Amino Acid Effect on the Enzyme Activity.

Measurements were Carried in the Standard Assay Mixture for 1 hr. Activities are in % Compared to no Effector Present

Amino acid	Concentration (mM)	
	1.0	10.0
<i>L</i> -Arginine	100	100
<i>L</i> -Aspartate	38	7
<i>L</i> -Cysteine	97	102
<i>L</i> -Glutamate	93	78
<i>L</i> -Histidine	95	98
<i>L</i> -Leucine	98	102
<i>L</i> -Proline	100	101
<i>L</i> -Threonine	99	100
<i>L</i> -Tyrosine	101	104

aspartate was competitive and the K_i value was 0.80 mM (Figure 9). This result means the active site has positive charge and binding seems not exclusively dependent on the charge. *L*-glutamate, larger one carbon than *L*-aspartate, didn't show such drastic inhibition, which means that steric effect is another important effect in binding substrates and active site.

In this paper, we showed that *Panax ginseng* and *Cortnellus shiike* contain a certain inhibitor of β -glucuronidase and *L*-aspartate is another inhibitor. All the inhibition patterns were competitive and the K_i values were 9.22×10^{-2} , 0.102 mg/ml and 0.80 mM, respectively. There was little suppression by salts and the effect of ionic strength was verified. All the inhibitions were seemed to be related to the charges on the active site and the interacting substrates.

References

- (1) B. Hultberg & P.A. Ockerman, *Clin. Chim. Acta.* **39**, 49 (1972).
- (2) M. Himeno, Y. Hashigichi & K. Kato, *J. Biochem.* **76**, 1243 (1974).
- (3) C.-W. Lin, M.L. Orcutt, & W.H. Fishman, *J. Biol. Chem.* **250**, 4737 (1975).
- (4) G.T. Mills, J. Paul & E.E.B. Smith, *Biochem. J.* **53**, 232 (1953); R. Mintel & J. Westley, *J. Biol. Chem.* **241**, 3386 (1966).
- (5) R.T. Dean, *Biochem. J.* **138**, 407 (1974).
- (6) P.D. Stahl & O. Touster, *J. Biol. Chem.* **246**, 5398 (1971).
- (7) J.W. Owens, K.L. Gammon & P.D. Stahl, *Archs Biochem. Biophys.* **166**, 258 (1975).
- (8) D.R.P. Tulsiani, H. Six & O. Touster, *Proc. Nat'l Acad. Sci. USA* **75**, 3080 (1978).
- (9) H.W. Kim & C.H. Yang, *Korean Biochem. J.* **16**, 111 (1983).
- (10) K. Paigen, *Exp. Cell Res.* **25**, 286 (1961).
- (11) R.T. Swank & K. Paigen, *J. Mol. Biol.* **77**, 371 (1973).
- (12) B. Sylven, *Europ. J. Cancer* **4**, 463 (1968).
- (13) C. Hoch-Ligetti, H.H. Grantham & J.M. Arvin, *Cancer* **24**, 297 (1969).
- (14) S.K. Yang, D.W. McCourt, P.R. Roller & H.V. Gelboin, *Proc. Nat'l Acad. Sci. USA* **73**, 2594 (1976).
- (15) C. Malaveille, H. Bartsch, P.L. Glover & P. Slims, *Biochem. Biophys. Res. Commun.* **66**, 693 (1975).
- (16) H.V. Gelboin, *Cancer Res.* **29**, 1272 (1969). J.H. Glaser, K.J. Roozen, F.E. Brot & W.S. Sly, *Archs Biochem. Biophys.* **166**, 536 (1975).

- (17) H.W.S. King, M.H. Thompson & P. Brookes, *Cancer Res.* **35**, 1263 (1975).
- (18) P. Daudle, M. Daquesne, P. Vigny, P.L. Glover & P. Slims, *FEBS Lett.* **57**, 250 (1975).
- (19) K.H. Harper, *Br. J. Cancer* **12**, 121 (1958).
- (20) N. Kinoshita & H.V. Gelboin, *Science* **199**, 307 (1978).
- (21) W. Sakamoto, O. Nishikaze & E. Sakakibara, *Biochim. Biophys. Acta.* **329**, 72 (1973).
- (22) R.G. Harris, J.J.M. Rowe, P.S. Stewart & D.C. Williams, *FEBS Lett.* **29**, 189 (1973).
- (23) O.H. Lowry, N.J. Rosebrough, A.L. Farr & R.J. Randall, *J. Biol. Chem.* **193**, 265 (1951).
- (24) B.V. Plapp & R.D. Cole, *Biochemistry* **6**, 3676 (1967).
- (25) S. Tomino & K. Paigen, *J. Biol. Chem.* **250**, 1146 (1975)