

## Communications

### Synthesis of Cinnamyl Amines from the Baylis-Hillman Adducts: Transformation of Allylic Alcohols to Allylic Amines


Hong Jung Lee, Yun Mi Chung, Ka Young Lee, and Jae Nyong Kim\*

Department of Chemistry, Chonnam National University, Kwangju 500-757, Korea


Received June 1, 2000

The Baylis-Hillman reaction is one of the most powerful carbon-carbon bond-forming methods in organic synthesis.<sup>1</sup> The Baylis-Hillman adducts, which are allylic alcohol derivatives, can be formed most often by the reaction of activated vinyls and carbonyl compounds.<sup>1</sup> Besides the usefulness of these Baylis-Hillman adducts themselves, further derivatization with various nucleophilic reagents toward synthetically useful compounds has been studied in depth by us and other groups.<sup>2</sup>

Recently, we have reported on the reaction of the Baylis-Hillman adducts of *N*-tosylimines and *N,N*-dimethylformamide dimethyl acetal (DMF-DMA).<sup>3</sup> We could obtain the *N*-methyl-*N*-tosylallylic amine derivatives stereoselectively as shown in Scheme 1.<sup>3</sup> As a continuous work, we tried on the reaction of the Baylis-Hillman adducts of benzaldehydes **1a-f**



Scheme 1



Scheme 2



**f** and *N,N*-dimethylformamide dimethyl acetal. From the reaction we could obtain *N,N*-dimethylcinnamyl amine derivatives **2a-f** in moderate to good yields stereoselectively (Scheme 2). *N,N*-Dialkylcinnamyl amine moiety constitutes an important scaffold in many biologically active compounds.<sup>4</sup>

As shown in Scheme 2 and in Table 1, the reaction of the Baylis-Hillman adducts **1a-f** and DMF-DMA in DMF afforded the corresponding amines **2a-f** in 63–88% isolated yields in a highly stereoselective manner. Ester substituted derivatives **1a-c** gave the *E* isomer **2a-c** selectively, while for nitrile substituted derivatives **1d-f** the *Z* isomer **2d-f** pre-

Table 1. Synthesis of *N,N*-dimethylcinnamyl amine derivatives

Entry	B-H adducts	Products (% yields)
1		<b>2a</b> ( <i>E</i> , 73) <sup>a</sup>
2		<b>2b</b> ( <i>E</i> , 69) <sup>b</sup>
3		<b>2c</b> ( <i>E</i> , 72) <sup>a</sup>
4		<b>2d</b> ( <i>Z</i> , 83) <sup>c</sup>
5		<b>2e</b> ( <i>Z</i> , 88) <sup>c</sup>
6		<b>2f</b> ( <i>Z</i> , 63) <sup>c</sup>

<sup>a</sup>*Z* isomer was observed in trace amounts on tlc. <sup>b</sup>*Z* isomer was isolated in 20% yield. <sup>c</sup>*E* isomer was observed in trace amounts on tlc.



dominates. The reaction did not proceed at room temperature. Xylene can replace DMF as solvent (76% of **2a** was isolated in the case of **1a**). Without DMF-DMA no reaction was observed. The reaction of **1a** and dimethylamine hydrochloride (3.0 equiv) in DMF (reflux, 48 h) gave **2a** in 43% yield. However, the *E/Z* ratio was about 2 : 1 in this case. The reaction of **1a** and *N,N*-dimethylacetamide dimethyl acetal (2 equiv) in *N,N*-dimethylacetamide gave the expected  $\gamma,\delta$ -unsaturated amide derivative (**3**, 2-(2-dimethylcarbamoyl ethyl)-3-phenyl acrylic acid ethyl ester) via the normal Claisen rearrangement pathway in 69% isolated yield.<sup>5,6</sup>

The stereochemistry of the products (*E* for ester, *Z* for nitrile) can be explained as we have already proposed.<sup>2b,3</sup> Some representative spectroscopic data of selected products **2a**, **2d**, and **3** were presented.<sup>6</sup>

We tentatively propose the reaction mechanism as follows: (1) exchange of methoxy group in DMF-DMA with the Baylis-Hillman adduct **1a-f** to form the intermediate **I**, (2) concerted Michael type addition of dimethylamino group<sup>2b</sup> and concomitant elimination of methyl formate gave **2a-f** (Scheme 3).

In a brilliant works of Buchi and co-workers,<sup>7</sup> treatment of some allylic alcohols with DMF-DMA gave the homologous  $\beta,\gamma$ -unsaturated amides via the formation of carbene and subsequent [2,3]-sigmatropic rearrangement (Scheme 3). However, in our reaction conditions, we could not observe the corresponding amide products **4** at all. This discrepancy could be explained as follows. In the cases of simple allylic alcohols the carbene mechanism works.<sup>7</sup> However, for the good Michael acceptor such as **1a-f**, Michael type addition of dimethylamino group is the preferred pathway leading to **2a-f**. To confirm the hypothesis we examined the reaction of ethyl 2-hydroxymethyl-3-phenyl-2-propenoate (**5**)<sup>2b</sup> and DMF-DMA in the same reaction conditions (Scheme 4). In this case the  $\beta$ -position is not a good Michael acceptor, and as a result, **2a** was isolated in low yield (10%) with 90% of the recovered starting material.

**Acknowledgment.** We wish to thank the Ministry of Korea (BK-21 project) for financial support of this work.

## References and Notes

- (a) Brezezinski, L. J.; Rafel, S.; Leahy, J. W. *J. Am. Chem. Soc.* **1997**, *119*, 4317. (b) Rafel, S.; Leahy, J. W. *J. Org. Chem.* **1997**, *62*, 1521. (c) Baylis, A. B.; Hillman, M. E. D. German Patent 2155113 (*Chem. Abstr.* **1972**, *77*, 34174q). (d) Drewes, S. E.; Roos, G. H. P. *Tetrahedron* **1988**, *44*, 4653. (e) Basavaiah, D.; Rao, P. D.; Hyma, R. S. *Tetrahedron* **1996**, *52*, 8001. (f) Basavaiah, D.; Bharathi, T. K.; Gowriswari, V. V. L. *Tetrahedron Lett.* **1987**, *28*, 4351. (g) Ciganek, E. *Organic Reactions*; John Wiley & Sons: New York, 1997; Vol. 51, pp 201-350. (h) Iwabuchi, Y.; Nakatani, M.; Yokoyama, N.; Hatakeyama, S. *J. Am. Chem. Soc.* **1999**, *121*, 10219.
- (a) Lee, H. J.; Seong, M. R.; Kim, J. N. *Tetrahedron Lett.* **1998**, *39*, 6223. (b) Kim, H. S.; Kim, T. Y.; Lee, K. Y.; Chung, Y. M.; Lee, H. J.; Kim, J. N. *Tetrahedron Lett.* **2000**, *41*, 2613. (c) Kim, J. N.; Lee, K. Y.; Kim, H. S.; Kim, T. Y. *Org. Lett.* **2000**, *2*, 343. (d) Basavaiah, D.; Sarma, P. K. S. *J. Chem. Soc., Chem. Commun.* **1992**, 955. (e) Charette, A. B.; Cote, B.; Monroc, S.; Prescott, S. J. *Org. Chem.* **1995**, *60*, 6888. (f) Basavaiah, D.; Krishnamacharyulu, M.; Hyma, R. S.; Pandiaraju, S. *Tetrahedron Lett.* **1997**, *38*, 2141. (g) Chavan, S. P.; Ethiraj, K. S.; Kamat, S. K. *Tetrahedron Lett.* **1997**, *38*, 7415. (h) Perlmutter, P.; Tabone, M. *Tetrahedron Lett.* **1988**, *29*, 949. (i) Lawrence, R. M.; Perlmutter, P. *Chem. Lett.* **1992**, 305. (j) Foucaud, A.; El Guemmout, F. *Bull. Soc. Chim. Fr.* **1989**, 403.
- Lee, H. J.; Kim, H. S.; Kim, J. N. *Tetrahedron Lett.* **1999**, *40*, 4363.
- (a) Weingartner, H.; Rudorfer, M. V.; Buchsbaum, M. S.; Linnoila, M. *Science* **1983**, *221*, 472. (b) Loibner, H.; Pruckner, A.; Stutz, A. *Tetrahedron Lett.* **1984**, *25*, 2535. (c) Kundu, M. K.; Bhat, S. V. *Synth. Commun.* **1999**, *29*, 93. (d) Eagen, M. C.; Cromwell, N. H.; *J. Org. Chem.* **1974**, *39*, 3863. (e) Hbaieb, S.; Latiri, Z.; Amri, H. *Synth. Commun.* **1999**, *29*, 981, and references cited therein.
- Basavaiah, D.; Pandiaraju, S.; Krishnamacharyulu, M. *Synlett* **1996**, 747.
- Some representative spectroscopic data of products are as follows:  
**2a**: clear oil; <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.36 (t, *J* = 7.2 Hz, 3H), 2.24 (s, 6H), 3.30 (s, 2H), 4.29 (q, *J* = 7.2 Hz, 2H), 7.34-7.62 (m, 5H), 7.83 (s, 1H); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  14.30, 45.17, 54.37, 60.93, 128.39, 128.78, 130.32, 130.61, 135.40, 142.62, 168.58; Mass (70 eV) *m/z* (rel intensity) 58 (100), 91 (24), 115 (40), 131 (17), 172 (19), 204 (19), 218 (36), 233 (M<sup>+</sup>, 21).  
**2d**: clear oil; <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  2.32 (s, 6H), 3.21 (s, 2H), 7.08 (s, 1H), 7.40-7.80 (m, 5H); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  44.95, 63.84, 108.97, 118.64, 128.84, 128.90, 130.37, 133.24, 145.08.  
**3**: clear oil; <sup>1</sup>H NMR (CDCl<sub>3</sub>)  $\delta$  1.36 (t, *J* = 7.2 Hz, 3H), 2.53-2.60 (m, 2H), 2.85-2.91 (m, 2H), 2.94 (s, 3H), 3.01 (s, 3H), 4.28 (q, *J* = 7.2 Hz, 2H), 7.30-7.42 (m, 5H), 7.73 (s, 1H); <sup>13</sup>C NMR (CDCl<sub>3</sub>)  $\delta$  14.24, 23.49, 32.76, 35.32, 37.15, 60.82, 128.51, 128.57, 129.20, 131.80, 135.22, 139.93, 168.05, 172.11.
- (a) Buchi, G.; Cushman, M.; Wuest, H. *J. Am. Chem. Soc.* **1974**, *96*, 5563. (b) Chan, K.-K.; Saucy, G. *J. Org. Chem.* **1977**, *42*, 3828.