



chloride. In the previous paper<sup>9</sup>



(R, R', and R'' = H or alkyl; OR'' = alkyl, amine or halide group)




we reported the synthesis and structure of functionally substituted monoalkyltin trichloride,  $\text{CH}_3\text{OOCCH}_2\text{CH}(\text{COOCH}_3)\text{CH}_2\text{SnCl}_3$  from the reaction of dimethyl itaconate ( $\text{CH}_3\text{OOCCH}_2\text{C}(\text{COOCH}_3)=\text{CH}_2$ ) with  $\text{SnCl}_2$  and HCl gas. Recently, further study on this compound system is under way. With view to expanding this chemistry, the present paper describes the result of the reaction of dimethyl itaconate with metallic tin and gaseous hydrogen chloride.

Sn (29.67 g, 250 mmol) and dimethyl itaconate (79.08 g, 500 mmol) in 200 ml of ethylene glycol dimethyl ether (or diethyl ether) were placed in a 500 ml three-necked flask equipped with a mechanical stirrer, a reflux condenser, and a gas inlet tube. Through the stirred suspension was bubbled for 5 hours anhydrous hydrogen chloride (27.37 g, 750 mmol) at room temperature. The solvent was removed, and the residue was washed with cold ethyl ether to produce a white crystalline solid in 92% (based on tin consumed) yield. The solid product was recrystallized from dichloromethane-petroleum ether solvent mixture (1:1) resulting in colorless crystals (mp. 154-155°C (Ref.<sup>9</sup> 154-155°C)). Anal. Found: C, 22.00; H, 2.84. Calcd. for  $\text{C}_7\text{H}_{11}\text{O}_4\text{Cl}_3\text{Sn}$ : C, 21.88; H, 2.89.

As is seen in the above experimental result, the reaction of dimethyl itaconate with metallic tin in the presence of gaseous hydrogen chloride produces the monoalkyltin trichloride,  $\text{CH}_3\text{OOCCH}_2\text{CH}(\text{COOCH}_3)\text{CH}_2\text{SnCl}_3$  instead of the dialkyltin dichloride. The reaction seems to proceed in the following general Scheme 1.

At the beginning stage of the above reaction, formation of a large amount of stannous chloride could easily be identified, but the quantity of the stannous chloride was diminished as the reaction proceeded. The reaction of  $\alpha$ ,  $\beta$ -unsaturated carbonyl compounds with  $[\text{HSnCl}_3]$  has been well established.<sup>2</sup> We also carried out the equimolar reaction of metallic tin with dimethyl itaconate. However, only monoalkyltin trichloride was formed irrespective of the mole ratio. Thus the formation of monoalkyltin trichloride in the reaction system is intrinsic and is not the result of a particular synthetic strategy. Why the reaction affords only monoalkyltin trichloride, rather than dialkyltin dichloride is not clear. There is a temptation to invoke the steric hindrance of dimethyl itaconate.

In conclusion, unlike the precedented results, the reaction of dimethyl itaconate either with stannous chloride or with metallic tin in the presence of hydrogen chloride affords the same product: the reaction is the first example that only



Scheme 1.

monoalkyltin trichloride is specifically prepared irrespective of the synthetic route.

**Acknowledgement.** We gratefully acknowledge the support of this work by the Ministry of Science and Technology.

## References

1. R. E. Hutton and V. Oakes, *Organotin Compounds: New Chemistry and Applications*, ACS 157, Am. Chem. Soc., Washington DC, p. 123 (1976).
2. R. E. Hutton, J. W. Burley, and V. Oakes, *J. Organomet. Chem.*, **156**, 369 (1978).
3. P. G. Harrison, T. J. King, and M. A. Healy, *J. Organomet. Chem.*, **182**, 17 (1979).
4. R. A. Howie, E. S. Paterson, J. L. Wardell, and J. W. Burley, *J. Organomet. Chem.*, **259**, 71 (1983).
5. R. A. Howie, E. S. Paterson, J. L. Wardell, and J. W. Burley, *J. Organomet. Chem.*, **304**, 301 (1986).
6. B. K. Deb and A. K. Ghosh, *Polyhedron*, **5**, 863 (1986).
7. B. K. Deb and A. K. Ghosh, *Can. J. Chem.*, **65**, 1241 (1987).
8. S. W. Ng, C. Wei, V. G. Kumar Das, J. P. Charland, and F. E. Smith, *J. Organomet. Chem.*, **364**, 343 (1989).
9. O. S. Jung, J. H. Jeong, and Y. S. Sohn, *J. Organomet. Chem.*, **397**, 17 (1990).
10.  $\text{HSnX}_3$  species (X = Cl, Br) have been isolated as complexes with  $\text{Et}_2\text{O}$  or amines. The trichlorostannane has been prepared by reaction of  $\text{SnX}_4$  with  $\text{LiAlH}_4$  and also by addition of  $\text{SnX}_2$  to a solution of excess HX in  $\text{Et}_2\text{O}$ : See ref. 2 and therein ref. 5 and 6.
11. One carbonyl group involved in the six-membered chelate is released from the tin atom in solution, so that the tin atom is pentacoordinate in solution in contrast to hexacoordinate in the solid state as pointed out in ref. 9.

## Heteroleptic Crown Thioether Complexes: Synthesis and Characterization of $[\text{M}(\text{CO})_3(\text{9S3})]$ ( $\text{M} = \text{Cr}, \text{Mo}, \text{W}$ ). Crystal Structure of $[\text{W}(\text{CO})_3(\text{9S3})]$ ( $\text{9S3} = 1,4,7\text{-Trithiacyclononane}$ )

Hyun-Joon Kim, Youngkyu Do\*, Hyo Won Lee<sup>†</sup>,  
Jong Hwa Jeong<sup>‡</sup>, and Youn Soo Sohn<sup>‡</sup>


Department of Chemistry, Korea Advanced Institute of  
Science and Technology, Seoul 130-650

<sup>†</sup>Department of Chemistry, Chungbuk National  
University, Cheongju 360-763

<sup>‡</sup>Inorganic Chemistry Laboratory, Korea Institute of  
Science and Technology, Seoul 136-971

Received February 21, 1991

A nine-membered crown thioether 1,4,7-trithiacyclononane (9S3), among others, has been employed in the development of crown thioether chemistry owing to its unique  $\sigma$ -donor



**Figure 1.** ORTEP plot of  $[\text{W}(\text{CO})_3(9\text{S}3)]$  (**3**), showing thermal ellipsoids at 50% probability level. All methylene hydrogen atoms are omitted for clarity. Selected average distances ( $\text{\AA}$ ) and angles (deg) with calculated standard deviation in parentheses: W-S, 2.509(7); W-C (carbonyl carbon), 1.957(7); carbonyl C-O, 1.161(7); S-W-S, 82.6(1); C-W-C (carbonyl carbon atoms), 89.9(16); S-W-C (*trans*), 172.7(11); two sets of S-W-C (*cis*): 90.3(12), 97.3(12).

and  $\pi$ -acid character as well as its strong chelating ability stemmed from the retention of endodentate conformation on binding to a trigonal face of a metal.<sup>1-3</sup> To date most exploratory research on the coordination chemistry of 9S3 has proceeded to determine its scope. The observation<sup>3</sup> of large set of homoleptic complexes suggests that 9S3 generally stabilizes a wide variety of lower oxidation states of metal ions and lacks sufficient flexibility to accommodate conformations other than a trigonal endodentate conformation although examples<sup>4,5</sup> became realized in the direction of extending the scope of crown thioether chemistry.

While homoleptic complexes are noble and useful, their robustness and the absence of vacant coordination sites may lead to their low reactivity.<sup>6</sup> On the other hand, heteroleptic 9S3 complexes, which constitute a newly emerging class,<sup>3,5-7</sup> contain heteroleptic ligands which are potentially susceptible for further reaction. As part of our efforts to exploit the chemistry of heteroleptic class of crown thioether, the group 6 metal carbonyl complexes of 9S3,  $[\text{M}(\text{CO})_3(9\text{S}3)]$  (M=Cr, Mo, W), were synthesized and characterized, brief accounts of which are reported herein.

An equimolar (0.3 mmol) reaction mixture of  $[\text{M}(\text{CO})_3(\text{CH}_3\text{CN})_3]$ <sup>8</sup> and 9S3<sup>9</sup> in 20 ml of  $\text{CH}_3\text{CN}$  was allowed to react anaerobically at ambient temperature, producing yellow precipitate. The yellow solids were collected, washed with  $\text{Et}_2\text{O}$  and recrystallized from DMF- $\text{Et}_2\text{O}$  pair, affording analytically pure crystalline  $[\text{M}(\text{CO})_3(9\text{S}3)]$  (**1**: M=Cr; **2**: M=Mo; **3**: M=W)<sup>10</sup> in the yields of 30, 40 and 32% for **1**, **2** and **3**, respectively. The composition and crystal structure of **3** were established by single-crystal X-ray analysis.<sup>13</sup> Elemental analysis<sup>10a</sup> of **1** and the comparison of spectroscopic data for **1**, **2** and **3**, *vide infra*, indicate that the compound of the lightest congener, **1**, has the same structural core as its heav-

**Table 1.** Spectroscopic Data for  $[\text{M}(\text{CO})_3(9\text{S}3)]$

	M	IR <sup>a</sup> $\nu_{\text{CO}}/\text{cm}^{-1}$	<sup>1</sup> H-NMR <sup>b</sup> ppm <sup>c</sup>	<sup>13</sup> C- <sup>1</sup> H-NMR <sup>b</sup> /ppm <sup>c</sup>	
				9S3	CO
<b>1</b>	Cr	1910,1790	2.53 (s)	31.9	232.1
<b>2</b>	Mo	1915,1796	2.55-2.69 (complex)	31.3	222.7
<b>3</b>	W <sup>d</sup>	1905,1788	2.55-2.68 (complex)	32.6	216.2

<sup>a</sup>KBr. <sup>b</sup> $\text{d}_6$ -DMSO. <sup>c</sup>Relative to TMS. <sup>d</sup>Coupling due to <sup>183</sup>W ( $I=1/2$ , 14.28%) has not been resolved in both NMR spectra.

**Table 2.** Comparison of IR Data for a series of  $[\text{LMo}(\text{CO})_3]$

L	$\nu_{\text{CO}}$ , $\text{cm}^{-1}$	Ref.
9S3	1915,1796	This work
9N3	1850,1740-1700	20
$\text{C}_6\text{H}_6$	1969,1936,1921	21
$\text{C}_5\text{H}_5^-$	1901,1796,1746	22
$\text{HBPz}_3^-$	1897,1761	23
$\text{C}_2\text{B}_9\text{H}_{11}^{2-}$	1909,1876,1741	24
12P3	1945,1840	25

ier analogues.

The molecular structure of **3**, displayed in Figure 1, reveals that 9S3 and three carbonyl groups are situated facially, approaching the overall symmetry of  $\text{C}_3$ . The  $\text{WS}_3\text{C}_3$  coordination sphere deviates from the regular octahedral geometry. Constriction of the S-W-S bond angles (average 82.6°), a measure of the extent of distortion in  $\text{WS}_3\text{C}_3$  coordination geometry, does not lead to corresponding dilation of carbonyl trigonal face as indicated by the average C-W-C (carbonyl carbon atoms) bond angle of 89.9°. The structure of **3** provides rare example of the W-SR<sub>2</sub> (R=sp<sup>3</sup>-C) bonding distance whose average value of 2.509(7)  $\text{\AA}$  in **3** is shorter by *ca* 0.06  $\text{\AA}$  than those in previously reported compounds with W-SR<sub>2</sub> interaction.<sup>14</sup> It is also interesting to note that the average W-S distance in **3** is shorter than the average Mo-S distance of 2.520(15)  $\text{\AA}$  in **2**,<sup>11</sup> the lighter congener analogue of **3**.<sup>15</sup> This observation is consistent with the expectation that a soft base 9S3 interacts more strongly with the softer acid tungsten.<sup>16</sup>

The structurally characterized  $\text{C}_3$  molecular symmetric nature of **2**<sup>11</sup> and **3** provides explanation for the vibrational spectral pattern ( $\text{A}_1$  and E) of **2** and **3** in the region of  $\nu_{\text{CO}}$ , which and comparison of IR spectra of **1**, **2** and **3**, in turn, suggest the facial  $\text{CrS}_3\text{C}_3$  coordination geometry of **1** in the solid state. The carbonyl stretching frequencies for **1**, **2** and **3** are listed in Table 1 along with other spectroscopic data. Upon dissolution in DMSO, no liberation of 9S3 is observed in all three compounds,<sup>17</sup> indicating the integrity of  $[\text{M}(\text{CO})_3(9\text{S}3)]$  in solution. The complex <sup>1</sup>H-NMR patterns of **2** and **3** are ascribed to AA' BB' or ABCD coupling of methylene protons. On the other hand, for **1**, a relatively sharp singlet enables to speculate fluxionality in **1** such as a merry-go-round type rotation of 9S3 with respect to Cr center although no example of 9S3 complex with fluxionality is available to date.<sup>18</sup>

As stated earlier in this report, the unique  $\sigma$ -base and  $\pi$ -acid character of 9S3 might be attributed to the stability of  $[\text{M}(\text{CO})_3(9\text{S3})]$  toward oxidative decarbonylation reaction.<sup>19</sup> A relative measure of  $\pi$ -acidity of L in a series of isoelectronic complexes  $\text{LM}(\text{CO})_3$ , estimated by the comparison of  $\nu_{\text{CO}}$  in Table 2, suggests the  $\pi$ -acidity decreases in the order  $\text{C}_6\text{H}_6$ , 12P3, 9S3,  $\text{C}_2\text{B}_9\text{H}_{11}^{2-}$ ,  $\text{C}_5\text{H}_5^-$ ,  $\text{HBPz}_3^-$ , and 9N3. With this trend in mind, it is worth noting that in case of  $\text{L}=\text{C}_2\text{B}_9\text{H}_{11}^{2-}$ <sup>24</sup> and  $\text{C}_5\text{H}_5^-$ <sup>26</sup> oxidative decarbonylation reactions have been observed. Further studies on the reactivities of three 9S3 complexes and the development of heteroleptic crown thioether chemistry are under investigation.

**Acknowledgement.** This work was in part supported by the Korea Science and Engineering Foundation. YD thanks Yun Am Cultural Foundation.

### References

1. S. R. Cooper, *Acc. Chem. Res.*, **21**, 141 (1988).
2. M. Schröder, *Pure & Appl. Chem.*, **60**, 517 (1988).
3. S. R. Cooper and S. C. Rawle, *Struc. & Bonding*, **72**, 1 (1990).
4. One of 9S3 in dinuclear cation,  $[\text{Cu}_2(9\text{S3})_3]^{2+}$ , has considerably greater conformational flexibility: J. A. Clarkson, R. Yagbasan, P. J. Blower, and S. R. Cooper, *J. Chem. Soc., Chem. Commun.*, 1244 (1989).
5. The first 9S3 complex of  $d^0$  Re (VII) ion: H. -J. Küppers, B. Nuber, J. Weiss, and S. R. Cooper, *J. Chem. Soc., Chem. Commun.*, 979 (1990).
6. Deprotonation and ring-opening of coordinated 9S3 have been observed in the presence of base: A. J. Blake, A. J. Holder, T. H. Hyde, H. -J. Küppers, M. Schröder, S. Stotzel, and K. Wieghardt, *J. Chem. Soc., Chem. Commun.*, 1600 (1989).
7. (a) A. J. Blake, R. D. Crofts, G. Reid, and M. Schröder, *J. Organomet. Chem.*, **359**, 371 (1989); (b) A. J. Blake, R. O. Gould, A. J. Hoder, T. I. Hyde, G. Reid, and M. Schröder, *J. Chem. Soc., Dalton Trans.*, 1759 (1990).
8. B. L. Ross, J. L. Grasselli, W. M. Ritchey, and H. D. Kaesz, *Inorg. Chem.*, **2**, 1023 (1963).
9. P. J. Blower and S. R. Cooper, *Inorg. Chem.*, **26**, 2009 (1987).
10. (a) Anal. data for **1** ( $\text{C}_9\text{H}_{12}\text{O}_3\text{S}_3\text{Cr}$ ). Calcd (found): C, 34.2 (34.2); H, 3.82 (3.83); (b) Anal. data for **3** ( $\text{C}_9\text{H}_{12}\text{O}_3\text{S}_3\text{W}$ ). Calcd (found): C, 24.1 (24.1); H, 2.70 (2.66); (c) Analysis of **2** was omitted since its synthesis<sup>11,12</sup> and structure<sup>11</sup> have been previously reported.
11. M. T. Ashby and D. L. Lichtenberger, *Inorg. Chem.*, **24**, 636 (1985).
12. (a) D. Sellmann and L. Zapf, *Angew. Chem. Int. Ed. Engl.*, **23**, 807 (1984); (b) D. Sellmann and L. Zapf, *J. Organomet. Chem.*, **57**, 289 (1985).
13. Suitable crystals of **3** were grown by diffusing  $\text{Et}_2\text{O}$  into a DMF solution. Crystallographic data for **3**:  $\text{C}_9\text{H}_{12}\text{O}_3\text{S}_3\text{W}$ , monoclinic, space group  $P2_1/n$ ,  $a=9.134(1)$ ,  $b=11.060(2)$ ,  $c=12.619(2)$  Å,  $\beta=92.74(1)^\circ$ ,  $V=1273.3(3)$  Å<sup>3</sup>,  $R(R_w)=0.0310$  (0.0335). A parallelepiped yellow crystal ( $0.21 \times 0.25 \times 0.41$  mm) was sealed in a glass capillary and mounted on an Enraf-Nonius CAD4 diffractometer. Diffraction data were collected at ambient temperature using  $\text{MoK}_\alpha$  radiation to a maximum  $2\theta$  value of  $50^\circ$  with  $w$ - $2\theta$  scan mode. Data were corrected for Lorentz, polarization and absorption effects. Intensity data for 1791 reflections with  $I > 3\sigma(I)$  were used for Patterson method and SHELX 76.
14. A. G. Orpen, L. Brammer, F. H. Allen, O. Kennerd, D. G. Watson, and R. Taylor, *J. Chem. Soc., Dalton Trans.*, S1 (1989).
15. The 0.01 Å shortening of the average M-S distance in **3** can be counted as significant because the covalent radius of W is longer than Mo by 0.01 Å.<sup>16</sup>
16. The covalent radii for Mo(0) and W(0) are 1.45 and 1.46 Å, respectively: I. S. Butler and J. F. Harrod, *Inorganic Chemistry-Principles and Applications*, Benjamin/Cummings Pub., p 48 (1989).
17. Chemical shifts (ppm) of free 9S3: <sup>1</sup>H-NMR ( $\text{CDCl}_3/d_6$ -DMSO), 3.13/3.07(s); <sup>13</sup>C-<sup>1</sup>H-NMR ( $\text{CD}_3\text{CN}$ ), 35.4.
18. A ligand commutation involving 12S4 has been reported: E. W. Abel, P. D. Beer, I. Moss, K. G. Orrell, V. Sik, P. A. Bates, and M. B. Hursthouse, *J. Chem. Soc., Chem. Commun.*, 978 (1987); *J. Organomet. Chem.*, **341**, 559 (1988).
19. At our hands, only starting complexes have been obtained from the attempted oxidative decarbonylation of  $[\text{M}(\text{CO})_3(9\text{S3})]$  with elemental sulfur at room temperature. Simultaneous use of  $\text{Me}_3\text{NO}$  proved to be also ineffective.
20. P. Chaudhuri, K. Wieghardt, Y. -H. Tsai, and C. Krüger, *Inorg. Chem.*, **23**, 427 (1984).
21. C. Barbeau and J. Turcotte, *Can. J. Chem.*, **54**, 1612 (1976).
22. D. E. Crotty, E. R. Corey, T. J. Anderson, M. D. Glick, and J. P. Oliver, *Inorg. Chem.*, **16**, 920 (1977).
23. S. Trofimenko, *J. Am. Chem. Soc.*, **91**, 588 (1969).
24. Y. W. Kim, M. S. Thesis, KAIST (1989).
25. B. N. Diel, R. C. Haltiwanger, and A. D. Norman, *J. Am. Chem. Soc.*, **104**, 4700 (1982).
26. S. C. O'Neal, W. T. Pennington, and J. W. Kolis, *Organometallics*, **8**, 2281 (1989).

### New Synthesis of Acylferrocene from Ferrocenecarboxaldehyde by Rh(I) Catalyst

Chul-Ho Jun\*, Jung-Bu Kang, and Jin-Yong Kim

Agency for Defense Development, Taejeon 300-600

Received February 21, 1991

The C-H bond activation by transition metal complexes has been one of the recent interests in organometallic chemistry.<sup>1</sup> It has been reported that C-H bond activation of the aldimine by Rh(I) generated iminoacylrhodium(III) hydride complex which inserted mono-olefin and conjugate dienes to form iminoacylrhodium(III) alkyl<sup>2</sup> and alkyl-substituted  $\eta^3$ -allyl complexes,<sup>3</sup> respectively. They were readily reductive-eliminated to give corresponding ketimines, potential precursors for ketone since hydrolysis of them produce ketones. Consequently aldehyde can be easily converted to ketone