

syl- α -D-mannopyranoside (6b) with superoxide was carried out in the same manner as described above. **6b** (0.13 g, 0.29 mmole) produced a mixture (0.033 g) of four compounds, which are not characterized yet; R_f 0.84 for **6b**, 0.66, 0.41 and 0.34 (toluene : ethyl acetate : ethanol, 5 : 5 : 2); $^1\text{H-NMR}$ of compounds having R_f 0.41 and 0.34 indicated hydrolysis had occurred.

5,6-anhydro-3-O-benzyl-1,2-O-isopropylidene- β -L-idofuranose (11) (0.046 g) was obtained from 3-O-benzyl-1,2-O-isopropylidene-5,6-di-O-mesyl- α -D-glucofuranose (**10**) (0.21 g, 0.46 mmole); R_f 0.42 for **10**, 0.59 for **11** (toluene : ethyl acetate, 5 : 3); $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ of **11** were identical to those of the authentic **11** that was synthesized by a known method.¹² (Table 2)

Acknowledgement. This work was supported by Research Center for New Bio-Materials in Agriculture, Seoul National University-KOSEF and Academic Foundation of Soong Sil University.

References

1. J. S. Fillipo, Jr., Cheun-Ing Chern, and J. S. Valentine, *J. Org. Chem.*, **40**, 1678 (1975).
2. E. J. Corey, K. C. Nicolaou, M. Shibasaki, Y. Machida, and C. S. Shiner, *Tetrahedron Lett.*, **37**, 3183 (1975).
3. D. H. Ball and F. W. Parrish, *Advan. Carbohydr. Chem.*, **23**, 233 (1968) and **24**, 139 (1969).
4. L. Hough and A. C. Richardson, *Comprehensive Org. Chem.*, (eds. D. H. R. Barton and W. D. Ollis), Vol. 5, p. 700, Pergamon Press, Oxford (1979).
5. K. Bock and C. Pedersen, *Advan. Carbohydr. Chem. and Biochem.*, **41**, 27 (1983).
6. N. Baggett, *Carbohydr. Chem.*, (ed. J. F. Kennedy) p. 381, Clarendon Press, Oxford (1988).
7. H. H. Baer and D. J. Astles, *Carbohydr. Res.*, **126**, 343 (1984).
8. V. S. Rao and A. S. Perlin, *Can J. Chem.*, **59**, 333 (1981).
9. N. K. Richtmyer, *Methods in Carbohydr. Chem.*, (eds. R. L. Whistler and M. L. Wolfrom) Vol. 1, p. 107, Academic Press (1962).
10. G. W. J. Fleet, N. M. Carpenter, S. Petursson, and N. G. Ramsden, *Tetrahedron Lett.*, **31**(3), 409 (1990).
11. J. E. Nam Shin and A. S. Perlin, *Carbohydr. Res.*, **84**, 315 (1980).
12. R. L. Whistler and W. C. Lake, *Methods in Carbohydr. Chem.*, (eds. R. L. Whistler and J. N. BeMiller) Vol. 6, p. 286, Academic Press (1972).
13. J. E. Nam Shin, *Soong Sil Univ., Papers and Essays, Part III. Natural Sciences*, Vol. 13, 331 (1983).
14. Ed. P. M. Collins, *Carbohydrates*, p. 362, Chapman and Hall Chemistry Sourcebooks (1987).
15. L. F. Wiggins, *Methods in Carbohydr. Chem.*, (eds. R. L. Whistler and M. L. Wolfrom) Vol. 2, p. 188, Academic Press (1963).

Bonding of Electron Deficient Thallium-Metal Cluster Compound

Sungkwon Kang

Department of Chemistry, College of Natural Sciences, Chungnam National University,
Taejon 305-764. Received September 8, 1992


Molecular orbital calculations at the extended Hückel level have been carried out for an electron deficient cluster, $\text{Ti}_3(\text{FeL}_3)_2(\text{FeL}_4)_3^{-3}$, where $\text{L}=\text{CO}$ or H^- . The LUMO, $2a_2''$, is destabilized by the secondary interaction of the LUMO with $1a_2''$ on $(\text{FeL}_3)_2$ fragment. This is one of six skeletal bonding orbitals which are associated with Tl- FeL_3 bonds. Overlap population analysis has been applied to account for two kinds of Tl-Fe bonds. Replacement of the terminal C_{3v} FeL_4 by the C_{2v} FeL_4 units in cluster results in slight energy stabilization of the cluster.

Introduction

A large number of clusters with both main group atoms and transition metals have been prepared and structurally determined.^{1,2} It is possible to predict the cage geometries of these cluster compounds through a set of electron counting rules.^{2,3} The chemical bonding of the cluster compound of $[\text{Et}_4\text{N}]_6[\text{Ti}_6\text{Fe}_{10}(\text{CO})_{36}]$ (**1**), however, has not been clearly understood yet. From our previous X-ray diffraction study⁴, the inter-Thallium distances were 3.71-3.77 Å, which are in the range of very weak Tl-Tl interactions.⁵ In the preliminary calculations the Tl-Tl overlap population was only 0.025.⁴ Tl^I-Tl^I interaction for the various molecular and solid-state structures was studied by C. Janiak and R. Hoffmann.⁶

In this paper we discuss the analysis of molecular orbital

calculations of compound **1** at the semiempirical extended Hückel level. The simple approach is to consider a dimer of a model system $\text{Ti}_3(\text{FeL}_3)_2(\text{FeL}_4)_3^{-3}$ (**2**), where $\text{L}=\text{Two}$ electron donor of CO or H^- .






Figure 1. The energy levels of equilateral triangle C_{3v} $3Tl^{+1}$ fragment.

The Fragment Orbitals

To investigate the electronic structures of the monomer cluster compound, we divided it into three fragments, Tl_3^{+1} (3), $2FeL_3^{-2}$ (4), and $3FeL_4$ (5). The fragment of Tl_3^{+1} with symmetry D_{3h} are shown along with the orbital diagrams in Figure 1. Tl_3^{+1} is a hypothetical fragment model in the cluster compound.

From the interaction of the three equivalent Tl orbitals in the Tl_3^{-1} fragments, the energy splitting is not large because of the long inter-Thallium distances (3.71-3.77 Å). Three thallium p orbitals produce two kinds of fragment orbitals. One is a radial orbital which includes the s orbitals and the other comprises two tangential orbitals. In Figure 1, the a_2'' orbital is π -type bonding, while the e'' set is π -type orbitals with out-of-phase combination. The $2e'$ set is the mixtures of tangential and radial p orbitals. The $2a_1'$ orbital is made with an in-phase combination of the radial p orbitals with a small portion of the s orbitals. At higher energy level is an antibonding $3e'$ set of p orbitals mixed with s orbitals.

As shown in Figure 2, fragment 4 ($2FeL_3^{-2}$) comprises two C_{3v} ML_3 units. The valence MO's for the C_{3v} ML_3 system have already been described in a number of papers.^{2,7} These are shown on the far right and left sides in the figure.

The $1a_1$ and $1e$ sets at the low energy levels are pure d orbitals of Fe atom which correspond to the t_{2g} set in an octahedral system. In the $2e$ set the orbitals are primarily metal orbitals of d_{yz} and d_{zx} mixed with some p orbitals which direct toward the missing ligands. The $2a_1$ is a $s-p_z$ hybrid orbital of the metal. When two C_{3v} FeL_3^{-2} fragments are combined together, each fragment orbital interacts in the D_{3h} symmetry. The low-lying two three-degenerate orbitals ($1a_1 + 1e$), which are shown in the left and right sides of the figure, respectively, form six occupied orbitals of $1a_1'$,




Figure 2. Valence orbitals for an $2FeL_3^{-2}$ fragment.

$1e'$, $1a_2''$, and $1e''$ (shown on the center of the figure). All these orbitals are at the low energy levels with doubly occupied. Therefore, these orbitals cannot interact strongly with the Tl_3^{+1} fragment orbitals in the cluster compound. The important valence orbitals in the fragment 4 are obtained from the symmetry adapted linear combinations of the three higher orbitals in FeL_3^{-2} , i.e., the $2a_1$ and $2e$ set. In the middle of the figure, the energy levels of the $2e'$ and $2e''$ orbitals do not split because of long inter-iron distances (3.13 Å) in the cluster compound. The energy levels of the $2a_1'$ and $2a_2''$ orbitals split somewhat because $4p$ orbitals are more diffusive than $3d$ orbitals.

Fragment 5, $3FeL_4$, is a simple combination of three C_{3v} FeL_4 units in a triangle. The valence orbitals of C_{3v} FeL_4 system can be easily derived from a D_{3h} FeL_5 system. Removing one ligand in axial position, a_1' orbital is stabilized considerably by forming the a_1 orbital and the hybridization toward the missing ligand (shown in Figure 3).

Two e sets, e' and e'' , are not perturbed when one axial ligand is removed. The important valence orbitals in the $3FeL_4$ fragment are derived from symmetry adapted linear combinations of the two sets of orbitals in C_{3v} FeL_4 , i.e., the a_1 and $1e$ set. These orbital shapes are shown in top view in 6 and 7.

The $Tl_3(FeL_3)_2(FeL_4)_3^{-3}$ Cluster Compound

Figure 4 represents the interaction diagram for a Tl_3^{-1} fragment interacting with two metal fragments.

On the left side of Figure 4 are the orbitals of Tl_3^{+1} . The




Figure 5. Contour plots of $2a_2''$, $3a_1'$, e'' , and $2e_3'$ orbitals in $Tl_3(FeL_3)_2(FeL_4)_3^{-3}$ cluster. The contours have the values of 0.0125, 0.025, 0.05, 0.075, 0.10, 0.15, and 0.20. The different phases of the wavefunction are represented by full and dashed lines. The orbitals except for $3a_1'$ (B) have been plotted in the yz plane which is defined in Figure 4. The $3a_1'$ (B) orbital has been plotted in xy plane.

FeL_4 fragments.

The computed Tl-Fe (of terminal FeL_4) overlap populations are 0.120 larger than the others of μ_3 - FeL_3 units (for which all Tl-Fe distances were fixed to be equal). This difference is consistent with the X-ray observation. Namely, the average distance of the terminal Tl- FeL_4 is 2.54 Å, while that of the bridging Tl- FeL_3 is 2.67 Å. The Tl-Fe overlap population difference can be explained easily. Two interaction types, σ - and π -bonding (**10** and **11**), are involved in the terminal Tl- FeL_4 bonds.

The filled sp hybrid orbital of Tl atom strongly interacts with the empty s , p_y , and $d_{x^2-y^2}$ hybrid orbital of FeL_4 unit in a given coordinate system, shown in **10**. And the filled d_{xy} and d_{yz} of the terminal FeL_4 unit also interact with empty p_x and p_z of Tl atom, **11**. These are back-bonding interactions. Total overlap populations from these two interactions are





Table 1. Parameters Used in the Extended Hückel Calculations

Atom	Orbital	H_{ii} , eV				
ζ_1	ζ_2	C_1^a	C_2^a			
Fe	3d	-12.70	5.35	1.80	0.5366	0.6678
	4s	-9.17	1.90			
	4p	-5.37	1.90			
Tl	6s	-11.60	2.30			
	6p	-5.80	1.60			
C	2s	-21.40	1.625			
	2p	-11.40	1.625			
O	2s	-32.30	2.275			
	2p	-14.80	2.275			
H	1s	-13.60	1.30			

^a Contraction coefficient used in the double- ζ expansion.

computed to be 0.3453. The main interactions in Tl- FeL_3 are those between the s , p_z orbitals of Tl atoms and the empty s , p orbitals of bridging metal units. The contribution of the metal d orbital for skeletal bonding is only 16% of total overlap population between the Tl atoms and bridging FeL_3 units.

Many cluster compounds have C_{2v} ML_4 fragment instead of C_{3v} ML_4 .⁸ This is one of candidates for the replacement of fragment **5**, $3FeL_4$. The valence orbitals of C_{2v} FeL_4 unit are shown in **12**.



The overlap populations of 0.3285 between the filled sp hybrid of Tl atom and LUMO $2a_1$ in **12** are larger than those of C_{3v} FeL_4 system, 0.3064. The HOMO b_2 in **12** interacts with empty p_x of Tl atom. This is back-bonding interaction which is also stronger than C_{3v} FeL_4 system. Our calculations find that the cluster compound with C_{2v} FeL_4 unit is slightly more stable than the cluster with C_{3v} .

Acknowledgement. This work was supported by NON DIRECTED RESEARCH FUND, Korea Research Foundation, 1990.

Appendix

The calculations have been carried out with the extended Hückel formalism,⁹ using the weighted H_{ij} formalism.¹⁰ The atomic parameters listed in Table 1 were obtained from pre-

vious work.^{6,11} In the molecular model, L is a two electron donor, *i.e.*, CO or H⁻. All Ti-Ti, Ti-Fe, Fe-C, Fe-H, and C-O distances were fixed at 3.75, 2.67, 1.78, 1.60, and 1.14 Å, respectively. The L-Fe-L angles were fixed at 90.0°.

References

- (a) B. F. G. Johnson, "Transition Metal Clusters", Wiley, Chichester, 1980; (b) Ch. Elschenbroich and A. Salzer, "Organometallics", VCH, Weinheim, 1992.
- T. A. Albright, J. K. Burdett, and M.-H. Whangbo, "Orbital Interactions in Chemistry", Wiley, New York, 1985.
- (a) J. K. Burdett, "Molecular shapes", Wiley, New York 1980; (b) K. Wade, *Adv. Inorg. Chem. Radiochem.*, **18**, 1 (1976); (c) D. M. P. Mingos, *Inorg. Chem.*, **24**, 114 (1985); (d) B. K. Teo, *Inorg. Chem.*, **24**, 1627 (1985); (e) W. N. Lipscomb, "In Born Hydride Chemistry", E. L. Muetterties, Ed., Academic Press, New York, pp. 39, 1975.
- K. H. Whitmire, R. R. Ryan, H. J. Wasserman, T. A. Albright, and S. K. Kang, *J. Am. Chem. Soc.*, **108**, 6831 (1986).
- (a) J. Beck and J. Strähle, *Z. Naturforsch.*, **41b**, 1381 (1986); (b) S. Harvey, M. F. Lappert, C. L. Raston, B. W. Skelton, G. Strivastava, and A. H. White, *J. Chem. Soc., Chem. Commun.*, 1216 (1988); (c) K. H. Whitmire, J. M. Cassidy, A. L. Rheingold, and R. R. Ryan, *Inorg. Chem.*, **27**, 1347 (1988); (d) J. M. Cassidy and K. H. Whitmire, *Inorg. Chem.*, **28**, 1432 (1989).
- C. Janiak and R. Hoffmann, *J. Am. Chem. Soc.*, **112**, 5924 (1990).
- (a) T. A. Albright, *Tetrahedron*, **38**, 1339 (1982); (b) K. H. Whitmire, T. A. Albright, S. K. Kang, M. R. Churchill, and J. C. Fettinger, *Inorg. Chem.*, **25**, 2799 (1986); (c) J. A. S. Howell, N. F. Ashford, D. T. Dixon, J. C. Kola, T. A. Albright, and S. K. Kang, *Organometallics*, **10**, 1852 (1991).
- (a) M. R. Churchill, F. J. Hollander, and J. P. Hutchinson, *Inorg. Chem.*, **16**, 2655 (1977); (b) M. R. Churchill and B. G. DeBoer, *Inorg. Chem.*, **16**, 878 (1977); (c) L. F. Dahl and J. F. Blount, *Inorg. Chem.*, **4**, 1965 (1965).
- R. Hoffmann, *J. Chem. Phys.*, **39**, 1397 (1963); R. Hoffmann and W. N. Lipscomb, *J. Chem. Phys.*, **36**, 2179 (1962); **37**, 2872 (1962).
- J. H. Ammeter, H.-B. Bürgi, J. C. Thibeault, and R. Hoffmann, *J. Am. Chem. Soc.*, **100**, 3686 (1978).
- T. A. Albright, P. Hofmann, and R. Hoffmann, *J. Am. Chem. Soc.*, **99**, 7546 (1977).

Dioxygen Binding to Dirhodium(II, II), (II, III), and (III, III) Complexes. Spectroscopic Characterization of $[\text{Rh}_2(\text{ap})_4(\text{O}_2)]^+$, $\text{Rh}_2(\text{ap})_4(\text{O}_2)$, and $[\text{Rh}_2(\text{ap})_4(\text{O}_2)]^-$, where ap=2-anilinopyridinate Ion

Jae-Duck Lee*, Chao-Liang Yao, Françoise J. Capdevielle, Baocheng Han, John L. Bear[†], and Karl M. Kadish[†]

Department of Chemistry, University of Houston, Houston, Texas 77204-5641, U.S.A.

Received September 9, 1992

The neutral, reduced, and oxidized 2,2-trans isomers of $\text{Rh}_2(\text{ap})_4$ (ap=2-anilinopyridinate) were investigated with respect to dioxygen binding in CH_2Cl_2 containing 0.1 M tetrabutyl-ammonium perchlorate. $\text{Rh}_2(\text{ap})_4$ binds dioxygen in nonaqueous media and forms a $\text{Rh}^{\text{II}}\text{Rh}^{\text{III}}$ superoxide complex, $\text{Rh}_2(\text{ap})_4(\text{O}_2)$. This neutral species was isolated and is characterized by UV-visible and IR spectroscopy, mass spectrometry and cyclic voltammetry. It can be reduced by one electron at $E_{1/2} = -0.45$ V *vs.* SCE in CH_2Cl_2 and gives $[\text{Rh}_2(\text{ap})_4(\text{O}_2)]^-$ as demonstrated by the ESR spectrum of a frozen solution taken after controlled potential reduction. The superoxide ion in $[\text{Rh}_2(\text{ap})_4(\text{O}_2)]^-$ is axially bound to one of the two rhodium ions, both of which are in a +2 oxidation state. $\text{Rh}_2(\text{ap})_4(\text{O}_2)$ can also be stepwise oxidized in two one-electron transfer steps at $E_{1/2} = 0.21$ V and 0.85 V *vs.* SCE in CH_2Cl_2 and gives $[\text{Rh}_2(\text{ap})_4(\text{O}_2)]^+$ followed by $[\text{Rh}_2(\text{ap})_4(\text{O}_2)]^{2+}$. ESR spectra demonstrate that the singly oxidized complex is best described as $[\text{Rh}^{\text{II}}\text{Rh}^{\text{III}}(\text{ap})_4(\text{O}_2)]^+$ where the odd electron is delocalized on both of the two rhodium ions and the axial ligand is molecular oxygen.

Introduction

$\text{Rh}_2(\text{ap})_4$ (where ap=2-anilinopyridinate) can exist in four different isomeric forms, two of which have been isolated and structurally characterized.^{1,2} The 4,0 isomer has one rhodium ion bound by four anilino nitrogens and the other by four pyridyl nitrogens. In contrast, each rhodium atom of the $\text{Rh}_2(\text{ap})_4$ 2,2-trans isomer is bound in a *trans* fashion

by two pyridyl nitrogens and two anilino nitrogens of the bridging ligands.

Our laboratory recently reported electrochemical properties of the 4,0 isomer in the presence of O_2 and demonstrated the formation of $\text{Rh}^{\text{II}}\text{Rh}^{\text{III}}(\text{ap})_4(\text{O}_2)$, where the dioxygen ligand was bound to the rhodium ion which was complexed by the four pyridyl nitrogens.³ The 2,2-trans isomer of Rh_2 , whose structure is shown in Figure 1, will also irreversibly bind dioxygen and gives a stable $\text{Rh}_2(\text{ap})_4(\text{O}_2)$ complex in CH_2Cl_2 or THF solutions under O_2 . This complex was isola-

*On leave of absence from Dong-A University, Korea.