

2. Ahuja, S. in *Chiral Separations by Liquid Chromatography*; Ahuja, S., Ed.; ACS Symposium Series 471, American Chemical Society: Washington, DC 1991, chap. 1.
3. Perrin, S. R.; Pirkle, W. H. in *Chiral Separations by Liquid Chromatography*; Ahuja, S., Ed.; ACS Symposium Series 471, American Chemical Society: Washington, DC 1991, chap. 3.
4. Taylor, D. R.; Maher, K. *J. Chromatogr. Sci.* 1992, 30, 67.
5. (a) Pirkle, W. H.; Finn, J. M.; Schreiner, J. L.; Hamper, B. C. *J. Am. Chem. Soc.* 1981, 103, 3064; (b) Hyun, M. H.; Kim, M. H. *J. Liq. Chromatogr.* 1990, 13, 3229.
6. (a) Pirkle, W. H.; Hyun, M. H.; Bank, B. *J. Chromatogr.* 1984, 316, 585; (b) Pirkle, W. H.; Pochapsky, T. *J. Am. Chem. Soc.* 1986, 108, 352; (c) Hyun, M. H.; Park, Y.-W.; Baik, I.-K. *Tetrahedron Lett.* 1988, 29, 4735; (d) Hyun, M. H.; Baik, I.-K.; Pirkle, W. H. *J. Liq. Chromatogr.* 1988, 11, 1249.
7. (a) Hyun, M. H.; Pirkle, W. H. *J. Chromatogr.* 1987, 393, 357; (b) Oi, N.; Kitahara, H.; Mathumoto, Y.; Nakajima, H.; Horikawa, Y. *J. Chromatogr.* 1989, 462, 382; (c) Tambe, A.; Siret, L.; Caude, M.; Begos, A.; Rosset, R. *Chirality* 1990, 2, 106.
8. Recently, a commercialized CSP containing both π -basic and -acidic functional groups has been reported. Pirkle, W. H.; Welch, C. J. *J. Liq. Chromatogr.* 1992, 15, 1947.
9. Doyle, T. D.; Brunner, A.; Smith, E. *US Patent 4,919,803*, Apr. 24, 1990.
10. Pirkle, W. H.; Welch, C. J.; Lamm, B. *J. Org. Chem.* 1992, 57, 3854.
11. (a) Pirkle, W. H.; Hyun, M. H. *J. Org. Chem.* 1984, 49, 3043; (b) Pirkle, W. H.; Hyun, M. H.; Bank, B. *J. Chromatogr.* 1984, 316, 585; (c) Hyun, M. H.; Kim, M. S.; Ryoo, J.-J. *Bull. Kor. Chem. Soc.* 1993, 14, 9.
12. Pirkle, W. H.; Welch, C. J. *J. Liq. Chromatogr.* 1991, 14, 1.
13. Hyun, M. H.; Cho, S. M.; Ryoo, J.-J.; Kim, M. S. *J. Liq. Chromatogr.* 1994, 17, 317.

Kinetic Studies on the Aminolysis of 2-Phenyl-1-propyl Arenesulfonates in Methanol

Han Joong Koh, Igor V. Shpan'ko[†], and Ikchoon Lee*

Department of Chemistry, Inha University, Incheon 402-751

[†]*Department of Chemistry, Donetsk State University, Donetsk, Ukraine*

Received March 2, 1994


The results of kinetic studies on the reactions of 2-phenyl-1-propyl arenesulfonates with anilines and benzylamines in methanol at 55.0°C are reported. The transition state variation with the substituents in the nucleophile (X) and leaving group (Z) is in accord with that expected from a negative ρ_{XZ} value: A stronger nucleophile and nucleofuge lead to a greater extent of bond-making and -breaking. Somewhat greater magnitude of ρ_{XZ} compared to the nearly constant value for the similar processes at a primary carbon atom has been interpreted to result from a partial contribution of the concurrent frontal displacement path.

Introduction

The cross-interaction constants, ρ_{ij} in Eq. (1) where i and j represent substituents in the nucleophile (X), substrate (Y) or leaving group (Z) in Scheme 1, are useful as a mechanistic tool for organic reactions in solution.¹

$$\log(k_{ij}/k_{HH}) = \rho_i\sigma_i + \rho_j\sigma_j + \rho_{ij}\sigma_i\sigma_j \quad (1)$$

We have shown for nucleophilic substitution reactions that: (i) The magnitudes of ρ_{XY} and ρ_{YZ} are directly proportional to the extent of bond-making and -breaking respectively in the transition state (TS), provided the fall-off of $|\rho_{ij}|$ (by ca. 2.8) due to an intervening nonconjugative group, e.g. CH₂ or CO, between the substituent and the reaction center is accounted for.² (ii) A positive [negative] ρ_{XZ} , which can be alternatively given as Eq. (2), leads to an earlier [later] TS along the



Scheme 1.

$$\rho_{XZ} = \frac{\partial \rho_Z}{\partial \sigma_X} = \frac{\partial \rho_X}{\partial \sigma_Z} \quad (2)$$

reaction coordinate for a stronger nucleophile ($\delta\sigma_X < 0$) and/or a stronger nucleofuge, i.e., a better leaving group ($\delta\sigma_Z > 0$).¹ (iii) The magnitude of ρ_{XZ} , is a measure of the TS tightness; the greater the $|\rho_{XZ}|$, the tighter is the TS (the shorter is the d^*_{XZ} in Scheme 1).^{1,3}

Table 3. Cross-interaction Constants, ρ_{XZ} and β_{XZ} , for Some Nucleophilic Substitution Reactions

Reaction	Solvent	Temp. °C	ρ_{XZ}	β_{XZ}
1. $\text{XC}_6\text{H}_4\text{NH}_2 + \text{CH}_3\text{CH}_2\text{OSO}_2\text{C}_6\text{H}_4\text{Z}^8$	MeOH	65	0.33	0.19
	MeCN	65	0.34	0.21
2. $\text{XC}_6\text{H}_4\text{NH}_2 + (\text{CH}_3)_3\text{SiCH}_2\text{OSO}_2\text{C}_6\text{H}_4\text{Z}^{12}$	MeOH	65	0.31	0.18
	MeCN	65	0.33	0.20
$\text{XC}_6\text{H}_4\text{CH}_2\text{NH}_2 + (\text{CH}_3)_3\text{SiCH}_2\text{OSO}_2\text{C}_6\text{H}_4\text{Z}^{12}$	MeCN	65	0.08	0.12
3. $\text{XC}_6\text{H}_4\text{NH}_2 + (\text{CH}_3)_3\text{CCH}_2\text{OSO}_2\text{C}_6\text{H}_4\text{Z}^{13}$	MeOH	55	0.31	0.18
$\text{XC}_6\text{H}_4\text{CH}_2\text{NH}_2 + (\text{CH}_3)_3\text{CCH}_2\text{OSO}_2\text{C}_6\text{H}_4\text{Z}^{13}$	MeOH	55	0.11	0.15
4. $\text{XC}_6\text{H}_4\text{NH}_2 + \text{C}_6\text{H}_5\text{CH}(\text{CH}_3)\text{OSO}_2\text{C}_6\text{H}_4\text{Z}^{14}$	MeOH	25	-0.56	-0.32
5. $\text{XC}_6\text{H}_4\text{NH}_2 + \text{C}(\text{CH}_3)_3\text{CHC}_6\text{H}_5\text{OSO}_2\text{C}_6\text{H}_4\text{Z}^{15}$	MeOH	35	0.0	0.0
6. $\text{XC}_6\text{H}_4\text{NH}_2 + \text{C}_6\text{H}_5\text{CH}_2\text{OSO}_2\text{C}_6\text{H}_4\text{Z}^{16}$	MeOH	30	-0.10	-0.06
7. $\text{XC}_6\text{H}_4\text{NH}_2 + \text{C}_6\text{H}_5\text{CH}_2\text{CH}_2\text{OSO}_2\text{C}_6\text{H}_4\text{Z}^4$	MeOH	65	-0.45	-0.28
8. $\text{XC}_6\text{H}_4\text{CH}_2\text{NH}_2 + \text{C}_6\text{H}_5\text{CH}_2\text{CH}_2\text{OSO}_2\text{C}_6\text{H}_4\text{Z}^{17}$	MeOH	65	-0.13	-0.21
9. $\text{XC}_6\text{H}_4\text{NH}_2 + \text{C}_6\text{H}_5\text{CH}_2\text{CH}(\text{CH}_3)\text{SO}_2\text{C}_6\text{H}_4\text{Z}^5$	MeOH	65	0.16	0.02
10. $\text{XC}_6\text{H}_4\text{NH}_2 + \text{C}_6\text{H}_5\text{CH}(\text{CH}_3)\text{CH}_2\text{OSO}_2\text{C}_6\text{H}_4\text{Z}^a$	MeOH	55	-0.39(0.999) ^b	-0.23(0.999)
$\text{XC}_6\text{H}_4\text{CH}_2\text{NH}_2 + \text{C}_6\text{H}_5\text{CH}(\text{CH}_3)\text{CH}_2\text{OSO}_2\text{C}_6\text{H}_4\text{Z}^a$	MeOH	55	-0.29(0.996)	-0.42(0.995)

^aThis work; ^bThe values in parenthesis are the correlation coefficients at 99% confidence level.

Table 4. The Kinetic Isotope Effects (k_H/k_D) for the Reaction of 2-Phenyl-1-propyl Z-arenesulfonates with Deuterated X-anilines ($\text{XC}_6\text{H}_4\text{ND}_2$) in Deuterated Methanol (CH_3OD) at 55.0°C

X	Z	MeOH ($\text{M}^{-1} \text{s}^{-1}$)	MeOD ($\text{M}^{-1} \text{s}^{-1}$)	k_H/k_D
<i>p</i> -CH ₃ O	<i>p</i> -NO ₂	79.2 (± 1) ^a $\times 10^{-6}$	88.6 (± 2) $\times 10^{-6}$	0.894 \pm 0.007 ^b
<i>p</i> -CH ₃ O	<i>p</i> -CH ₃	15.8 (± 0.6) $\times 10^{-6}$	17.5 (± 0.2) $\times 10^{-6}$	0.901 \pm 0.008
<i>p</i> -Cl	<i>p</i> -NO ₂	9.77 (± 0.5) $\times 10^{-6}$	10.5 (± 0.7) $\times 10^{-6}$	0.924 \pm 0.010
<i>p</i> -Cl	<i>p</i> -CH ₃	2.75 (± 0.4) $\times 10^{-6}$	2.94 (± 0.5) $\times 10^{-6}$	0.935 \pm 0.007


^aStandard deviation, ^bStandard error.¹⁸

reflecting a greater degree of steric crowding due to a greater degree of bond formation.

It is interesting to note in Table 3 that the reaction type with a negative ρ_{XZ} value is rather rare, and occurs only in the $\text{S}_{\text{N}}2$ processes at a benzylic and primary 2-phenylethyl carbon centers. The magnitude of β_Z in Table 2, is relatively smaller, especially for the reactions with benzylamines, than those for the normal $\text{S}_{\text{N}}2$ processes, $\beta_Z \cong -0.30 \sim -0.45$,⁸ suggesting a relatively low degree of bond cleavage in the TS. This is in line with a relatively greater magnitude of ρ_{XZ} and β_{XZ} for **III** in Table 3. For the reactions with anilines, the magnitude of ρ_{XZ} ($= -0.39$) and β_{XZ} ($= -0.23$) is somewhat greater than those for other $\text{S}_{\text{N}}2$ processes at a primary carbon center ($\rho_{XZ} = 0.31\text{--}0.33$ and $\beta_{XZ} = 0.18\text{--}0.19$). It is to be noted that very little change in $|\rho_{XZ}|$ may be expected for the 10°C rise in temperature.⁹ The magnitude of ρ_{XZ} and β_{XZ} is especially greater for the reactions with benzylamine; the magnitude is more than twice of that for other reactions at a primary carbon center, reactions 2, 3, and 8 in Table 3. Since the magnitude of ρ_{XZ} (and β_{XZ}) is a measure of the TS tightness, these larger values obtained in this work seem to reflect a somewhat tighter TS for the reactions studied in this work, Eq. (3). In view of our earlier results, however, the following alternative interpretation seems more plausible.

The structure of the 2-phenylethyl arenesulfonate analogue, **IV**, suggests that the normal rear-side $\text{S}_{\text{N}}2$ attack by

an aniline molecule must be sterically inhibited due to the relatively large size of the (Y) benzene ring (e.g. the Taft steric constants¹⁰ are -0.07 and -0.38 for CH_3 and C_6H_5) lending probability to a front-side attack. In the front-side $\text{S}_{\text{N}}2$ attack the nucleophile and leaving group can come close together leading to a greater magnitude of ρ_{XZ} (and β_{XZ}). Indeed such a frontal attack has been predicted for the reactions of **I** based on the large magnitude of ρ_{XZ} (and β_{XZ}) observed.⁴



- I**, $R_1 = R_2 = \text{H}$
II, $R_1 = \text{H}, R_2 = \text{CH}_3$
III, $R_1 = \text{CH}_3, R_2 = \text{H}$

IV

For the 1-methyl analogue, **III**, however, the frontal attack should be somewhat limited by an extra methyl group on C_β . In any of the reactions of **I-III** it is hard to conceive of the total front-side attack process, since the normal rear-side attack process is intrinsically more favorable and can compete with the frontal process.¹¹ Therefore, the relatively

greater magnitude of ρ_{XZ} (β_{XZ}) obtained for I-III appears most likely to result from partial involvement of the frontal process. The extent of such contribution may decrease in the order I>II>III, considering an additional steric inhibition by the methyl group on C_α and C_β in II and III respectively. The increment of the magnitude, $\Delta\rho_{XZ} = \rho_{XZ(obsd)} - \rho_{XZ(normal)}$, is 0.14, 0.06 and 0.05 for the reactions of I, II and III, respectively, assuming the normal $|\rho_{XZ}|$ value in MeOH at 65.0°C of 0.31⁶ and 0.10⁷ for the reactions at a primary and secondary carbon atom. The decrease in the increment, $\Delta\rho_{XZ}$, should reflect the decrease in the extent of the frontal process contribution.

The greater increase in ρ_{XZ} and β_{XZ} for the reactions of III with benzylamines may arise from a greater contribution of the frontal process due to a greater degree of steric inhibition in the rear-side attack by the relatively larger nucleophile, benzylamine.

We therefore conclude that in the S_N2 reactions of 2-phenylethyl arenesulfonates, I-III, with anilines and benzylamines the frontal attack process contribute partially. The contribution of the frontal process results in a greater magnitude of ρ_{XZ} and β_{XZ} , and the contribution is greater, the greater is the steric inhibition in the rear-side attack, and the lower is the steric requirement in the front-side attack.

Experimental

Materials. Merck analytical grade methanol was used without further purification. The nucleophiles, aniline, were Aldrich G.R. purchased, which were redistilled or recrystallized before use. Preparation of deuterated anilines were as described previously.^{3,6} The analysis (NMR spectroscopy) of the deuterated anilines showed more than 99% deuterium content, so no corrections to kinetic isotope effects for incomplete deuterium were made.

Substrates, 2-phenyl-1-propyl arenesulfonates, were prepared by reacting Aldrich G.R. 1-phenyl-2-propanol with benzenesulfonyl chlorides.¹⁹ The substrates synthesized were confirmed by spectral and elementary analyses as follows.

$C_6H_5CH(CH_3)CH_2OSO_2C_6H_5$. ν_{max} (neat) 1360, 1195 (SO₂), 1020 (SO) and 815 cm⁻¹ (S-O-C); δ_H (CDCl₃) 1.2 (β -CH₃, 3H, d), 2.9 (CH, 1H, m), 4.1 (-CH₂-, 2H, d), 7.2-7.4 (phenyl, 10H, m). (Found: C, 73.7; H, 6.7. Calc. for C₁₅H₁₆O₃S: C, 73.8, H, 6.8%).

$C_6H_5CH(CH_3)CH_2OSO_2C_6H_4-p-CH_3$. mp. 45°C, ν_{max} (KBr) 1350, 1165 (SO₂), 1020 (SO) and 810 cm⁻¹ (S-O-C); δ_H (CDCl₃) 1.3 (β -CH₃, 3H, d), 1.9 (*p*-CH₃, 3H, s), 2.8 (CH, 1H, m), 4.0 (-CH₂-, 2H, d), 7.2-7.8 (phenyl, 9H, m). (Found: C, 74.2; H, 7.1. Calc. for C₁₆H₁₈O₃S: C, 74.4; H, 7.0%).

$C_6H_5CH(CH_3)CH_2OSO_2C_6H_4-p-Cl$. mp. 74-75°C, ν_{max} (KBr) 1355, 1180 (SO₂), 1010 (SO) and 810 cm⁻¹ (S-O-C); δ_H (CDCl₃) 1.3 (β -CH₃, 3H, d), 2.9 (CH, 1H, m), 4.2 (-CH₂-, 2H, d), 7.2-7.8 (phenyl, 9H, m). (Found: C, 74.2; H, 6.1. Calc. for C₁₅H₁₅O₃S: C, 74.1; H, 6.2%).

$C_6H_5CH(CH_3)CH_2OSO_2C_6H_4-p-NO_2$. mp. 108°C, ν_{max} (KBr) 1530, 1345 (NO₂), 1180 (SO₂) and 812 cm⁻¹ (S-O-C); δ_H (CDCl₃) 1.3 (β -CH₃, 3H, d), 2.8 (CH, 1H, m), 4.2 (-CH₂-, 2H, d), 7.2-7.9 (phenyl, 9H, m). (Found: C, 62.4; H, 5.1. Calc. for C₁₅H₁₅O₅NS: C, 62.3; H, 5.2%).

Kinetic procedures. The rate measurements for the reactions of 2-phenyl-1-propyl arenesulfonates with anilines

were attempted in acetonitrile but the rates were found to be too slow for our conductivity apparatus. Rates were measured conductimetrically at 55.0±0.05°C in methanol. The conductivity bridge used in this work was a self-made computer interface automatic A/D converter conductivity bridge. Substrates were injected with a syringe. Pseudo-first order rate constants, k_{obs} , were determined by the Guggenheim method²⁰ with a large excess of aniline; [2-phenyl-1-propyl arenesulfonate]=10⁻³ mol dm⁻³ and [aniline]=0.025-0.50 mol dm⁻³. Second-order rate constant, k_2 , were obtained from the slope of the plot of k_{obs} vs. [aniline] with more than four concentrations of aniline. The intercept of this plot was almost zero ranging from 1.6×10⁻⁷ to 7.1×10⁻¹¹ s⁻¹ indicating that many side reactions including methanolysis are insignificant during the rate measurements.

Product analysis. 2-phenyl-1-propyl arenesulfonate was reacted with excess aniline with stirring for more than 15 half-lives at 55.0°C in methanol, and the products were isolated by evaporating the solvent under reduced pressure. The TLC analysis of the product mixture gave three spots (silica gel, glass plate, 10% ethyl acetate/*n*-hexane).

R_f values. 0.62 (C₆H₅CH(CH₃)CH₂NHC₆H₅), 0.43 (C₆H₅-CH(CH₃)CH₂OSO₂C₆H₅), 0.30 (C₆H₅NH₂), 0.02 (C₆H₅NH₃⁺-OSO₂C₆H₅). The product mixture was treated with column chromatography (silica gel, 10% ethyl acetate/*n*-hexane). Analysis of the products, C₆H₅CH(CH₃)CH₂NHC₆H₅: δ_H (CDCl₃) 1.2 (β -CH₃, 3H, d), 2.7 (CH, 1H, m), 3.0 (NH, 1H, broad), 4.2 (-CH₂-, 2H, d) and 7.2-7.6 (phenyl, 10H, m). C₆H₅OSO₂-NH₃⁺-C₆H₅: δ_H (CDCl₃) 4.8 (NH₃⁺, 3H, broad) and 7.1-7.7 (phenyl, 10H, m). The analysis of the reaction mixture, 2-phenyl-2-propyl arenesulfonate with aniline, by NMR at appropriate intervals for at least 15 half-lives under exactly the same reaction conditions did not give any detectable change due to aryl participation.

Acknowledgement. We thank the Inha University for support of this work. H. J. Koh also thanks the Inha University for postdoctoral fellowship.

References

- (a) Lee, I. *Chem. Soc. Rev.* **1990**, *19*, 317; (b) Lee, I. *Adv. Phys. Org. Chem.* **1992**, *27*, 57.
- Lee, I. *J. Phys. Org. Chem.* **1992**, *5*, 736.
- Koh, H. J.; Lee, H. W.; Lee, I. *J. Chem. Soc. Perkin Trans. 2* **1994**, *2*, 125.
- Lee, I.; Choi, Y. H.; Lee, H. W.; Lee, B. C. *J. Chem. Soc. Perkin Trans. 2* **1988**, *2*, 1539.
- Lee, I.; Lee, W. H.; Lee, H. W. *J. Org. Chem.* **1991**, *56*, 4682.
- Koh, H. J.; Lee, H. W.; Lee, I. *J. Chem. Soc. Perkin Trans. 2* **1994**, *2*, 253.
- Oh, H. K.; Kwon, Y. B.; Cho, I. H.; Lee, I. *J. Chem. Soc. Perkin Trans. 2*, in press.
- Lee, I.; Choi, Y. H.; Rhyu, K. W.; Shim, C. S. *J. Chem. Soc. Perkin Trans. 2* **1989**, *2*, 1881.
- Lee, J. H. Ph. D. Thesis, Han Yang University, Seoul, **1992**.
- Taft, R. W. In *Steric Effects in Organic Chemistry*; Newman, M. ed.; Wiley: New York, 1956, Chapter 13.
- (a) Lowry, T. H.; Richardson, K. S. *Mechanism and Theory in Organic Chemistry*, 3rd ed.; Harper and Row:

- New York, 1987; p 331-333; (b) Yang, K.; Koo, I. S.; Kang, D. H.; Lee, I. *J. Phys. Org. Chem.* Submitted for publication.
12. Oh, H. K.; Shin, C. H.; Lee, I. *J. Chem. Soc. Perkin Trans. 2* 1993, 2, 2411.
 13. Oh, H. K.; Kwon, Y. B.; Lee, I. *J. Phys. Org. Chem.* 1993, 6, 357.
 14. Lee, I.; Kim, H. Y.; Kang, H. K.; Lee, H. W. *J. Org. Chem.* 1988, 53, 2678.
 15. Lee, I.; Choi, M. S.; Lee, H. W. *J. Chem. Res.* 1994, (S)92-93, (M)0568-0587.
 16. Lee, I.; Sohn, S. C.; Oh, Y. J.; Lee, B.-S. *Tetrahedron.* 1986, 42, 4713.
 17. Lee, I.; Lee, W. H.; Lee, H. W. *J. Phys. Org. Chem.* 1990, 3, 545.
 18. Crumpler, T. B.; You, J. H. *Chemical Computations and Errors*; John Wiley: New York, 1940; p 178.
 19. Tipson, R. S. *J. Org. Chem.* 1944, 9, 235.
 20. Guggenheim, E. A. *Philos. Mag.* 1926, 2, 538.